

Quantum Field Theory on a Highly Symmetric Lattice

Marco Aliberti

Università degli Studi di Torino

July 9, 2023

Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated.


Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form

$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated. There are two possible approaches:

Perturbative




Why Lattice Quantum Chromodynamics?

In quantum field theory scattering amplitudes in the form


$$\langle f|i\rangle = \int_{\phi_i}^{\phi_f} \mathcal{D}[\phi] e^{-S[\phi]}$$

need to be evaluated. There are two possible approaches:

Perturbative




Non-Perturbative




Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops

Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated

Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative




- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$

Perturbative vs Non-Perturbative

Perturbative




Non-Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
 - UV divergencies need to be eliminated
 - Fails predicting quantities with essential singularities as $g \rightarrow 0$
- No straightforward approach


Perturbative vs Non-Perturbative

Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$


Non-Perturbative



- No straightforward approach
- Can have a natural cut-off for high momenta \Rightarrow No UV divergencies


Perturbative vs Non-Perturbative

Perturbative



- Straightforward series expansion in powers of small $g \Leftrightarrow$ Feynman diagrams with n loops
- UV divergencies need to be eliminated
- Fails predicting quantities with essential singularities as $g \rightarrow 0$

Non-Perturbative



- No straightforward approach
- Can have a natural cut-off for high momenta \Rightarrow No UV divergencies
- Can predict quantities with essential singularities as $g \rightarrow 0$

What is a Lattice?

Definition: Lattice Λ

$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z} \}$, with $\{e_i\}$ any basis of \mathbb{R}^n




Figure: A bidimensional lattice.

What is a Lattice?

Definition: Lattice Λ

$\Lambda = \{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z} \}$, with $\{e_i\}$ any basis of \mathbb{R}^n




Figure: A bidimensional lattice.

Hypercubic lattice

$\{e_i\}$ is the canonical basis of \mathbb{R}^n
 a is called *lattice spacing*.




Figure: A square lattice.

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites
- **Vector fields** $U_\mu(x) \rightarrow U_\mu(n)$ on links




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites
- **Vector fields** $U_\mu(x) \rightarrow U_\mu(n)$ on links
- Object with k indices on k -simplexes




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Lattice Field Theory

Basic idea

Fields can take values only in given parts of the lattice, $x \rightarrow n \in \Lambda$.

Examples:

- **Scalar fields** $\Phi(x) \rightarrow \Phi(n)$ on sites
- **Vector fields** $U_\mu(x) \rightarrow U_\mu(n)$ on links
- Object with k indices on k -simplexes




Figure: A (hyper)cubic lattice in \mathbb{R}^3 .

Attention!

Spinorial fields are trickier to be discretized.

Gauge-Invariant Observables and Wilson Action

The Yang-Mills continuum action is

$$S = \frac{1}{4} \int d^4x F^{a\mu\nu}(x) F_{\mu\nu}^a(x).$$

On the lattice, every closed path is gauge-invariant.




Figure: Gauge-invariant paths on a bidimensional lattice.[1]

Gauge-Invariant Observables and Wilson Action

The Yang-Mills continuum action is
$$S = \frac{1}{4} \int d^4x F^{a\mu\nu}(x) F_{\mu\nu}^a(x).$$

On the lattice, every closed path is gauge-invariant.

Definition: Plaquette $U_{\mu\nu}(n)$

$$U_\mu(n) U_\nu(n + \mu) U_\mu^\dagger(n + \nu) U_\nu^\dagger(n)$$




Figure: Gauge-invariant paths on a bidimensional lattice.[1]

Gauge-Invariant Observables and Wilson Action

The Yang-Mills continuum action is
$$S = \frac{1}{4} \int d^4x F^{a\mu\nu}(x) F_{\mu\nu}^a(x).$$

On the lattice, every closed path is gauge-invariant.

Definition: Plaquette $U_{\mu\nu}(n)$

$$U_\mu(n) U_\nu(n + \mu) U_\mu^\dagger(n + \nu) U_\nu^\dagger(n)$$

Wilson's Idea

$$S = \frac{\beta}{2N} \sum_{n,\mu,\nu} \Re \text{Tr} (1 - U_{\mu\nu}(n))$$




Figure: Gauge-invariant paths on a bidimensional lattice.[1]

Polyakov Loops and Potential

If the time coordinate is taken to be periodic, more closed paths arise.

Polyakov Loop

$$P(n) = \text{Tr} \prod_{t=0}^{T-1} U_t(n)$$




Figure: Gauge-invariant paths on a bidimensional lattice.[1]

Polyakov Loops and Potential

If the time coordinate is taken to be periodic, more closed paths arise.

The expectation value of two Polyakov loops is the potential.

Polyakov Loop

$$P(n) = \text{Tr} \prod_{t=0}^{T-1} U_t(n)$$

Potential

$$V(R) = -\frac{1}{T} \log \langle P(0) P^\dagger(R) \rangle$$




Figure: Gauge-invariant paths on a bidimensional lattice.[1]

Lattice symmetries

- [1] Dibakar Sigdel. “Two Dimensional Lattice Gauge Theory with and without Fermion Content”. In: *FIU Electronic Theses and Dissertations* 3224 (2016). DOI: 10.25148/etd.FIDC001748. URL: https://digitalcommons.fiu.edu/etd/3224?utm_source=digitalcommons.fiu.edu%2Fetd%2F3224&utm_medium=PDF&utm_campaign=PDFCoverPages.