GREEEN tallgrass prairie restoration: RLQ analyses (NW Station)

Markus Bauer

2025-09-02

Contents

Preparation **1**
Checks . . . 2
Finalize preparation . . . 3

Statistics **4**
Descriptive analyses . . . 4
Ordinations . . . 4
RLQ analyses . . . 4
Global test . . . 4
Traits by environment crossed table . . . 5

Plots **5**

Save **9**

Session info **9**

Technichal University of Munich, TUM School of Life Sciences, Chair of Restoration Ecology, Emil-Ramann-Straße 6, 85354 Freising, Germany

*markus1.bauer@tum.de

ORCiD ID: 0000-0001-5372-4174 Google Scholar GitHub: markus1bauer GitHub: TUM-Restoration-Ecology

Preparation

```
library(here)
library(tidyverse)
library(ade4)
```

Packages

```
sites <- read_csv(
  here("data", "processed", "data_processed_sites.csv"),
  col_names = TRUE, na = c("na", "NA", ""), col_types =
    cols(
    .default = "?",
    seeding_time = "f"
  )) %>%
```

```r
  arrange(id_plot_year) %>%
  filter(
    site == "NW Station",
    richness_type == "seeded_richness",
    treatment_id %in% c("1", "2", "3", "4")
  ) %>%
  mutate(
    treatment = str_c(seeding_time, herbicide, seeded_pool, sep = "_"),
    treatment = factor(treatment)
  ) %>%
  filter(treatment != "unseeded_0_0") %>%
  select(
    id_plot_year, year, treatment, water_cap, cover_non_seeded,
    cover_seeded_grass
    )

traits <- read_csv(
  here("data", "processed", "data_processed_traits.csv"),
  col_names = TRUE, na = c("na", "NA", ""), col_types = cols(.default = "?")
  ) %>%
  filter(seeded == "1", family != "Poaceae") %>%
  select(accepted_name, sla, height, seedmass) %>%
  mutate(seedmass = log(seedmass)) %>%
  drop_na()

species <- read_csv(
  here("data", "processed", "data_processed_species.csv"),
  col_names = TRUE, na = c("na", "NA", ""), col_types = cols(.default = "?")
  ) %>%
  filter(plot_size %in% c("1", "25")) %>%
  mutate(abundance = 1) %>%
  select(id_plot_year, accepted_name, abundance) %>%
  semi_join(sites, by = "id_plot_year") %>%
  semi_join(traits, by = "accepted_name") %>%
  arrange(id_plot_year)
```

Load data

Checks

```r
species %>%
  group_by(id_plot_year) %>%
  count()
```

Seeded species richness

```
## # A tibble: 47 x 2
## # Groups:   id_plot_year [47]
##    id_plot_year     n
##    <chr>        <int>
##  1 NWS_11_2018      1
##  2 NWS_13_2015      1
##  3 NWS_13_2016      1
##  4 NWS_13_2018      1
```

```
##  5 NWS_14_2015      4
##  6 NWS_14_2016      3
##  7 NWS_14_2017      6
##  8 NWS_14_2018      3
##  9 NWS_15_2016      1
## 10 NWS_15_2017      1
## # i 37 more rows
```

```
traits %>% anti_join(species, by = "accepted_name")
```

Check non-occuring seeded species

```
## # A tibble: 15 x 4
##    accepted_name                   sla height seedmass
##    <chr>                         <dbl>  <dbl>    <dbl>
##  1 Asclepias syriaca             158.   1.23     -5.08
##  2 Chamaecrista fasciculata      297.   0.73     -4.87
##  3 Eryngium yuccifolium           91.7  1.01     -5.57
##  4 Helianthus occidentalis       184.   0.537    -6.11
##  5 Lespedeza virginica            12.6  0.708    -6.26
##  6 Liatris cylindracea           110.   0.564    -5.73
##  7 Lupinus perennis              167.   0.326    -3.84
##  8 Silphium terebinthinaceum     171.   0.716     2.37
##  9 Solidago juncea                13.0  0.783    -7.79
## 10 Solidago nemoralis             12.2  0.61     -7.84
## 11 Solidago speciosa             263.   0.71     -8.27
## 12 Symphyotrichum oolentangiense  14.3  0.362    -8.68
## 13 Tradescantia ohiensis         103.   0.921    -5.61
## 14 Verbena stricta               183.   0.765    -6.89
## 15 Zizia aptera                   10.0  0.34     -6.42
```

Finalize preparation

```
traits <- traits %>%
  semi_join(species, by = "accepted_name") %>%
  column_to_rownames(var = "accepted_name")

sites <- sites %>%
  semi_join(species, by = "id_plot_year") %>%
  column_to_rownames(var = "id_plot_year") %>%
  select(-water_cap)

species <- species %>%
  pivot_wider(
    names_from = "accepted_name", values_from = "abundance", values_fill = 0
    ) %>%
  column_to_rownames(var = "id_plot_year")
```

Statistics

Descriptive analyses

```
sites %>%
  group_by(year) %>%
  count()
```

```
## # A tibble: 4 x 2
## # Groups:   year [4]
##    year     n
##   <dbl> <int>
## 1  2015     9
## 2  2016    11
## 3  2017    13
## 4  2018    14
```

Ordinations

```
dudi_l <- dudi.coa(species, scannf = FALSE)
dudi_r <- dudi.hillsmith(
  sites,
  row.w = dudi_l$lw,
  scannf = FALSE
  )
dudi_q <- dudi.pca(
  traits,
  row.w = dudi_l$cw,
  scannf = FALSE
  )
```

RLQ analyses

```
m1 <- rlq(
  dudiR = dudi_r,
  dudiL = dudi_l,
  dudiQ = dudi_q,
  scannf = FALSE
  )
```

Global test

```
randtest(m1, nrepet = 999, modeltype = 6)
```

```
## class: krandtest lightkrandtest
## Monte-Carlo tests
## Call: randtest.rlq(xtest = m1, nrepet = 999, modeltype = 6)
##
## Number of tests:   2
##
## Adjustment method for multiple comparisons:   none
## Permutation number:   999
##      Test       Obs    Std.Obs   Alter Pvalue
```

```
## 1 Model 2 0.3573247  2.8003928 greater  0.021
## 2 Model 4 0.3573247 -0.2756001 greater  0.572
```

Traits by environment crossed table

```
m1$tab
```

```
##                              sla      height    seedmass
## year               -0.0004125142 -0.01124349  0.11849854
## treat.fall_0_33     0.3068994066  0.32467864  0.06251722
## treat.spring_0_33  -0.6019771690 -0.03359695 -0.26293183
## treat.spring_1_33  -0.1720737061 -0.53007330  0.04589307
## cover_non_seeded    0.0149149180  0.01038072 -0.21287043
## cover_seeded_grass  0.0205877656  0.07473488  0.11899757
```

Plots

```
plot(m1)
```




```
s.label(m1$lR)
```


d = 1
NWS_13_2018
NWS_57_2018
NWS_9_2018
NWS_18_2016
NWS_57_2017
NWS_9_2016
NWS_57_2016
NWS_36_2018
NWS_29_2018
NWS_57_2015
NWS_44_2017
NWS_47_2016
NWS_14_2016
NWS_9_2015
NWS_49_2018
NWS_47_2015
NWS_27_2015
NWS_42_2015
NWS_14_2015
NWS_36_2015

```
s.label(m1$lQ)
```


d = 1
Desmodium illinoense
Asclepias tuberosa
Echinacea purpurea
Lespedeza capitata
Rudbeckia hirta
Coreopsis lanceolata
Dalea purpurea
Monarda punctata
Symphyotrichum ericoides
Symphyotrichum laeve
Penstemon hirsutus
Monarda fistulosa

```
s.arrow(m1$l1)
```


d = 0.5
treat.spring_1_3
year
cover_seeded_grass
treat.fall_0_33
cover_non_seeded
treat.spring_0_33

```
s.arrow(m1$c1)
```



Save

Session info

```
## R version 4.5.0 (2025-04-11 ucrt)
## Platform: x86_64-w64-mingw32/x64
## Running under: Windows 11 x64 (build 26100)
##
## Matrix products: default
##   LAPACK version 3.12.1
##
## locale:
## [1] LC_COLLATE=German_Germany.utf8  LC_CTYPE=German_Germany.utf8
## [3] LC_MONETARY=German_Germany.utf8 LC_NUMERIC=C
## [5] LC_TIME=German_Germany.utf8
##
## time zone: America/New_York
## tzcode source: internal
##
## attached base packages:
## [1] stats     graphics  grDevices utils     datasets  methods   base
##
## other attached packages:
##  [1] ade4_1.7-23     lubridate_1.9.4 forcats_1.0.0   stringr_1.5.1
```

```
##  [5] dplyr_1.1.4     purrr_1.1.0     readr_2.1.5     tidyr_1.3.1
##  [9] tibble_3.3.0    ggplot2_3.5.2   tidyverse_2.0.0 here_1.0.1
## [13] tinytex_0.57
##
## loaded via a namespace (and not attached):
##  [1] bit_4.6.0          gtable_0.3.6       crayon_1.5.3       compiler_4.5.0
##  [5] Rcpp_1.1.0         tidyselect_1.2.1   parallel_4.5.0     scales_1.4.0
##  [9] yaml_2.3.10        fastmap_1.2.0      R6_2.6.1           generics_0.1.4
## [13] knitr_1.50         MASS_7.3-65        rprojroot_2.0.4    pillar_1.11.0
## [17] RColorBrewer_1.1-3 tzdb_0.5.0         rlang_1.1.6        utf8_1.2.6
## [21] stringi_1.8.7      xfun_0.52          bit64_4.6.0-1      timechange_0.3.0
## [25] cli_3.6.5          withr_3.0.2        magrittr_2.0.3     digest_0.6.37
## [29] grid_4.5.0         vroom_1.6.5        rstudioapi_0.17.1  hms_1.1.3
## [33] lifecycle_1.0.4    vctrs_0.6.5        evaluate_1.0.4     glue_1.8.0
## [37] farver_2.1.2       rmarkdown_2.29     tools_4.5.0        pkgconfig_2.0.3
## [41] htmltools_0.5.8.1
```