

Groundfish Distribution Centroids

Markus Min

2026-03-25

This plot shows the latitudinal centroid of the distribution of each of the species:

yellowtail



rougheyen






Figure 1: Latitudinal centroid of each species. Note that the longitudinal centroid is not shown, as for many species this is on land.

CPUE Maps

Black Rockfish

Canary Rockfish

Copper Rockfish

Dover Sole

Lingcod

Petrale Sole

Quillback Rockfish

Rex Sole

Rougheye/Blackspotted Rockfish

Sablefish

Vermilion & Sunset Rockfish

Yellowtail Rockfish




Figure 2: CPUE map for Black Rockfish.




Figure 3: CPUE map for Canary Rockfish.




Figure 4: CPUE map for Copper Rockfish.




Figure 5: CPUE map for Dover Sole.




Figure 6: CPUE map for Lingcod.




Figure 7: CPUE map for Petrale Sole.




Figure 8: CPUE map for Quillback Rockfish.




Figure 9: CPUE map for Rex Sole.




Figure 10: CPUE map for Rougheye/Blackspotted Rockfish.




Figure 11: CPUE map for Sablefish.




Figure 12: CPUE map for Vermilion & Sunset Rockfish.




Figure 13: CPUE map for Yellowtail Rockfish.