


Griffiths & Squibb Figure 6 · from the paper to WSPRadar

G3ZIL – G4HZX | 5–7 April 2017 | Panels B and C use the same 6,459 frozen paired observations


A Published Figure 6



- R1-R4: fixed paper contour regions
- ✕ Matching density peaks
- ☆ Reconstructed global maximum
- P1-P3: isolated scatter witnesses


R1 Early-night negative branch
R3 Strongest evening island

B Reconstruction



R2 Morning positive branch
R4 Late-night negative branch

C WSPRadar view



Same observations, native WSPRadar presentation

R1-R4 map the same paper bounds onto the app's axis.

P1-P3 mark matched native spots within hourly cells.

Unsmoothed 1-hour × 1-dB cells; pooled median +5 dB.

Markers: hourly medians. Band: hourly middle 50% (IQR).

C uses the app's median-centered nonlinear dB axis.
Read the dB labels; vertical pixel positions differ from A/B.
These medians and IQRs are calculated from frozen pairs;
they are not measurements extracted from the paper.

A/B alignment: printed-axis calibration, with no fitted shift or scale. R1-R4 are contour regions, not confidence intervals.

Matching peaks use the fixed ±4-pixel readout + half-cell allowances; boxes show the source regions alone. P1-P3 match folded time/SNR, not identified dates or stations.

B reconstruction: frozen same-cycle pairs → production 1-hour × 1-dB count grid → fixed Gaussian comparison smoother ($\sigma = 1$ hour, 1 dB; UTC wraps).

The full −17 to +40 dB sample range enters smoothing; A/B display −15 to +25 dB. Faint points are native pairs; gray ✕ mark other density peaks. C retains the full app view.

Source: Griffiths & Squibb, Practical Wireless, October 2017, p. 25, Fig. 6 (5–7 April). Frozen demo: 5 April 00:00–7 April 23:45 UTC; distance <10,000 km.

Illustration from the existing frozen fixture, not a fresh live run. Author smoothing, contour levels and exact population are unknown: agreement concerns selected features, not the complete distribution.