


Zander Figure 4 · from the paper to WSPRadar

Experiment A, 14 MHz | The same 166 reconstructed pairs, shown with coarse and fine bins

A Published Figure 4

Original raster; orange line: paper’s normal curve.



Rounded paper statistics

Mean -6.8 dB · Reported σ 3.5 dB

Independent archive reconstruction

Mean -6.783 dB · Sample s 3.515 dB


166 receiver-cycle pairs · 37 receiver identities · 12 cycles

All 9 visible regions fall within the original readout allowance.




B Reconstruction in the paper’s bins

Digitized paper bars over the reconstruction.




Agreement check · reconstruction minus digitized density

Maximum |residual| $0.0001414 \text{ dB}^{-1}$



C WSPRadar · 1 dB bins

Same pairs, percent-of-sample bar heights.



Same observations, finer display

The 1 dB bins reveal uneven counts and empty bins that are combined in the paper’s coarser histogram. The mean and standard deviation do not change.

B shows density per dB.
C shows percent of all 166 pairs per 1 dB bin; its percentages sum to 100%.

No fitted curve, time shift or density rescaling.

METHOD Coarse density = count $\div$ (166 $\times$ region width). Eight 2.1 dB regions plus one 4.2 dB region; no clipping or subset renormalization.

LIMIT Agreement supports the pooled-pair reconstruction; the paper leaves exact selectors and histogram sample weighting unresolved.

SOURCE Jens Zander (2022), arXiv:2209.08989v1, Figure 4. Graphical allowances are not statistical confidence intervals. Frozen fixture data and tests are unchanged.