


The crow and the fox

(Solution)

A diagram synthesising all available information is given in the problem. We add to that diagram a Oxy frame, where the origin O is the fox's position, and the angle α , which shows the direction of v_0 with respect to the x -axis.



a) The initial conditions are¹:

- for the stone:

$$\text{position: } \begin{cases} x_0^S = 0 \\ y_0^S = 0 \end{cases} \quad \text{velocity: } \begin{cases} v_{x,0}^S = v_0 \cos \alpha \\ v_{y,0}^S = v_0 \sin \alpha \end{cases}$$

- for the cheese:

$$\text{position: } \begin{cases} x_0^C = L \\ y_0^C = H \end{cases} \quad \text{velocity: } \begin{cases} v_{x,0}^C = 0 \\ v_{y,0}^C = 0 \end{cases}$$

¹For ease of writing, $\|\vec{v}_0\|$ is simply written v_0

1. Since both considered systems (system 1: the stone and system 2: the cheese) are only subject to gravity, their equations of motions are given by Newton's second law :

$$\ddot{\vec{r}}^S(t) = \vec{g} \quad \text{et} \quad \ddot{\vec{r}}^C(t) = \vec{g} \quad \text{avec} \quad \vec{r}(t) = \begin{pmatrix} x(t) \\ y(t) \\ z(t) \end{pmatrix}$$

Projecting both of these equations on the Oxy frame, we get four equations of motion:

- Equation for the stone, projected onto the Ox horizontal axis:

$$\ddot{x}^S(t) = 0.$$

Let's integrate this once:

$$\dot{x}^S(t) = cte.$$

The initial condition $v_{x,0}^S \equiv \dot{x}^S(t = t_0) = v_0 \cos \alpha$ allows to compute the constant

$$\dot{x}^S(t) = v_0 \cos \alpha.$$

To find the position, we integrate a second time:

$$x^S(t) = (v_0 \cos \alpha)t + cte'.$$

The initial condition $x_0^S = 0$ gives us $cte' = 0$, and so

$$x^S(t) = (v_0 \cos \alpha)t.$$

- Equation for the stone, projected on Oy :

$$\ddot{y}^S(t) = -g.$$

We integrate once and use the initial condition $v_{y,0}^S = v_0 \sin \alpha$

$$\dot{y}^S(t) = -gt + v_0 \sin \alpha.$$

And we integrate and use the initial condition $y_0^S = 0$ again to find

$$y^S(t) = -\frac{1}{2}gt^2 + (v_0 \sin \alpha)t.$$

- Equations for the cheese projected onto Ox and Oy : Using the same method as above, we find

$$\begin{cases} \ddot{x}^C(t) = 0 \\ \ddot{y}^C(t) = -g \end{cases} \quad \begin{cases} \dot{x}^C(t) = 0 \\ \dot{y}^C(t) = -gt \end{cases} \quad \begin{cases} x^C(t) = L \\ y^C(t) = -\frac{1}{2}gt^2 + H. \end{cases}$$

2. The cheese and the stone do collide if there is a time $t = t_{coll}$ at which they have the same position, that is, if the following conditions are met:

$$x^S(t_{coll}) = x^C(t_{coll}) \quad \text{ET} \quad y^S(t_{coll}) = y^C(t_{coll}).$$

The condition $x^S(t_{coll}) = x^C(t_{coll})$ gives

$$(v_0 \cos \alpha)t_{coll} = L \Rightarrow t_{coll} = \frac{L}{v_0 \cos \alpha} = \frac{\sqrt{L^2 + H^2}}{v_0}. \quad (1)$$

While the condition $y^S(t_{coll}) = y^C(t_{coll})$ gives

$$-\frac{1}{2}gt_{coll}^2 + (v_0 \sin \alpha)t_{coll} = -\frac{1}{2}gt_{coll}^2 + H \Rightarrow t_{coll} = \frac{H}{v_0 \sin \alpha} = \frac{\sqrt{L^2 + H^2}}{v_0}. \quad (2)$$

Here, we used $\tan \alpha = \frac{\sin \alpha}{\cos \alpha} = \frac{H}{L}$. The solutions (??) and (??) are identical: there is, indeed, a cheese-stone collision at time

$$t_{coll} = \frac{\sqrt{L^2 + H^2}}{v_0}. \quad (3)$$

We find that the expression for t_{coll} does not depend on g : g accelerates both objects vertically and in the same fashion (mathematically, that's the term in $-\frac{1}{2}gt^2$), which has no influence on the horizontal motion of the objects, and so none on t_{coll} either.

- b) We just proved that the the stone and the cheese will always collide, no matter the initial speed v_0 . Isn't that surprising? However, if v_0 is too small, the cheese will hit the ground before the collision happens: we must restrict v_0 in that respect. This is not contradicting the previous statement: the collision would just "happen" at $y < 0$, that is, "in the ground". We would just have to dig a big enough hole to be able to see it.

In order for the collision to happen above ground, we must ensure $y^C(t_{coll}) > 0$, that is

$$\begin{aligned} y^C(t_{coll}) &= -\frac{1}{2}gt_{coll}^2 + H > 0 \\ \Rightarrow H &> \frac{1}{2}gt_{coll}^2 = \frac{1}{2}g \frac{L^2 + H^2}{v_0^2} \\ \Rightarrow v_0 &> \sqrt{g \frac{L^2 + H^2}{2H}}. \end{aligned} \quad (4)$$

c) Let's check that the results we found are in the right dimensions.

- Equation ?? : $t_{coll} = \frac{\sqrt{L^2 + H^2}}{v_0} : \frac{([m]^2)^{1/2}}{[m/s]} = \frac{[m]}{[m/s]} = [s]$. We find units of time.
- Equation ?? : $v_0 > \sqrt{g \frac{L^2 + H^2}{2H}} : \left([m/s^2] \frac{[m]^2}{[m]}\right)^{1/2} = \frac{[m]^{1/2} [m]}{[s] [m]^{1/2}}$. We find units of speed.

Let's see a few limiting cases to check if we find what we expect.

- If v_0 tends towards infinity, t_{coll} must tend to 0 (Eq.??) and the collision will happen above ground (Eq.??)
- If v_0 tends to 0, t_{coll} must tend towards infinity (Eq.??) and the collision cannot happen above ground since Eq.?? is never satisfied.
- If g tends to 0, the right-hand side of Eq.?? tends to 0, so the collision will happen above ground, whatever the speed $v_0 > 0$.
- If H tends to 0, the right-hand side of Eq.?? tends towards infinity, so the collision cannot happen above ground.

Projet ExoSet La section de physique de l'EPFL met à disposition de ses étudiants une collection de problèmes puisés dans les séries des enseignants de première année. Les utilisateurs de cette plateforme sont tenus de faire un usage loyal (fair use) des ressources documentaires en ligne mises à leur disposition, reproduction et diffusion interdite.

Soumis par: F. Blanc, O. Schneider, J.-Ph. Brantut, J.-M. Fürbringer