



Rationality of Real Conic Bundles with Quartic Discriminant Curve

Mattie Ji

Brown University

Advised by Lena Ji

2022 Mathematics REU Program - University of Michigan



Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles

Irrational Conic Bundles

Main Theorem 2

Next Steps

For this talk,

- This talk is intended to be accessible to undergraduates who has taken a first course in Abstract Algebra (ex. MATH1530)
- Unless otherwise specified, we will work over \mathbb{R} as our ground field



Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

1 Background in Algebraic Geometry

2 What is a Conic Bundle

3 What is Rationality

4 Main Theorem 1

Rational Conic Bundles
Irrational Conic Bundles

5 Main Theorem 2

6 Next Steps



What are Conics?

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality


Main Theorem 1

Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

In Ancient Greece, conics (or conic sections) are defined as the intersection of a cone and a plane, by “slicing” a cone in creative ways.



1

¹Figure taken from
https://en.wikipedia.org/wiki/File:Conic_Sections.svg



In Algebraic Geometry, our classical conic sections become part of affine spaces.

Definition

Let $n \geq 0$ an integer, the **affine space** of dimension n is \mathbb{R}^n , which we will denote as \mathbb{A}^n

Definition

An (affine) **algebraic variety** $V \subset \mathbb{A}^n$ is the set of common \mathbb{R} -roots of a collection of polynomials $\{F_i\}_{i \in I}$ where $F_i \in \mathbb{R}[x_1, \dots, x_n]$. We write V as

$$V = \mathbb{V}(\{F_i\}_{i \in I})$$



Example of Affine Algebraic Varieties

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

Example:

In the affine space \mathbb{A}^n

- $\mathbb{V}(0) = \mathbb{A}^n$, $\mathbb{V}(1) = \emptyset$

Take $n = 2$, then

- $\mathbb{V}(y - x)$ is the line $y = x$ through \mathbb{R}^2
- $\mathbb{V}(y - x^2)$ is the quadratic line $y = x^2$ in \mathbb{R}^2
- the classical conic section C is the variety

$$C = \mathbb{V}(ax^2 + by^2 + c + dxy + ey + fx)$$

where $a, b, c, d, e, f \in \mathbb{R}$

The theory of affine varieties is great, but we can generalize conics with what's known as “**projective spaces**”.

Definition

The set of 1-dimensional subspaces of \mathbb{A}^{n+1} is called the **projective space** of dimension n , denoted as \mathbb{P}^n . In other words, they are just the set of lines going through the origin in \mathbb{A}^{n+1} .

Notations:

- We will denote the line through 0 and (a_0, \dots, a_n) as $[a_0 : \dots : a_n]$ in \mathbb{P}^n .
- Sometimes we will denote \mathbb{P}^n as $\mathbb{P}_{[x_0, \dots, x_n]}^n$ to emphasize its coordinates.



Why Projective Spaces?

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

Q: Why do we want to study conics in projective spaces rather than affine spaces?

A: There are 2 reasons:

- Geometrically, projective spaces are a natural compactification of affine spaces.
- Algebraically, we can turn conic sections into a class of what's called "homogeneous polynomials", which is generally nicer to work with.



Embedding the Affine Plane

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1


Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

We can embed the affine plane \mathbb{A}^2 into \mathbb{P}^2 by identifying \mathbb{A}^2 with the subset $U_Z = \{[X : Y : Z] \in \mathbb{P}^2 \mid Z \neq 0\}$ via:

$$\varphi_Z : U_Z \rightarrow \mathbb{A}^2, [X : Y : Z] \mapsto \left(\frac{X}{Z}, \frac{Y}{Z}\right)$$



This gives a compactification $\mathbb{P}^2 = \mathbb{A}^2 \sqcup \mathbb{P}^1$



Homogeneous Polynomials

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

Definition

A polynomial $F \in \mathbb{R}[x_0, \dots, x_n]$ is called **homogeneous of degree d** if it is a sum of degree d monomials.

For example, in $\mathbb{R}[x, y, z]$,

$$6x^5 + 7y^5 + \pi x^4 y + 3x^2 y^2 z + 9z^5$$

is a homogeneous polynomial of degree 5.

Observation:

Let F be a homogeneous polynomial of degree d and $\lambda \in \mathbb{R}$,

$$F(\lambda a_0, \dots, \lambda a_n) = \lambda^d F(a_0, \dots, a_n)$$

for all $(a_0, \dots, a_n) \in \mathbb{R}^{n+1}$. In particular, if (a_0, \dots, a_n) is a root of F , then so is $(\lambda a_0, \dots, \lambda a_n)$.



Connection to Conics

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

Classically, conic sections have been considered as real roots of the polynomial

$$f(x, y) = ax^2 + by^2 + c + dxy + ey + fx \in \mathbb{R}[x, y]$$

With our embedding, we can homogenize $f(x, y)$ into:

$$F(X, Y, Z) = aX^2 + bY^2 + cZ^2 + dXY + eYZ + fXZ$$

Then we note that on $Z = 1$, $F(X, Y, Z)$ becomes $f(x, y)$. This is in fact a **bijective correspondence**.

Definition:

A **plane conic** $C \subset \mathbb{P}_{[X:Y:Z]}^2$ is the real roots of a homogeneous polynomial of degree 2 in $\mathbb{R}[X, Y, Z]$.



Matrices and Conics

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

Take any homogenous polynomial of degree 2

$$F(X, Y, Z) := aX^2 + bY^2 + cZ^2 + dXY + eYZ + fXZ$$

We note that this polynomial has an associated **symmetric matrix**

$$M_F = \begin{bmatrix} a & \frac{d}{2} & \frac{f}{2} \\ \frac{d}{2} & b & \frac{e}{2} \\ \frac{f}{2} & \frac{e}{2} & c \end{bmatrix}$$

such that

$$F(X, Y, Z) = \begin{bmatrix} X & Y & Z \end{bmatrix} M_F \begin{bmatrix} X \\ Y \\ Z \end{bmatrix}$$

This also makes $F(X, Y, Z)$ into what's called a **quadratic form** of 3 variables.



Smoothness of Conics

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

It turns out that the rank of the matrix M_F determines the geometry of the conic C .

Fact:

Let $C_{\mathbb{C}}$ be all the complex roots of F (in particular $C \subseteq C_{\mathbb{C}}$)

- If M_F has rank 3, then $C_{\mathbb{C}}$ is a **smooth conic**
- If M_F has rank 2, then $C_{\mathbb{C}}$ is the union of **two distinct lines** meeting at a point.
- If M_F has rank 1, then $C_{\mathbb{C}}$ is a **double line**.



Example: $\text{rank}(M_F) = 3$

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles

Irrational Conic Bundles


Main Theorem 2

Next Steps

Let $M_F = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$, $F(X, Y, Z) = X^2 + Y^2 - Z^2$.

Then C is a smooth conic.

On the chart ($Z \neq 0$),





Example: $\text{rank}(M_F) = 2$

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1


Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

Let $M_F = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $F(X, Y, Z) = X^2 - Y^2$.

Then C is the union of two lines meeting at the origin.
On the chart ($Z \neq 0$),





Example: $\text{rank}(M_F) = 1$

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles


Irrational Conic Bundles

Main Theorem 2

Next Steps

Let $M_F = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $F(X, Y, Z) = X^2$.

Then C is a line, we say it's "double" because of the square.
On the chart ($Z \neq 0$),





Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

- 1 Background in Algebraic Geometry
- 2 What is a Conic Bundle
- 3 What is Rationality
- 4 Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles
- 5 Main Theorem 2
- 6 Next Steps



Definition:

A **conic bundle** over \mathbb{P}^2 is a “nice”² morphism $\pi : X \rightarrow \mathbb{P}^2$ such that

- X is a smooth variety
- The fiber over every point $p \in \mathbb{P}^2$ is a conic
- The generic fiber is a smooth conic

²A proper flat \mathbb{R} -morphism



In our research, we are interested in the **conic bundle**

$\pi_2 : Y_{\tilde{\Delta}/\Delta} \rightarrow \mathbb{P}_{[u:v:w]}^2$ where:

- $Y_{\tilde{\Delta}/\Delta}$ is a variety defined by the equation³:

$$z^2 = Q_1(u, v, w)t_0^2 + 2Q_2(u, v, w)t_0t_1 + Q_3(u, v, w)t_1^2$$

- $Q_1, Q_2, Q_3 \in \mathbb{R}[u, v, w]$ are homogenous polynomials of degree 2
- π_2 is the standard projection that forgets z, t_0 , and t_1

³This looks like a very specific choice, but it turns out that every degree 4 conic bundle $X \rightarrow \mathbb{P}^2$ is birationally equivalent to some π_2 “up to a class in $\mathbb{Z}/2\mathbb{Z}$ ” (Theorem 2.6 of [FJS⁺] based on [Bru08])



Why is π_2 a conic bundle?

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

Intuitively, every point in \mathbb{P}^2 should correspond to some conic in $Y_{\tilde{\Delta}/\Delta}$.

Example of Fibers for π_2 :

Concretely, take the point $[1 : 2 : 3] \in \mathbb{P}_{[u:v:w]}^2$, then fiber of $[1 : 2 : 3]$ is exactly the solutions satisfying:

$$0 = Q_1(1, 2, 3) t_0^2 + 2Q_2(1, 2, 3) t_0 t_1 + Q_3(1, 2, 3) t_1^2 - z^2$$

This forms a conic in $\mathbb{P}_{[t_0:t_1:z]}^2$.



Quadric Surface Bundle

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

We are also interested in the map $\pi_1 : Y_{\tilde{\Delta}/\Delta} \rightarrow \mathbb{P}_{[t_0:t_1]}^1$, which is the standard projection that forgets z, u, v , and w .

Example of Fibers for π_1 :

Similarly, take the point $[1 : 3] \in \mathbb{P}_{[t_0:t_1]}^1(\mathbb{R})$, then fiber of $[1 : 3]$ is exactly the solutions satisfying:


$$\begin{aligned} z^2 &= Q_1(u, v, w)(1)^2 + 2Q_2(u, v, w)(1)(3) + Q_3(u, v, w)(3)^2 \\ &= (1)Q_1(u, v, w) + 2(3)Q_2(u, v, w) + (9)Q_3(u, v, w) \end{aligned}$$

This forms a degree 2 surface (known as a **quadric**) in $\mathbb{P}_{[u:v:w:z]}^3(\mathbb{R})$.

$\pi_1 : Y_{\tilde{\Delta}/\Delta} \rightarrow \mathbb{P}_{[t_0:t_1]}^1$ is an example of a **quadric surface bundle**.




Putting π_1 and π_2 together, we have the commutative diagram:



In this talk, we refer to this as the **double cover model**.



This commutative diagram also induces a diagram between **their real points**:





The Discriminant Curve

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

We would like to identify if a given fiber of π_2 is smooth:

Smoothness Criterion

Given fixed $[u : v : w] \in \mathbb{P}_{[u:v:w]}^2(\mathbb{R})$, we can rewrite its associated conic as:

$$0 = Q_1(u, v, w)t_0^2 + 2Q_2(u, v, w)t_0t_1 + Q_3(u, v, w)t_1^2 + (-1)z^2 (*)$$

This gives the symmetric matrix:

$$M = \begin{bmatrix} Q_1(u, v, w) & Q_2(u, v, w) & 0 \\ Q_2(u, v, w) & Q_3(u, v, w) & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

The conic $(*)$ is smooth if and only if $\det(M) \neq 0$.



The Discriminant Curve

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2


Next Steps

Smoothness Criterion:

Let M be the symmetric matrix associated to each conic of $Y_{\tilde{\Delta}/\Delta}$, the curve defined by $\det(M) = 0$ is called the **discriminant curve** Δ :

$$\Delta = (Q_1 Q_3 - Q_2^2 = 0) \subset \mathbb{P}_{[u:v:w]}^2$$

The fiber of $s \in \mathbb{P}_{[u:v:w]}^2$ is smooth if and only if $s \notin \Delta$





The Double Cover of Δ

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles

Irrational Conic Bundles

Main Theorem 2


Next Steps

Fact:

There exists a curve $\tilde{\Delta} \subset \mathbb{P}^4_{[u:v:w:r:s]}$ defined by

$$\tilde{\Delta} := (Q_1 - r^2 = Q_2 - rs = Q_3 - s^2 = 0)$$

such that the projection $\tilde{\Delta} \rightarrow \Delta$ is a double cover.
 $\tilde{\Delta}$ is called the **double cover** of Δ .





Quartic Plane Curves

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

$Q_1Q_3 - Q_2^2$ is a degree 4 homogeneous real polynomial.

Definition

The roots of a degree 4 homogenous polynomial over \mathbb{P}^2 is known as a **quartic**.

Theorem (Zeuthen, 1874 [Zeu74])

Let Δ be a smooth quartic over \mathbb{R} , then $\Delta(\mathbb{R})$ can be classified into 1 of the 6 following topological types:

- 1 No real points
- 2 One oval
- 3 Two nested ovals
- 4 Two non-nested ovals
- 5 Three ovals
- 6 Four ovals



Example: Four Ovals

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles
Irrational Conic Bundles


Main Theorem 2

Next Steps

The homogeneous equation defines a smooth quartic whose real component has 4 ovals:

$$0 = -11u^4 - 5u^2v^2 - 2v^4 + 11u^2w^2 + 4v^2w^2 - 3w^4$$

The real components on the chart ($w \neq 0$)





Example: Two Nested Ovals

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles

Irrational Conic Bundles


Main Theorem 2

Next Steps

This homogeneous equation defines a smooth quartic whose real component has 2 nested ovals:

$$0 = -3u^4 - \frac{7}{10}u^2v^2 - \frac{169}{400}v^4 + \frac{67}{6}u^2w^2 + \frac{949}{240}v^2w^2 - \frac{121}{576}w^4$$

The real components on the chart ($w \neq 0$):





Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

- 1 Background in Algebraic Geometry
- 2 What is a Conic Bundle
- 3 What is Rationality**
- 4 Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles
- 5 Main Theorem 2
- 6 Next Steps



Overview of Rationality

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

- A variety X is **k -rational** over a field k if there exists non-empty open sets $U \subset X$ and $V \subset \mathbb{P}_k^{\dim X}$ such that U and V are isomorphic over k .
- If X is not rational over k , we say that X is **irrational** over k .

There are two relevant facts about rationality:

- **Lang–Nishimura Lemma:** If X is a projective k -rational variety, then $X(k)$ is non-empty.
- **General Topological Fact:** If X is a smooth projective \mathbb{R} -rational variety, then $X(\mathbb{R})$ is connected.

In particular,

$Y_{\tilde{\Delta}/\Delta}$ is \mathbb{R} -rational $\implies Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is non-empty and connected

Q: Why do we care about rationality?

A: Given an algebraic variety, a natural question to ask is how simple it is:

- One notion of simplicity is its **closeness to projective spaces**.
- It turns out that rational algebraic varieties are **“birationally equivalent”** to projective spaces.
- Conic bundles have been a very rich source of examples of varieties with varying levels of similarity to projective space.



Criterion for Rationality

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

The \mathbb{C} -rationality of $Y_{\tilde{\Delta}/\Delta}$ is quite clear:

Theorem (Iskovskikh, 1987 [Isk87])

$Y_{\tilde{\Delta}/\Delta}$ is always \mathbb{C} -rational

The \mathbb{R} -rationality of $Y_{\tilde{\Delta}/\Delta}$ is more complicated, but the work of S. Frei, L. Ji, S. Sankar, B. Viray, and I. Vogt previously gave a very useful criterion:

Proposition (Proposition 6.1, Frei et al, 2022 [FJS⁺])

Let $Y_{\tilde{\Delta}/\Delta}$ be defined as before. If $\tilde{\Delta}(\mathbb{R}) \neq \emptyset$, then $Y_{\tilde{\Delta}/\Delta}$ is \mathbb{R} -rational

In our REU, we instead investigate what happens when $\tilde{\Delta}(\mathbb{R}) = \emptyset$.



Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

- 1 Background in Algebraic Geometry
- 2 What is a Conic Bundle
- 3 What is Rationality
- 4 Main Theorem 1**
Rational Conic Bundles
Irrational Conic Bundles
- 5 Main Theorem 2
- 6 Next Steps



Main Result 1

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

Theorem (Ji and Ji, 2022 [JJ22]):

Let $\mathcal{CB}_{\emptyset/*}$ denote the set of real conic bundles $X \rightarrow \mathbb{P}^2$ of the form $Y_{\tilde{\Delta}/\Delta}$ with smooth Δ of topological type $*$ and smooth $\tilde{\Delta}$ that has no real points,

- ① $\mathcal{CB}_{\emptyset/\emptyset}$ contains both rational and irrational members
- ② $\mathcal{CB}_{\emptyset/1\text{-oval}}$ contains both rational and irrational members⁴
- ③ $\mathcal{CB}_{\emptyset/2}$ non-nested ovals, $\mathcal{CB}_{\emptyset/2}$ nested ovals, and $\mathcal{CB}_{\emptyset/3\text{-ovals}}$ contains both rational and irrational members⁵
- ④ $\mathcal{CB}_{\emptyset/4\text{-ovals}}$ contains only rational members⁶.

⁴The case of irrational 1 oval was done in [FJS⁺]

⁵The case of irrational 2 non-nested ovals was done in [FJS⁺]

⁶We proved this based on ideas joint with S. Frei, S. Sankar, B. Viray, and I. Vogt

It turns out that there's another rationality construction for rational examples when $\tilde{\Delta}(\mathbb{R})$ is empty:


Theorem (Witt, 1937 [Wit37])

If $\pi_1 : Y_{\tilde{\Delta}/\Delta} \rightarrow \mathbb{P}_{[t_0:t_1]}^1$ is surjective on real points, then $Y_{\tilde{\Delta}/\Delta}$ is \mathbb{R} -rational.

Using this criterion, we can find and check that rational members exist for all topological types of $\Delta(\mathbb{R})$.

Example of Rational One Oval with $\tilde{\Delta}(\mathbb{R}) = \emptyset$

Take $Q_1 := -u^2 + uv - w^2$, $Q_2 := 3u^2 + uv - v^2 + w^2$, and $Q_3 = -u^2 - 2uv - 2w^2$, then one can verify that $\tilde{\Delta}(\mathbb{R}) = \emptyset$, π_1 is surjective on real points, and $\Delta(\mathbb{R})$ is one oval, as seen on the chart ($w \neq 0$):





Obstruction by Disconnected Real Loci

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

Now we will move on to irrational conic bundles:

- Recall that if $Y_{\tilde{\Delta}/\Delta}$ is \mathbb{R} -rational, then $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is **connected** and non-empty.
- What if we take $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ to be **disconnected**?

The key insight is looking at $\pi_1(\mathbb{R}) : Y_{\tilde{\Delta}/\Delta}(\mathbb{R}) \rightarrow \mathbb{P}_{[t_0:t_1]}^1(\mathbb{R})$:

$Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is disconnected $\iff \pi_1(Y_{\tilde{\Delta}/\Delta}(\mathbb{R}))$ is disconnected

Observation:

Take any point $[a : b] \in \mathbb{P}_{[t_0:t_1]}^1(\mathbb{R})$, then fiber of $[a : b]$ is exactly the solutions satisfying:

$$z^2 = Q_1(u, v, w)(a)^2 + 2Q_2(u, v, w)(ab) + Q_3(u, v, w)(b)^2$$

This equation has a solution if and only if the matrix

$$M_{[a:b]} := \left[\begin{array}{c|c} M_1a^2 + 2M_2ab + M_3b^2 & 0 \\ \hline 0 & -1 \end{array} \right]$$

is **NOT** negative definite, where M_1, M_2, M_3 are the symmetric matrices associated to Q_1, Q_2, Q_3 respectively.



Key Observation (Continued)

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

We will denote the signature of a real symmetric matrix M as (p, n) , where p is the number of positive eigenvalues and n is the number of negative eigenvalues.

Lemma (Adapted from Degtjarev Et al., [DIK00])

Given the setup with $Y_{\tilde{\Delta}/\Delta}$, let $M_{[t_0:t_1]}$ be the matrix defined the same as previously, then the signature of $M_{[t_0:t_1]}$ can only change by ± 1 at real solutions of a degree 6 polynomial⁷

$$\Gamma(t) := -\det(M_1 t^2 + 2M_2 t + M_3)$$

Notably, on each interval defined by real points of $\Gamma(t)$, the signature of $M_{[t_0:t_1]}$ stays the same.

⁷It is possible that one of the roots is the point at infinity, but we can without loss choose an appropriate basis to avoid this.



Disconnected Examples

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

Now, as long as we can find examples where at least two non-adjacent intervals with signature $(0, 4)$, this will produce a disconnected $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$.

Irrational Example with $\Delta(\mathbb{R})$ 3 ovals

Define $Q_1 := -u^2 - v^2 - w^2$, $Q_2 := u^2 + 5v^2 + 9w^2$, and $Q_3 := -24v^2 - 80w^2$. Then we have that

$$\Gamma(t) = t^6 - 30t^5 + 340t^4 - 1800t^3 + 4384t^2 - 3840t$$

with 6 real roots

$$t = 0, 2, 4, 6, 8, 10$$



Disconnected Examples

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles

Irrational Conic Bundles


Main Theorem 2

Next Steps

Example (Continued)

The signatures of $M_{[t_0:t_1]}$ follow the pattern:

$$(0, 4), (1, 3), (0, 4), (1, 3), (0, 4), (1, 3)$$





The Explicit Construction

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles

Irrational Conic Bundles

Main Theorem 2

Next Steps

The choice of Q_1, Q_2, Q_3 in the previous example may seem arbitrary, but you can “reverse engineer” the appropriate Q_1, Q_2, Q_3 by considering

$$\begin{bmatrix} a_0t^2 + b_0t + c_0 & 0 & 0 \\ 0 & a_1t^2 + b_1t + c_1 & 0 \\ 0 & 0 & a_2t^2 + b_2t + c_2 \end{bmatrix}$$

$$= t^2(M_1) + t(2M_2) + M_3$$

and compare the coefficients.



Limitations of the Topological Obstruction

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

During the REU, we also showed that

Proposition (Ji and Ji, 2022 [JJ22])

If $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is disconnected, then $\Delta(\mathbb{R})$ is either 2 non-nested ovals, 2 nested ovals, or 3 ovals. More precisely,

- $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ has at most 3 connected components
- If $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ has 2 connected components, then $\Delta(\mathbb{R})$ is either 2 non-nested ovals or 2 nested ovals
- If $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ has 3 connected components, then $\Delta(\mathbb{R})$ is 3 ovals.

Each case described above does occur.



Remark:

When $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ has 2 connected components, the signatures of $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is one of

- ① $(0, 4), (1, 3), (0, 4), (1, 3), (2, 2), (1, 3)$
- ② $(0, 4), (1, 3), (0, 4), (1, 3)$

If pattern (1) occurs, then $\Delta(\mathbb{R})$ is 2 nested ovals.

If pattern (2) occurs, $\Delta(\mathbb{R})$ has always been 2 non-nested ovals experimentally. **It's unknown if the pattern (2) implies that $\Delta(\mathbb{R})$ is 2 non-nested ovals.**



The Case where $Y_{\tilde{\Delta}/\Delta}(\mathbb{R}) = \emptyset$

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

- Recall that if $Y_{\tilde{\Delta}/\Delta}$ is \mathbb{R} -rational, then $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is connected and **non-empty**.
- What if we take $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ to be **empty** instead?

Irrational Example where $\Delta(\mathbb{R}) = \emptyset$

Define $Q_1 := -2u^2 - 3v^2 - 5w^2$, $Q_2 := u^2 + 2v^2 + 3w^2$, and $Q_3 := -2u^2 - 4v^2 - 2w^2$. Then one can verify that the associated $\Delta(\mathbb{R})$ and $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ are both empty, hence $Y_{\tilde{\Delta}/\Delta}$ is irrational over \mathbb{R} .

This doesn't get us really far, as $Y_{\tilde{\Delta}/\Delta}(\mathbb{R}) = \emptyset$ implies that $\Delta(\mathbb{R}) = \emptyset$.



Irrational One Oval

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

S. Frei, L. Ji, S. Sankar, B. Viray, and I. Vogt showed that

Irrational Example (Theorem 1.3(2), [FJS⁺])

Define $Q_1 := -u^2 - v^2 - 3w^2$, $Q_2 := 3u^2 + 5v^2$, and $Q_3 := -7u^2 - 23v^2 - 12w^2$, then $\Delta(\mathbb{R})$ is one oval, and the associated $Y_{\tilde{\Delta}/\Delta}$ is irrational over any subfield of \mathbb{R} .

The main idea behind showing this is irrational relied on what's known as an **Intermediate Jacobian Torsor (IJT) obstruction**:

- Classically, this has been done over \mathbb{C} by **Clemens–Griffiths** (1972, [CG72]).
- Over non-closed fields, a refinement of this technique was introduced by **Hassett–Tschinkel** (2021, [HT21]) and **Benoist–Wittenberg** (2019, [BW]).



Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

- 1 Background in Algebraic Geometry
- 2 What is a Conic Bundle
- 3 What is Rationality
- 4 Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles
- 5 Main Theorem 2
- 6 Next Steps



Main Result 2

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

A natural question from our discovery would be to characterize the rationality of $Y_{\tilde{\Delta}/\Delta}$ in each topological type of $\Delta(\mathbb{R})$. So far, we have showed that⁸:

Theorem (Ji and Ji, 2022 [JJ22]):

Let $Y_{\tilde{\Delta}/\Delta}$ be as before,

- ① If $\Delta(\mathbb{R})$ is 4 ovals, then $Y_{\tilde{\Delta}/\Delta}$ is rational
- ② If $\Delta(\mathbb{R})$ is 3 ovals, then $Y_{\tilde{\Delta}/\Delta}$ is rational $\iff Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is connected
- ③ If $\Delta(\mathbb{R})$ is 2 non-nested ovals, then $Y_{\tilde{\Delta}/\Delta}$ is rational $\iff Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is connected and the IJT obstruction vanishes
- ④ If $\Delta(\mathbb{R})$ is empty, then $Y_{\tilde{\Delta}/\Delta}$ is rational $\iff Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is non-empty and the IJT obstruction vanishes

⁸(1) and (2) were based on or inspired by ideas joint with S. Frei, S. Sankar, B. Viray, and I. Vogt



Proof Sketch

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1

Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2


Next Steps

For (1) and (2),

Fact:

Every smooth real quartic $\Delta(\mathbb{R})$ has 28 bitangents.

- When $\Delta(\mathbb{R})$ is 4 ovals, it has 28 real bitangents.
- When $\Delta(\mathbb{R})$ is 3 ovals, it has 16 real bitangents.





Proof Sketch

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

Define W to be the sub-variety of $Y_{\tilde{\Delta}/\Delta}$ given by the equation

$$0 = Q_1(u, v, w)t_0^2 + 2Q_2(u, v, w)t_0t_1 + Q_3(u, v, w)t_1^2$$

There's an induced projection map $\pi'_1 : W \rightarrow \mathbb{P}^1$ defined similar to π_1 .

Viewing W as $W_{\mathbb{C}}$, $W_{\mathbb{C}}$ has 56 geometric lines:

- 32 of those lines each give a geometric section of π'_1 , call this collection E
- Note that if π'_1 has a section over \mathbb{R} , then $\pi_1(\mathbb{R})$ is surjective, hence $Y_{\tilde{\Delta}/\Delta}$ is rational

In particular, if $(Q_1Q_3 - Q_2^2 < 0)_{\mathbb{R}}$ contains some non-orientable part of \mathbb{P}^2 , then each bitangent of $\Delta(\mathbb{R})$ would correspond to two lines in $W_{\mathbb{C}}$.



Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

Fact:

If $\Delta(\mathbb{R})$ is 4 ovals or 3 ovals with $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is connected, then $(Q_1Q_3 - Q_2^2 < 0)_{\mathbb{R}}$ is not orientable.

Then the rest is really a combinatorics argument, as there'd be some real bitangent of $\Delta(\mathbb{R})$ that will end up giving a line in E , which corresponds to a section over \mathbb{R} .



For (3) and (4), the forward directions are already given. For the converse, we rely on the following proposition

Proposition (Ji and Ji, 2022 [JJ22]):

If $(Q_1Q_3 - Q_2^2 < 0)_{\mathbb{R}}$ contains some non-orientable part of \mathbb{P}^2 ,
 $\pi_1(\mathbb{R})$ is surjective \iff the IJT obstruction vanishes for Y

- For both (3) and (4), one can show that the converse conditions forces $\pi_1(\mathbb{R})$ to be surjective, hence $Y_{\tilde{\Delta}/\Delta}$ is rational.
- We leave this as an exercise to the audience.



Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

- 1 Background in Algebraic Geometry
- 2 What is a Conic Bundle
- 3 What is Rationality
- 4 Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles
- 5 Main Theorem 2
- 6 Next Steps



Next Steps

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2
Next Steps

The next steps would be to characterize the rationality when $\Delta(\mathbb{R})$ is two nested ovals or one oval.

Experimentally, if $\Delta(\mathbb{R})$ is 2 nested ovals, we are confident that the following statement is most likely true:

- $Y_{\tilde{\Delta}/\Delta}$ is rational \iff the IJT obstruction vanishes
 $\iff Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is connected

We actually already know that:

- $Y_{\tilde{\Delta}/\Delta}$ is rational \implies the IJT obstruction vanishes $\implies Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ is connected



The case when $\Delta(\mathbb{R})$ is one oval is a lot more complicated.

- Experimentally, when π_1 has signature $(0, 4)$, $(1, 3)$, $\Delta(\mathbb{R})$ is one oval, $\tilde{\Delta}(\mathbb{R}) = \emptyset$, $(Q_1Q_3 - Q_2^2 < 0)$ is inside the oval
- This situation is tricky because all of our current tools of rationality and irrationality constructions do not satisfy this scenario.

Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle


What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps

If you are interested to learn more, please check out our paper:

 Lena Ji and Mattie Ji. [Rationality of real conic bundles with quartic discriminant curve, 2022](https://arxiv.org/abs/2208.08916)
(<https://arxiv.org/abs/2208.08916>)

I would like to thank

- Lena Ji for advising this project
- Sarah Frei, Brendan Hassett, Soumya Sankar, Bianca Viray, and Isabel Vogt for helpful discussions and conversations
- János Kollár for the question that motivated this project
- The University of Michigan Mathematics REU Program
- The National Science Foundation (Karen Smith's NSF grant DMS-2101075) for funding this project
- My letter of recommendation writers Nicole Looper and Jungang Li



Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps



Nils Bruin.

The arithmetic of Prym varieties in genus 3.

Compos. Math., 144(2):317–338, 2008.



Olivier Benoist and Olivier Wittenberg.

Intermediate Jacobians and rationality over arbitrary fields.

arXiv e-prints, page arXiv:1909.12668.



C. Herbert Clemens and Phillip A. Griffiths.

The Intermediate Jacobian of the cubic threefold.

Ann. of Math., 95(2):281–356, 1972.



Aleksandr Degtjarev, Il'ja V. Itenberg, and Viatcheslav Kharlamov.

Real Enriques surfaces, pages 211–212.

Springer, 2000.



Sarah Frei, Lena Ji, Soumya Sankar, Bianca Viray, and Isabel Vogt.

Curve classes on conic bundle threefolds and applications to rationality.

arXiv e-prints, page arXiv:2207.07093.



Brendan Hassett and Yuri Tschinkel.

Rationality of complete intersections of two quadrics over nonclosed fields.

Enseign. Math., 67(1-2):1–44, 2021.

With an appendix by Jean-Louis Colliot-Thélène.



V. A. Iskovskikh.

On the rationality problem for conic bundles.

Duke Math. J., 54(2):271–294, 1987.



Rationality of
Real Conic
Bundles with
Quartic
Discriminant
Curve

Mattie Ji

Background in
Algebraic
Geometry

What is a Conic
Bundle

What is
Rationality

Main Theorem 1
Rational Conic Bundles
Irrational Conic Bundles

Main Theorem 2

Next Steps



[Lena Ji and Mattie Ji.](#)

Rationality of real conic bundles with quartic discriminant curve, 2022.



[Ernst Witt.](#)

Theorie der quadratischen Formen in beliebigen Körpern.
J. Reine Angew. Math., 176:31–44, 1937.



[Hieronymus Georg Zeuthen.](#)

Sur les différentes formes des courbes planes du quatrième ordre.

Mathematische Annalen, 7:410–432, 1874.