



Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

Main Result

Main Theorem  
Rational Conic Bundles

Irrational Conic  
Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

# Rationality of Real Conic Bundles with Quartic Discriminant Curve

Mattie Ji

Brown University

Advised by Lena Ji

2022 Mathematics REU Program - University of Michigan



## Rationality of Real Conic Bundles with Quartic Discriminant Curve

Mattie Ji

### Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

### Main Result

Main Theorem  
Rational Conic Bundles

### Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

- 1 Background
  - Plane Conics
  - Conic Bundles
  - Real Quartic Plane Curves
  - Rationality
- 2 Main Result
  - Main Theorem
  - Rational Conic Bundles
- 3 Irrational Conic Bundles
  - Disconnected Real Loci
  - Empty Real Loci
  - Irrational One Oval Example



## Rationality of Real Conic Bundles with Quartic Discriminant Curve

Mattie Ji

### Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

### Main Result

Main Theorem  
Rational Conic Bundles

### Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

Throughout this talk,

- Unless otherwise specified, we will work over  $\mathbb{R}$  as our ground field
- We will denote the projective  $n$ -space over the ground field as  $\mathbb{P}^n$
- Sometimes, to emphasize the coordinates  $[X_0 : \dots : X_n]$  of  $\mathbb{P}^n$ , we will denote  $\mathbb{P}^n$  as  $\mathbb{P}^n_{[X_0 : \dots : X_n]}$

# Overview of Conics

- A **plane conic**  $C \subset \mathbb{P}^2$  is the curve defined

$$C := (F(X, Y, Z) = 0)$$

where

$F(X, Y, Z) = aX^2 + bY^2 + cZ^2 + dXY + eYZ + fXZ$   
is a homogeneous polynomial of degree 2 in  $\mathbb{R}[X, Y, Z]$

- This  $C$  has an associated **symmetric matrix**  $M_F$

$$M_F := \begin{bmatrix} a & \frac{d}{2} & \frac{f}{2} \\ \frac{d}{2} & b & \frac{e}{2} \\ \frac{f}{2} & \frac{e}{2} & c \end{bmatrix}$$

such that

$$F(X, Y, Z) = \begin{bmatrix} X & Y & Z \end{bmatrix} M_F \begin{bmatrix} X \\ Y \\ Z \end{bmatrix}$$

- $C$  is **smooth** if and only if  $M_F$  has **rank 3**



## Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

## Main Result

Main Theorem

Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci

Empty Real Loci

Irrational One Oval  
Example

## Definition:

A **conic bundle** over  $\mathbb{P}^2$  is a “nice”<sup>1</sup> morphism  $\pi : X \rightarrow \mathbb{P}^2$  such that

- $X$  is a smooth variety
- The fiber over every point  $p \in \mathbb{P}^2$  is a conic
- The generic fiber is a smooth conic

---

<sup>1</sup>A proper flat  $\mathbb{R}$ -morphism



# Conic Bundles

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

## Main Result

Main Theorem  
Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

In our research, we are interested in the **conic bundle**

$\pi_2 : Y_{\tilde{\Delta}/\Delta} \rightarrow \mathbb{P}_{[u:v:w]}^2$  where:

- $Y_{\tilde{\Delta}/\Delta}$  is a variety defined by the equation<sup>2</sup>:

$$z^2 = Q_1(u, v, w)t_0^2 + 2Q_2(u, v, w)t_0t_1 + Q_3(u, v, w)t_1^2$$

- $Q_1, Q_2, Q_3 \in \mathbb{R}[u, v, w]$  are homogenous polynomials of degree 2
- $\pi_2$  is the standard projection that forgets  $z, t_0$ , and  $t_1$

We are also interested in the map  $\pi_1 : Y_{\tilde{\Delta}/\Delta} \rightarrow \mathbb{P}_{[t_0:t_1]}^1$ , which is the standard projection that forgets  $z, u, v$ , and  $w$ .

---

<sup>2</sup>This looks like a very specific choice, but it turns out that every degree 4 conic bundle  $X \rightarrow \mathbb{P}^2$  is birationally equivalent to some  $\pi_2$  “up to a class in  $\mathbb{Z}/2\mathbb{Z}$ ” (Theorem 2.6 of [FJS<sup>+</sup>] based on [Bru08])



# Why is $\pi_2$ a conic bundle?

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

## Main Result

Main Theorem

Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci

Empty Real Loci

Irrational One Oval  
Example

Intuitively, every point in  $\mathbb{P}^2$  should correspond to some conic in  $Y_{\tilde{\Delta}/\Delta}$ .

## Example of Fibers for $\pi_2$ :

Concretely, take the point  $[1 : 2 : 3] \in \mathbb{P}_{[u:v:w]}^2$ , then fiber of  $[1 : 2 : 3]$  is exactly the solutions satisfying:

$$0 = Q_1(1, 2, 3) t_0^2 + 2Q_2(1, 2, 3) t_0 t_1 + Q_3(1, 2, 3) t_1^2 - z^2$$

This forms a conic in  $\mathbb{P}_{[t_0:t_1:z]}^2$ .



## Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

## Main Result

Main Theorem

Rational Conic Bundles


Irrational Conic  
Bundles

Disconnected Real Loci

Empty Real Loci

Irrational One Oval  
Example

Putting  $\pi_1$  and  $\pi_2$  together, we have the commutative diagram:



In this talk, we refer to this as the **double cover model**.





## Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

## Main Result

Main Theorem

Rational Conic Bundles


Irrational Conic  
Bundles

Disconnected Real Loci

Empty Real Loci

Irrational One Oval  
Example

This commutative diagram also induces a diagram between **their real points**:





# The Discriminant Curve

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

## Main Result

Main Theorem

Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci

Empty Real Loci


Irrational One Oval  
Example

## Smoothness Criterion:

Let  $M$  be the symmetric matrix associated to each conic of  $Y_{\tilde{\Delta}/\Delta}$ , the curve defined by  $\det(M) = 0$  is called the **discriminant curve**  $\Delta$ :

$$\Delta = (Q_1 Q_3 - Q_2^2 = 0) \subset \mathbb{P}_{[u:v:w]}^2$$

The fiber of  $s \in \mathbb{P}_{[u:v:w]}^2$  is smooth if and only if  $s \notin \Delta$





# The Double Cover of $\Delta$

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

## Main Result

Main Theorem  
Rational Conic Bundles

## Irrational Conic Bundles


Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

## Fact:

There exists a curve  $\tilde{\Delta} \subset \mathbb{P}^4_{[u:v:w:r:s]}$  defined by

$$\tilde{\Delta} := (Q_1 - r^2 = Q_2 - rs = Q_3 - s^2 = 0)$$

such that the projection  $\tilde{\Delta} \rightarrow \Delta$  is a double cover.  
 $\tilde{\Delta}$  is called the **double cover** of  $\Delta$ .





# Quartic Plane Curves

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

## Main Result

Main Theorem

Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci

Empty Real Loci

Irrational One Oval  
Example

$Q_1Q_3 - Q_2^2$  is a degree 4 homogeneous real polynomial.

## Definition

The roots of a degree 4 homogenous polynomial over  $\mathbb{P}^2$  is known as a **quartic**.

## Theorem (Zeuthen, 1874 [Zeu74])

Let  $\Delta$  be a smooth quartic over  $\mathbb{R}$ , then  $\Delta(\mathbb{R})$  can be classified into 1 of the 6 following topological types:

- ① No real points
- ② One oval
- ③ Two nested ovals
- ④ Two non-nested ovals
- ⑤ Three ovals
- ⑥ Four ovals



# Example: Four Ovals

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

## Main Result

Main Theorem

Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci


Empty Real Loci

Irrational One Oval  
Example

The homogeneous equation defines a smooth quartic whose real component has 4 ovals:

$$0 = -11u^4 - 5u^2v^2 - 2v^4 + 11u^2w^2 + 4v^2w^2 - 3w^4$$

The real components on the chart ( $w \neq 0$ )





# Example: Two Nested Ovals

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

## Main Result

Main Theorem

Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci

Empty Real Loci


Irrational One Oval

Example

This homogeneous equation defines a smooth quartic whose real component has 2 nested ovals:

$$0 = -3u^4 - \frac{7}{10}u^2v^2 - \frac{169}{400}v^4 + \frac{67}{6}u^2w^2 + \frac{949}{240}v^2w^2 - \frac{121}{576}w^4$$

The real components on the chart ( $w \neq 0$ ):





# Overview of Rationality

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

Main Result

Main Theorem

Rational Conic Bundles

Irrational Conic  
Bundles

Disconnected Real Loci

Empty Real Loci

Irrational One Oval  
Example

- A variety  $X$  is  **$k$ -rational** over a field  $k$  if there exists non-empty open sets  $U \subset X$  and  $V \subset \mathbb{P}_k^{\dim X}$  such that  $U$  and  $V$  are isomorphic over  $k$ .
- If  $X$  is not rational over  $k$ , we say that  $X$  is **irrational** over  $k$ .

There are two relevant facts about rationality:

- **Lang–Nishimura Lemma:** If  $X$  is a projective  $k$ -rational variety, then  $X(k)$  is non-empty.
- **General Topological Fact:** If  $X$  is a smooth projective  $\mathbb{R}$ -rational variety, then  $X(\mathbb{R})$  is connected.

In particular,

$Y_{\tilde{\Delta}/\Delta}$  is  $\mathbb{R}$ -rational  $\implies Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  is non-empty and connected



# Criterion for Rationality

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

## Main Result

Main Theorem  
Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

The  $\mathbb{C}$ -rationality of  $Y_{\tilde{\Delta}/\Delta}$  is quite clear:

**Theorem (Iskovskikh, 1987 [Isk87])**

$Y_{\tilde{\Delta}/\Delta}$  is always  $\mathbb{C}$ -rational

The  $\mathbb{R}$ -rationality of  $Y_{\tilde{\Delta}/\Delta}$  is more complicated, but the work of S. Frei, L. Ji, S. Sankar, B. Viray, and I. Vogt previously gave a very useful criterion:

**Proposition (Proposition 6.1, [FJS<sup>+</sup>])**

Let  $Y_{\tilde{\Delta}/\Delta}$  be defined as before. If  $\tilde{\Delta}(\mathbb{R}) \neq \emptyset$ , then  $Y_{\tilde{\Delta}/\Delta}$  is  $\mathbb{R}$ -rational

In our REU, we instead investigate what happens when  $\tilde{\Delta}(\mathbb{R}) = \emptyset$ .





## Rationality of Real Conic Bundles with Quartic Discriminant Curve

Mattie Ji

### Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

### Main Result

Main Theorem  
Rational Conic Bundles

### Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

- 1 Background
  - Plane Conics
  - Conic Bundles
  - Real Quartic Plane Curves
  - Rationality
- 2 Main Result
  - Main Theorem
  - Rational Conic Bundles
- 3 Irrational Conic Bundles
  - Disconnected Real Loci
  - Empty Real Loci
  - Irrational One Oval Example



## Theorem:

Let  $\mathcal{CB}_{\emptyset/*}$  denote the set of real conic bundles  $X \rightarrow \mathbb{P}^2$  of the form  $Y_{\tilde{\Delta}/\Delta}$  with smooth  $\Delta$  of topological type  $*$  and smooth  $\tilde{\Delta}$  that has no real points,

- ①  $\mathcal{CB}_{\emptyset/\emptyset}$  contains both rational and irrational members
- ②  $\mathcal{CB}_{\emptyset/1\text{-oval}}$  contains both rational and irrational members<sup>3</sup>
- ③  $\mathcal{CB}_{\emptyset/2}$  non-nested ovals,  $\mathcal{CB}_{\emptyset/2}$  nested ovals, and  $\mathcal{CB}_{\emptyset/3\text{-ovals}}$  contains both rational and irrational members<sup>4</sup>
- ④  $\mathcal{CB}_{\emptyset/4\text{-ovals}}$  contains only rational members<sup>5</sup>.

<sup>3</sup>The case of irrational 1 oval was done in [FJS<sup>+</sup>]

<sup>4</sup>The case of irrational 2 non-nested ovals was done in [FJS<sup>+</sup>]

<sup>5</sup>We proved this based on ideas joint with S. Frei, S. Sankar, B. Viray, and I. Vogt

It turns out that there's another rationality construction for rational examples when  $\tilde{\Delta}(\mathbb{R})$  is empty:


### Theorem (Witt, 1937 [Wit37])

If  $\pi_1 : Y_{\tilde{\Delta}/\Delta} \rightarrow \mathbb{P}_{[t_0:t_1]}^1$  is surjective on real points, then  $Y_{\tilde{\Delta}/\Delta}$  is  $\mathbb{R}$ -rational.

Using this criterion, we can find and check that rational members exist for all topological types of  $\Delta(\mathbb{R})$ .

## Example of Rational One Oval with $\tilde{\Delta}(\mathbb{R}) = \emptyset$

Take  $Q_1 := -u^2 + uv - w^2$ ,  $Q_2 := 3u^2 + uv - v^2 + w^2$ , and  $Q_3 = -u^2 - 2uv - 2w^2$ , then one can verify that  $\tilde{\Delta}(\mathbb{R}) = \emptyset$ ,  $\pi_1$  is surjective on real points, and  $\Delta(\mathbb{R})$  is one oval, as seen on the chart ( $w \neq 0$ ):





## Rationality of Real Conic Bundles with Quartic Discriminant Curve

Mattie Ji

### Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

### Main Result

Main Theorem  
Rational Conic Bundles

### Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

- 1 Background
  - Plane Conics
  - Conic Bundles
  - Real Quartic Plane Curves
  - Rationality
- 2 Main Result
  - Main Theorem
  - Rational Conic Bundles
- 3 Irrational Conic Bundles
  - Disconnected Real Loci
  - Empty Real Loci
  - Irrational One Oval Example



# Obstruction by Disconnected Real Loci

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

## Main Result

Main Theorem

Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci

Empty Real Loci

Irrational One Oval  
Example

- Recall that if  $Y_{\tilde{\Delta}/\Delta}$  is  $\mathbb{R}$ -rational, then  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  is **connected** and non-empty.
- What if we take  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  to be **disconnected**?

The key insight is looking at  $\pi_1(\mathbb{R}) : Y_{\tilde{\Delta}/\Delta}(\mathbb{R}) \rightarrow \mathbb{P}_{[t_0:t_1]}^1(\mathbb{R})$ :  
 $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  is disconnected  $\iff \pi_1(Y_{\tilde{\Delta}/\Delta}(\mathbb{R}))$  is disconnected

## Observation:

Take any point  $[a : b] \in \mathbb{P}_{[t_0:t_1]}^1(\mathbb{R})$ , then fiber of  $[a : b]$  is exactly the solutions satisfying:

$$z^2 = Q_1(u, v, w)(a)^2 + 2Q_2(u, v, w)(ab) + Q_3(u, v, w)(b)^2$$

This equation has a solution if and only if the matrix

$$M_{[a:b]} := \left[ \begin{array}{c|c} M_1 a^2 + 2M_2 ab + M_3 b^2 & 0 \\ \hline 0 & -1 \end{array} \right]$$

is **NOT** negative definite, where  $M_1, M_2, M_3$  are the symmetric matrices associated to  $Q_1, Q_2, Q_3$  respectively.



# Key Observation (Continued)

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

Main Result

Main Theorem  
Rational Conic Bundles

Irrational Conic  
Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

We will denote the signature of a real symmetric matrix  $M$  as  $(p, n)$ , where  $p$  is the number of positive eigenvalues and  $n$  is the number of negative eigenvalues.

Lemma (Adapted from Degtjarev Et al., [DIK00])

Given the setup with  $Y_{\tilde{\Delta}/\Delta}$ , let  $M_{[t_0:t_1]}$  be the matrix defined the same as previously, then the signature of  $M_{[t_0:t_1]}$  can only change by  $\pm 1$  at real solutions of a degree 6 polynomial<sup>6</sup>

$$\Gamma(t) := -\det(M_1 t^2 + 2M_2 t + M_3)$$

Notably, on each interval defined by real points of  $\Gamma(t)$ , the signature of  $M_{[t_0:t_1]}$  stays the same.

---

<sup>6</sup>It is possible that one of the roots is the point at infinity, but we can without loss choose an appropriate basis to avoid this.





# Disconnected Examples

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

## Main Result

Main Theorem  
Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

Now, as long as we can find examples where at least two non-adjacent intervals with signature  $(0, 4)$ , this will produce a disconnected  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$ .

## Irrational Example with $\Delta(\mathbb{R})$ 3 ovals

Define  $Q_1 := -u^2 - v^2 - w^2$ ,  $Q_2 := u^2 + 5v^2 + 9w^2$ , and  $Q_3 := -24v^2 - 80w^2$ . Then we have that

$$\Gamma(t) = t^6 - 30t^5 + 340t^4 - 1800t^3 + 4384t^2 - 3840t$$

with 6 real roots

$$t = 0, 2, 4, 6, 8, 10$$



# Disconnected Examples

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

## Main Result

Main Theorem  
Rational Conic Bundles


## Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

## Example (Continued)

The signatures of  $M_{[t_0:t_1]}$  follow the pattern:

$$(0, 4), (1, 3), (0, 4), (1, 3), (0, 4), (1, 3)$$





# Limitations of the Topological Obstruction

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

## Main Result

Main Theorem  
Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

During the REU, we also showed that

### Proposition:

If  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  is disconnected, then  $\Delta(\mathbb{R})$  is either 2 non-nested ovals, 2 nested ovals, or 3 ovals. More precisely,

- $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  has at most 3 connected components
- If  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  has 2 connected components, then  $\Delta(\mathbb{R})$  is either 2 non-nested ovals or 2 nested ovals
- If  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  has 3 connected components, then  $\Delta(\mathbb{R})$  is 3 ovals.

Each case described above does occur.



# The Case where $Y_{\tilde{\Delta}/\Delta}(\mathbb{R}) = \emptyset$

Rationality of  
Real Conic  
Bundles with  
Quartic  
Discriminant  
Curve

Mattie Ji

## Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

## Main Result

Main Theorem  
Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

- Recall that if  $Y_{\tilde{\Delta}/\Delta}$  is  $\mathbb{R}$ -rational, then  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  is connected and **non-empty**.
- What if we take  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  to be **empty** instead?

## Irrational Example where $\Delta(\mathbb{R}) = \emptyset$

Define  $Q_1 := -2u^2 - 3v^2 - 5w^2$ ,  $Q_2 := u^2 + 2v^2 + 3w^2$ , and  $Q_3 := -2u^2 - 4v^2 - 2w^2$ . Then one can verify that the associated  $\Delta(\mathbb{R})$  and  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R})$  are both empty, hence  $Y_{\tilde{\Delta}/\Delta}$  is irrational over  $\mathbb{R}$ .

This doesn't get us really far, as  $Y_{\tilde{\Delta}/\Delta}(\mathbb{R}) = \emptyset$  implies that  $\Delta(\mathbb{R}) = \emptyset$ .



## Background

Plane Conics  
Conic Bundles  
Real Quartic Plane  
Curves  
Rationality

## Main Result

Main Theorem  
Rational Conic Bundles

## Irrational Conic Bundles

Disconnected Real Loci  
Empty Real Loci  
Irrational One Oval  
Example

## Question:

What about the case for one oval?

S. Frei, L. Ji, S. Sankar, B. Viray, and I. Vogt showed that

## Irrational Example (Theorem 1.3(2), [FJS<sup>+</sup>])

Define  $Q_1 := -u^2 - v^2 - 3w^2$ ,  $Q_2 := 3u^2 + 5v^2$ , and  $Q_3 := -7u^2 - 23v^2 - 12w^2$ , then  $\Delta(\mathbb{R})$  is one oval, and the associated  $Y_{\tilde{\Delta}/\Delta}$  is irrational over any subfield of  $\mathbb{R}$ .

We would like to thank

- Lena Ji for advising this project
- Sarah Frei, Soumya Sankar, Bianca Viray, and Isabel Vogt for helpful discussions and conversations
- János Kollár for the question that motivated this project
- The University of Michigan Mathematics REU Program
- The National Science Foundation (Karen Smith's NSF grant DMS-2101075) for funding this project
- My letter of recommendation writers Nicole Looper and Jungang Li



Nils Bruin.

The arithmetic of Prym varieties in genus 3.  
*Compos. Math.*, 144(2):317–338, 2008.



Aleksandr Degtjarev, Il'ja V. Itenberg, and Viatcheslav Kharlamov.

*Real Enriques surfaces*, pages 211–212.  
Springer, 2000.



Sarah Frei, Lena Ji, Soumya Sankar, Bianca Viray, and Isabel Vogt.

Curve classes on conic bundle threefolds and applications  
to rationality.  
*arXiv e-prints*, page arXiv:2207.07093.



## Rationality of Real Conic Bundles with Quartic Discriminant Curve

Mattie Ji

### Background

Plane Conics

Conic Bundles

Real Quartic Plane  
Curves

Rationality

### Main Result

Main Theorem

Rational Conic Bundles

### Irrational Conic Bundles

Disconnected Real Loci

Empty Real Loci

Irrational One Oval  
Example



V. A. Iskovskikh.

On the rationality problem for conic bundles.

*Duke Math. J.*, 54(2):271–294, 1987.



Ernst Witt.

Theorie der quadratischen Formen in beliebigen Körpern.

*J. Reine Angew. Math.*, 176:31–44, 1937.



Hieronymus Georg Zeuthen.

Sur les différentes formes des courbes planes du quatrième ordre.

*Mathematische Annalen*, 7:410–432, 1874.