

A Brief Introduction to Radiation Transport

M. I. Ortega

Department of Nuclear Engineering
University of California, Berkeley

Lawrence Livermore National Laboratory Mercury Group
Seminar, 2016

Outline

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

1 Preliminaries

- Cross Sections and Particle Distributions
- Particle Interactions

2 The Transport Equation

- Quantities of Interest
- Particle Balance
- Boundary Conditions
- Eigenvalue Problems

3 Numerical Solution of the Transport Equation

- Discretization
- Solution Methods

Nuclear Cross Sections

Neutron Cross Sections

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions

Particle
Interactions

The Transport
Equation

Quantities of
Interest

Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Particle interactions are incorporated through the concept of cross sections. Neutrons have a few interactions of interest in radiation transport: capture, fission, elastic, and inelastic scattering.

- Neutron Cross Sections:

- Absorption:

$$\sigma_a(E) = \sigma_c(E) + \sigma_f(E) \quad (1)$$

- Scattering (elastic and inelastic):

$$\sigma_s(E) = \sigma_n(E) + \sigma_{n'}(E) + \sum_{i=2}^N \sigma_{i,n}(E) \quad (2)$$


Nuclear Cross Sections

Example of Neutron Cross Sections

A Brief Introduction to Radiation Transport

M. I. Ortega

Cross Sections and Particle Distributions



Nuclear Cross Sections

Photon Cross Sections

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions

Particle
Interactions

The Transport
Equation

Quantities of
Interest

Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Photons have three interactions of interest in radiation transport: photoelectric, Compton scattering, pair production.

■ Photon Cross Sections

■ Absorption:

$$\sigma_a(E) = \sigma_{pe}(E) \quad (3)$$

■ Scattering:

$$\sigma_s(E) = \sigma_{cs}(E) + \sigma_{pp}(E) \quad (4)$$

Nuclear Cross Sections

Example of Photon Cross Section

A Brief
Introduction to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections and Particle
Distributions

Particle
Interactions

The Transport
Equation


Quantities of
Interest
Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods



Scattered Particle Distributions

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions

Particle
Interactions

The Transport
Equation

Quantities of
Interest

Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Express probability distribution laws governing scattered particle emission. The scattering cross section tells us the probability a scattered particle will be emitted in some specific direction with some specific energy:

$$\sigma_s(\vec{r}, E \rightarrow E', \hat{\Omega} \cdot \hat{\Omega}') dE' d\hat{\Omega}' \quad (5)$$

Fission Neutron Distributions

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions

Particle
Interactions

The Transport
Equation

Quantities of
Interest

Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Two quantities need to be defined in fission:

- $\nu(E)$ = mean number of fission neutrons produced in a fission caused by a neutron with energy E .
- $\chi(E)dE$ = probability that a fission neutron will have an energy dE about E .

$$\int_0^{\infty} \chi(E)dE = 1 \quad (6)$$

Particle Interactions

Assumptions

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions

Particle
Interactions

The Transport
Equation

Quantities of
Interest

Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

In the derivation of the TE, we assume the following:

- Particles are considered as points.
- Particles travel in straight lines between point collisions (neutral!).
- Particle-particle interactions can be neglected.
- Collisions are instantaneous (some exceptions like delayed neutron emission).
- Material properties are assumed to be isotropic.
- Nuclei properties and material properties are assumed constant and known (can be relaxed).
- Only expected or mean particle density distribution is considered.

Definitions

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions

Particle
Interactions

The Transport
Equation

Quantities of
Interest

Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Angular Flux

Let the particle density distribution in a six-dimensional phase space (space, direction, energy, and time) be $N(\vec{r}, \hat{\Omega}, E, t)$, we define the neutron angular flux as

$$\psi(\vec{r}, \hat{\Omega}, E, t) = vN(\vec{r}, \hat{\Omega}, E, t) \quad (7)$$

which is convenient for the calculation of reaction rates. Usually reaction rates are independent of direction, so we define the scalar flux:

$$\phi(\vec{r}, \hat{\Omega}, E, t) = \int_{4\pi} \psi(\vec{r}, \hat{\Omega}, E, t). \quad (8)$$

The Transport Equation

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

The TE is merely a balance equation (source - losses):

$$\frac{1}{v} \frac{\partial}{\partial t} \psi(\vec{r}, \hat{\Omega}, E, t) + \hat{\Omega} \cdot \nabla \psi(\vec{r}, \hat{\Omega}, E, t) + \sigma(\vec{r}, E) \psi(\vec{r}, \hat{\Omega}, E, t) = q(\vec{r}, \hat{\Omega}, E, t), \quad (9)$$

where $q(\vec{r}, \hat{\Omega}, E, t)$ includes both the external and fission source.

We need to impose boundary and initial conditions!

Boundary Conditions

Our Connection to the Real World

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

- We can impose an incoming angular flux. If no angular flux is imposed, the boundary condition is called vacuum.

$$\psi(\vec{r}, \hat{\Omega}, E, t) = \tilde{\psi}(\vec{r}, \hat{\Omega}, E, t), \quad \hat{\Omega} \cdot \hat{n} < 0, \vec{r} \in \Gamma \quad (10)$$

- Albedo (reflective) boundary condition:

$$\psi(\vec{r}, \hat{\Omega}, E, t) = \alpha(E)\psi(\vec{r}, \hat{\Omega}', E, t), \quad \hat{\Omega} \cdot \hat{n} < 0, \vec{r} \in \Gamma \quad (11)$$

These are usually the two boundary conditions of interest in applications.

Eigenvalue Problems in Radiation Transport

The Criticality Condition

A Brief
Introduction to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

- A problem of special interest is the criticality condition.
- Criticality is defined as a system capable of sustaining a chain reaction without an external source indefinitely. That is, there is a time-independent solution to the problem.
- In most circumstances, a problem under consideration is not exactly critical (no time-independent solution exists). We must reformulate the TE into an eigenvalue problem to determine how subcritical or supercritical the problem really is.
- Reformulation of the TE into an eigenvalue problem is not a unique process (that is, we can create an eigenvalue problem any way we want to).

Eigenvalue Problems in Radiation Transport

The Time-Independent Transport Equation

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

No guarantee non-trivial solution exists for the following problem:

$$\left[\hat{\Omega} \cdot \nabla + \sigma(\vec{r}, E) \right] \psi(\vec{r}, \hat{\Omega}, E) = \int dE' \int d\hat{\Omega}' \sigma_s(\vec{r}, E' \rightarrow E, \hat{\Omega}' \cdot \hat{\Omega}) \psi(\vec{r}, E', \hat{\Omega}') + \chi(E) \int dE' \nu \sigma_f(\vec{r}, E') \int d\hat{\Omega}' \psi(\vec{r}, E', \hat{\Omega}'). \quad (12)$$

We "force" a solution to exist. We consider two formulations: the time-absorption and multiplication eigenvalues.

Eigenvalue Problems in Radiation Transport

The α Eigenvalue

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions

Particle
Interactions

The Transport
Equation

Quantities of
Interest

Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Assume an asymptotic solution to the previous equation in the form

$$\psi(\vec{r}, E, \hat{\Omega}, t) = \psi_{\alpha}(\vec{r}, E, \hat{\Omega})e^{\alpha t}. \quad (13)$$

In general there will be a spectrum of eigenvalues for which there are solutions. At long times only nonnegative solutions will predominate, so we can define criticality as

$$\text{Re } \alpha_0 \begin{cases} > 0 & \textit{supercritical}, \\ = 0 & \textit{critical}, \\ < 0 & \textit{subcritical}. \end{cases} \quad (14)$$

Eigenvalue Problems in Radiation Transport

The Multiplication Eigenvalue

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance

Boundary
Conditions
**Eigenvalue
Problems**

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Assume you can adjust ν , the average number of neutrons emitted in fission, to obtain a time-independent solution. Our fission source term becomes:

$$\frac{\chi(E)}{k} \int dE' \nu \sigma_f(\vec{r}, E') \int d\hat{\Omega}' \psi(\vec{r}, E', \hat{\Omega}'). \quad (15)$$

We can define criticality as

$$k \begin{cases} > 1 & \text{supercritical,} \\ = 1 & \text{critical,} \\ < 1 & \text{subcritical.} \end{cases} \quad (16)$$

Discretizing the Transport Equation in Energy

Energy, Space, and Angle

A Brief
Introduction to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions

Particle
Interactions

The Transport
Equation

Quantities of
Interest

Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Energy Discretization

We divide energy into groups and define a corresponding angular flux in each group,

$$\psi_g(\vec{r}, \hat{\Omega}) = \int_g dE \psi(\vec{r}, \hat{\Omega}, E), \quad \int_g dE = \int_{E_g}^{E_{g-1}} dE \quad (17)$$

such that

$$\int_0^\infty dE' = \sum_{g'=1}^G \int_{g'} dE'. \quad (18)$$

Discretizing the Transport Equation in Energy Energy, Space, and Angle

A Brief
Introduction to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation


Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Spatial Discretization

Derivatives are expressed using either the finite difference or finite element method.



Discretizing the Transport Equation in Energy Energy, Space, and Angle

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation


Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Angular Discretization

The Discrete Ordinate Method is used to discretize angle. The TE equation is solved along specific directions and a quadrature formula is used to reconstruct the angular flux.



The Transport Sweep Algorithm

The TE in Matrix Form

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

We can rewrite the discretized TE equation into matrix form:

$$\mathbf{H}\psi = \chi \mathbf{f}^T \int d\hat{\Omega} \psi + \mathbf{q}^e \quad (19)$$

and recast it into the multiplication eigenvalue formulation:

$$\mathbf{H}\psi = \frac{1}{k} \chi \mathbf{f}^T \int d\hat{\Omega} \psi + \mathbf{q}^e. \quad (20)$$

We define a spatial fission neutron distribution source (scalar):

$$F(\vec{r}) = \mathbf{f}(\vec{r})^T \int d\hat{\Omega} \psi(\vec{r}, \hat{\Omega}). \quad (21)$$

The Transport Sweep Algorithm

The TE in Matrix Form

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

We rewrite our problem in terms of this distribution and doing some manipulations we obtain

$$AF = kF \quad (22)$$

where define the scalar multigroup transport operator as

$$A \equiv \mathbf{f}^T \int d\hat{\Omega} \mathbf{H}^{-1} \chi. \quad (23)$$

This is an eigenvalue problem that can be solved using the power method.

The Power Method

Updating the Eigenvector and Eigenvalue

We determine the eigenvector and eigenvalue using the power method. Assume the problem has a largest eigenvalue, the eigenvalue is positive, and is real and unique, we can obtain the better approximation to the eigenpair using the relationships

$$F_{i+1} = \frac{1}{k_i} A F_i, \quad (24)$$

and

$$k_{i+1} = k_i \frac{\int dV \ w A F_i}{\int dV \ w F_i}. \quad (25)$$

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions

Particle
Interactions

The Transport
Equation

Quantities of
Interest

Particle Balance

Boundary
Conditions

Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization

Solution
Methods

The Power Method

Why Does This Method Work?

Suppose there are distinct eigenvalue problem solutions:

$$AF^\ell = \kappa_\ell F^\ell, \quad \ell = 1, 2, \dots, \quad (26)$$

where

$$k = \kappa_1 > \kappa_2 > \kappa_3 \dots \quad (27)$$

Suppose we guess a linear combination of these solutions

$$F_0 = \sum_{\ell} \alpha_{\ell} F^{\ell}. \quad (28)$$

If we continue to apply on operator (algorithm) say n times, we obtain the following:

$$A^n F_0 = \sum_{\ell} \alpha_{\ell} \kappa_{\ell}^n F^{\ell} = \alpha_1 \kappa_1^n F^1 + \sum_{\ell > 1} \alpha_{\ell} \kappa_{\ell}^n F^{\ell}. \quad (29)$$

The Power Method

Why Does This Method Work?

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Factoring out κ_1^n (which is the value we are interested in) we see that

$$A^n F_0 \rightarrow \alpha_1 \kappa_1^n F^1, \quad n \rightarrow \infty. \quad (30)$$

since $(\kappa_\ell/\kappa_1)^n$ goes to zero as n goes to zero due to the ordering of the eigenvalues.

Inverting \mathbf{H}

We Haven't Solved Anything Yet...

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

Hidden in the math is the fact we need to invert \mathbf{H} . How do we do this?

- Iteratively! Matrix too large and process too expensive to invert directly.
- Gauss-Seidel, Jacobi, many other methods!
- A lot of research in acceleration (preconditioning) of problems. (Me hopefully)

Comparisons to Monte Carlo

Why Not Just Monte Carlo All the Time?

A Brief
Introduction
to Radiation
Transport

M. I. Ortega

Preliminaries

Cross Sections
and Particle
Distributions
Particle
Interactions

The Transport
Equation

Quantities of
Interest
Particle Balance
Boundary
Conditions
Eigenvalue
Problems

Numerical
Solution of
the Transport
Equation

Discretization
Solution
Methods

■ Advantages

- No random error. Error is caused by discretization and data only.
- Fast! Matrix iterative methods are well studied and are less expensive than random sampling.
- Give global solutions (solution defined at every point).
- No need for variance reduction. We are solving an equation and not simulating particles.

■ Disadvantage

- Difficult to represent complex geometries (requires mesh).
- Energy treatment is not continuous (multigroup approximation).
- Difficult to parallelize (can be done however).