




[A1] Copy number status of reportable (and likely copy number-affected) genes for sample IPD2225-D01-P01-A08; LocalApp-estimated sex from the tumor sample: Male; using only Illumina's panel of normals [PON]; no matched normal sample available; no tumor content adjustment applied (assumed tumor cell fraction: 1.0)




[A2] Copy number status of reportable (and likely copy number-affected) genes for sample IPD2225-D01-P01-A08; LocalApp-estimated sex from the tumor sample: Male; using only Illumina's panel of normals [PON]; no matched normal sample available; tumor content adjustment applied (assumed tumor cell fraction: 0.5)




[B2] Copy number status along the genome – bin-wise values for tumor sample IPD2225-D01-P01-A08;
no tumor content adjustment applied (assumed tumor cell fraction: 1.0)




[B3] Copy number status along the genome – bin-wise values for tumor sample IPD2225-D01-P01-A08;
tumor content adjustment applied (assumed tumor cell fraction: 0.5)




[C1] Allelic fractions of assumed somatic and germline variants in tumor sample IPD2225-D01-P01-A08
(variants at genomic sites with read depth < 50 have been excluded)




[C2] Allelic fractions of assumed somatic (SOM) variants in tumor sample IPD2225-D01-P01-A08
(variants at genomic sites with read depth < 50 have been excluded)



[C3] Allelic fractions of assumed germline (GL_DB) variants in tumor sample IPD2225-D01-P01-A08
(variants at genomic sites with read depth < 50 have been excluded)



[C4] Allelic fractions of assumed germline (GL_DB) variants in tumor sample IPD2225-D01-P01-A08
(variants at genomic sites with read depth < 50 have been excluded)



[C6] Tumor RNA sample (IPD2225-R03-P01-A08) allelic fractions of assumed germline (GL_DB) variants identified in tumor sample (IPD2225-D01-P01-A08)
(variants at genomic sites with read depth < 50 have been excluded)

