

Global Seismic Series: Statistical Analysis of Spatiotemporal Clustering in $M \geq 6.5$ Earthquake Catalogs, 1973–2026 CE

© 2026 Yaroslav Marshalkin

Yaroslav Marshalkin

Corresponding author: marshalkin@gmail.com

Telegram: @MRSHLKN | GitHub: github.com/marshalkin-ux/paleoseismic-clustering

Abstract.

We analyze an **analysis catalog of 4,267 unique $M \geq 6.5$ events** (modern window 1973–2026: 2,041). A Baiesi-Paczuski η detector with **great-circle distance** yields **27 algorithmic candidates** in the modern window. **Catalog-calibrated temporal ETAS** (GK mainshocks): mean=27.0, **$p_{\text{ETAS}}=1.0$** — N_{obs} consistent with the null. Under the temporal ETAS model used, no anomalous temporal clustering beyond calibrated ETAS; spatial component not modeled. **Limitation:** spatial component not modeled (Sec. 5.6).

Keywords: global seismicity; seismic series; earthquake clustering; heuristic metric with tectonic hint; Baiesi-Paczuski metric; ETAS validation; Monte Carlo; paleoseismology; remote triggering

1. INTRODUCTION

Large earthquakes do not occur as independent Poisson events. Following the 1992 Landers earthquake (M_w 7.3), Hill et al. [1993] documented remotely triggered seismicity at distances exceeding 1,000 km. Brodsky & Prejean [2006] showed that surface waves can initiate swarms in volcanic systems thousands of kilometres away. The 2004 Sumatra–Andaman earthquake (M_w 9.1) was followed by elevated activity in distant regions [Pollitz et al., 1998; Freed & Lin, 2001].

However, the systematic nature of such correlations remains debated. Michael [2011] found that $M \geq 7$ clustering in 1995–2011 is statistically indistinguishable from random fluctuations. Shearer & Stark [2012] reported no increase in global $M \geq 7$ and $M \geq 8$ rates after the 2004 Sumatra event. Kagan & Jackson [1999] confirmed elevated probability of paired events at short separation without resolving long-range links.

The ETAS model [Ogata, 1988] reproduces regional aftershock clustering but does not encode inter-plate correlations. The Baiesi–Paczuski [2004] metric and Zaliapin–Ben-Zion extensions [2008, 2013] provide objective cluster detection but typically use Euclidean distance, ignoring lithospheric connectivity.

Objective. Test (and if warranted, **falsify**) the hypothesis that physically meaningful multi-regional global series exist, with **primary inference on the modern window (1973–2026)**, using complementary null tests (permutation vs ETAS) and explicit detector liberalness assessment. **Scope.** We combine nearest-neighbor clustering with heuristic tectonic hint and ETAS validation; this complements global rate tests (Michael 2011; Shearer & Stark 2012) with a different η -linkage statistic but does not supersede their conclusions.

§1.1 Research question. Do physically meaningful multi-regional global series exist in $M \geq 6.5$ (1973–2026)? (a) Permutation: $p=0.0001$ rejects Poisson times only. (b) ETAS MLE: mean=27.0, $p_{\text{ETAS}}=1.0$. (c) Global-series hypothesis: **not tested** by temporal-only ETAS; spatial null open. (d) WLS control (App. B): $p=1.0$ — coupling illustration, not primary null.

2. DATA AND METHODS

2.1 Catalog compilation

Magnitude notation. Catalog thresholds and detector gates use **$M \geq 6.5$** (USGS ComCat, predominantly M_w); individual events are cited with catalog type where relevant (e.g. M_w 7.3).

Three catalogs were merged: **USGS ComCat** (1900–2026, primary instrumental); **ISC Bulletin** (relocated hypocenters); and **NOAA NGDC** (historical and paleoseismic records pre-1900, 47 events). Duplicates were merged using ± 30 days and ≤ 50 km tolerance; source priority: $\text{ISC} > \text{USGS} > \text{NOAA}$. After deduplication, the catalog contains **4,267** unique $M \geq 6.5$ events (from **4,418** CSV rows; ~ 151 $M < 6.5$ excluded from clustering). USGS ComCat: **2,088** raw (1973–2026) $\rightarrow$ **2,041** after merge/ISC. GK: 24 aftershocks (2,017/2,041); ZBZ: 1 dependent (2,040/2,041). `quality_score` is

metadata, not an inclusion filter. **Primary analysis set:** events 1900–2026 (4,218); 47 pre-1900 records remain in CSV (provenance) but are excluded from the primary detector pipeline and ETAS calibration window (1973–2026 only) — see Appendix A. **Final catalog:** 4,267 $M \geq 6.5$ events (4,418 CSV rows).

2.1.1 Catalog completeness

Maximum-curvature analysis yields $M_c = 6.55$. Maximum-likelihood b -value from 1,688 events above M_c :

$$b = 0.911 \pm 0.018 \text{ (n = 1,688 events)}$$

2.2 Tectonic heuristic (deprecated)

The Bird (2003) tectonic-path heuristic is **excluded from primary inference** (Appendix); **great-circle only**. Failure to validate shows metric **unsuitability**, not evidence against global series.

$$\eta_{ij} = t_{ij} * r_{ij}^{1.6} * 10^{(-b * m_i)}$$

*$b=1.0$ — deliberate Baiesi & Paczuski (2004) simplification; catalog $b=0.911 \pm 0.018$ for M_c /completeness and MC null only — **not** in the η formula. Equal $N=27$ **does not prove** candidate identity (Jaccard=1.0); upstream clusters at $b=0.911$ **not re-run** (~9.8% label mismatch).*

2.3 Analysis pipeline

Raw catalogs → dedup → exclude $M < 6.5$ (~151 NOAA) → GK declustering (primary) → η NN forest → sliding windows → merge overlapping → series criteria → MC/ETAS/FDR. GK: 24 aftershocks (2,017/2,041); ZBZ: 1 dependent (2,040/2,041) — different algorithms (GK: magnitude-dependent windows; ZBZ: η NN + KDE threshold). At global $M \geq 6.5$ scale ZBZ is permissive (sparse events → high η). GK is sole primary for inference; ZBZ sensitivity only, does not replace GK. If ZBZ were primary, $N=27$ likely unchanged (untested).

Why GK-ZBZ counts differ: not contradictory methods — GK removes short-range fore/aftershocks via fixed windows; ZBZ classifies only 1 event with exceptionally low η at global catalog scale. Under fixed gates GK/ZBZ/none all yield $N=27$ (sensitivity_declustering.json); different mainshock labels (24 vs 1) could matter for cluster analysis.

Algorithm specification (§3.4.1). GK WINDOWS, $T(M)/R(M)$ interpolation, magnitude-descending; aftershocks $[0, T]$, foreshocks $[-T/2, 0]$; haversine. find_nearest_neighbor: causal argmin η , $b=1.0$, GC km. identify_clusters: Union-Find, η_0 KDE. global_series: used[] mask, mean GC > 1500 km; merge 142→47 across epochs.

2.4 Clustering and detector criteria

1. GK declustering (primary) → mainshocks for η NN forest.
2. η NN forest: $b=1.0$, $r^{1.6}$; tectonic Bird 2003 ($1.5\times$ GC fallback).
3. Sliding windows 1, 2, 5 yr (1-yr step); merge 142→47.
4. **Detector gate:** $N\geq 4$; $M\geq 6.5$; mean pairwise GC>1500 km.
5. Flinn-Engdahl zone count — diagnostic only (not admission criterion).

2.5 Primary ETAS null

Primary ETAS null uses catalog-calibrated temporal MLE on GK mainshocks (calibrate_etas_mle.py). WLS calibration scripts are Appendix B reproducibility only.

2.6 Statistical validation

Permutation statement: $p=0.0001$ rejects **Poisson event times only** (Ogata 1988); does not confirm global series.

Permutation test: $n = 10,000$, $p \leq 0.0001$, $z = -6.17$ (modern). **ETAS (primary MLE):** mean = 27.0, $p_{\text{ETAS}} = 1.0$, $N = 27$. Interpretation — Sec. 4. GK mainshocks: $N = 27$. BH post-hoc — not discovery.

3. RESULTS

3.1 Identified series

Full historical analysis yields **47 detector candidates** (not validated global series): 27 modern ($p \leq 0.0001$), 15 early instrumental ($p = 0.007$; pre-1960 incompleteness caveat), 5 historical candidates (**not significant**, $p = 0.46$; 47 $M\geq 6.5$ events pre-1900). 142 cluster candidates before filtering. *Interpretation — Sec. 4.*

Serie s ID	Ev ent s	Re gio ns	M ma x	Years	Mean lat°	Mean lon°	Notes
1905-1910	193	43	8.8	1905-1910	-60...72	-180...180	Early instrumental; incomplete before ~1960
S047	53	5	8.0	1982-2024	-21.5...18.9	-175.5...155.2	Western Pacific
S170	46	12	9.1	2002-2023	-14.3...33.8	-113.1...167.3	Sumatra 2004 (M 9.1)
S095	25	4	7.9	1989-2017	-8.1...16.5	120.4...146.4	Western Pacific arc
S116	22	5	8.2	1993-2021	-31.7...10.8	-179.5...179.4	South Pacific

Table 1. Raw detector candidates — NOT validated series; illustrative only.

3.2 Spatial-temporal distribution

Elevated detector candidate frequency (not validated physical series) occurs in 1952-1965 and 2002-2016 (post-Sumatra period). Spatially, clusters concentrate along the circum-Pacific belt (Kamchatka, Kuril Islands, Japan, Tonga, Indonesia). Tectonic diagnostic: median $\Delta\log_{10}\eta = +0.28$ on random pairs (mostly $1.5\times$ GC fallback).

3.3 Consolidated sensitivity table (modern window)

Parameter	Setting	N_series
GC gate	1000 km	27
GC gate	1500 km (baseline)	27
GC gate	2000 km	27
Window	1 yr	53
Window	2 yr (baseline)	27
Window	5 yr	11
Window	10 yr	6
b in η	1.0 (BP 2004)	27
b in η	0.911 (catalog)	27
b overlap (series sets)	Jaccard=1.0; upstream ~9.8%	27
Declustering	GK / ZBZ / none	27 / 27 / 27
min_events (strict)	5 / 6 / 8	27 / 27 / 27
Catalog	GK mainshocks only	27
$\eta_0 \pm 20\%$	not_applied	—

Table 2. *N_series* sensitivity under fixed detector gates (mean GC > 1500 km, $N \geq 4$). Sources: *sensitivity_*.json*.

4. DISCUSSION AND CONCLUSIONS

Temporal ETAS (primary): no anomalous temporal clustering beyond the null (Sec. 3.1: $N_{\text{obs}}=27$, $\text{mean}=27.0$, $p_{\text{ETAS}}=1.0$). Spatial linkage not tested. Permutation rejects Poisson times only. Detector liberal (142 windows). Bird/WLS — supplementary only.

We used temporal ETAS only; the spatial component was not modeled; conclusions are strictly limited to temporal clustering. Rejecting global series as a **tested** null would require spatial ETAS (Ogata 1998); we **do not** claim that rejection here.

Limitation	Affected step	Impact on main conclusion
η_0 unverified at global scale	GK/ZBZ identify_clusters	KDE valley $\approx 7.1 \times 10^{-6}$; global_series skips $\eta_0 \rightarrow N=27$; mis-specified η_0 shifts GK/ZBZ labels only
$b = 1.0$ vs 0.911 in η	η upstream	$N_{\text{series}} = 27$ both; cluster labels not re-run
No spatial Ogata MLE	ETAS null	Temporal MLE primary; spatial MLE future work
142 windows + merge	Detector	Main source of liberalness
Calibrated $p_{\text{ETAS}} = 1.0$	ETAS test	Consistent with null; supports negative conclusion

Table 3. Limitation $\rightarrow$ affected step $\rightarrow$ impact on main conclusion (Sec. 5.6).

Impact analysis. $N = 27$ is computed by global_series() without η_0 filtering; $b = 1.0$ vs 0.911 leaves $N_{\text{series}} = 27$ under fixed gates, but upstream clusters at $b = 0.911$ were not re-verified. 142 sliding windows dominate detector liberalness.

APPENDIX A. PRE-1900 NOAA RECORDS

Forty-seven fragmentary paleoseismic/historical $M \geq 6.5$ records from NOAA NGDC are retained in data/processed/unified_catalog_full.csv for provenance (not removed). These 47 events are **excluded from the primary detector pipeline and ETAS calibration window**: pipeline_v2.py and calibrate_etas.py use the modern catalog 1973–2026 only ($N=2,041$). Epoch-stratified counts include pre-1900 descriptively via run_full_historical_analysis.py but do not enter primary significance claims.

APPENDIX B. CATALOG-MATCHED WLS NEGATIVE CONTROL

Reproducibility negative control only — not inference. Catalog-matched WLS (results/etas_calibration.json: $\mu \approx 0.103$, $K \approx 0.495$) yields mean=27.0, $p_{\text{ETAS}}=1.0$ ($n=1000$; multiseed stable). Illustrates **detector-calibration coupling**; **not** the primary null.

Component	Method
μ	GK mainshocks / T (closed form)
c, p	Omori MLE, Nelder–Mead on 24 delays
K, α	WLS (numpy.linalg.lstsq) on same 24 GK aftershocks

DATA AND CODE AVAILABILITY

Data and code: github.com/marshalkin-ux/paleoseismic-clustering (docs/data_availability.md). External DOI deposition (Zenodo) deferred — GitHub only.

REFERENCES

1. Baiesi M., Paczuski M. (2004). Scale-free networks of earthquakes and aftershocks. *Phys. Rev. E*, 69, 066106.
2. Zaliapin I. et al. (2008). Clustering analysis of seismicity and aftershock identification. *Phys. Rev. Lett.*, 101, 018501.
3. Zaliapin I., Ben-Zion Y. (2013). Earthquake clusters in southern California I. *J. Geophys. Res.*, 118, 2847–2864.
4. Bird P. (2003). An updated digital model of plate boundaries. *Geochem. Geophys. Geosyst.*, 4(3), 1027.
5. Hill D.P. et al. (1993). Seismicity remotely triggered by the magnitude 7.3 Landers earthquake. *Science*, 260, 1617–1623.
6. Brodsky E.E., Prejean S.G. (2006). New constraints on mechanisms of remotely triggered seismicity. *J. Geophys. Res.*, 111, B06312.
7. Pollitz F.F. et al. (1998). Viscosity of oceanic asthenosphere inferred from remote triggering. *Science*, 280, 1245–1249.
8. King G.C.P. et al. (1994). Static stress changes and the triggering of earthquakes. *BSSA*, 84, 935–953.
9. Stein R.S. (1999). The role of stress transfer in earthquake occurrence. *Nature*, 402, 605–609.
10. Michael A.J. (2011). Random variability explains apparent global clustering of large earthquakes. *Geophys. Res. Lett.*, 38, L21301.
11. Shearer P.M., Stark P.B. (2012). Global risk of big earthquakes has not recently increased. *PNAS*, 109(3), 717–721.
12. Ogata Y. (1988). Statistical models for earthquake occurrences and residual analysis. *J. Amer. Stat. Assoc.*, 83, 9–27.
13. Benjamini Y., Hochberg Y. (1995). Controlling the false discovery rate. *J. Roy. Stat. Soc. B*, 57(1), 289–300.
14. Gardner J.K., Knopoff L. (1974). Is the sequence of earthquakes in Southern California Poissonian? *BSSA*, 64, 1363–1367.
15. Woessner J., Wiemer S. (2005). Assessing the quality of earthquake catalogues. *BSSA*, 95, 684–698.
16. Young J.B. et al. (1996). The Flinn-Engdahl regionalization scheme: the 1995 revision. *Phys. Earth Planet. Int.*, 96, 223–297.

Acknowledgments. The author acknowledges USGS, NOAA NGDC, and the ISC for maintaining open seismic catalogs. This work received no targeted funding.