

Global Seismic Series: Statistical Analysis of Spatiotemporal Clustering in $M \geq 6.5$ Earthquake Catalogs, 1973–2026 CE

© 2026 Yaroslav Marshalkin

Yaroslav Marshalkin

Corresponding author: marshalkin@gmail.com

Telegram: @MRSHLKN | GitHub: github.com/marshalkin-ux/paleoseismic-clustering

Abstract.

We analyse an analysis catalog of 4,267 unique $M \geq 6.5$ events (modern window 1973–2026: 2,041 events; provenance: 4,418 CSV rows, ~151 NOAA $M < 6.5$ excluded from clustering) using the Baiesi-Paczuski metric η with tectonic-path distance (Bird 2003). The detector yields 47 algorithmic candidates (27 modern); historical NOAA records ($n=47$) reported in Appendix A only. Primary ETAS null (literature H&S 2003, decoupled): $\text{mean} \approx 15.4$, $p_{\text{ETAS}} \leq 0.001$ — exceeds local aftershock expectation; not teleseismic proof (Sec. 5.4). GK mainshocks: $N=27$ unchanged. Global-series hypothesis not supported (Sec. 5.4–5.6). Permutation $p=0.0001$ (1/10,001) rejects temporal Poisson null only. Limitations — Sec. 5.6.

Keywords: global seismicity; seismic series; earthquake clustering; heuristic metric with tectonic hint; Baiesi-Paczuski metric; ETAS validation; Monte Carlo; paleoseismology; remote triggering

1. INTRODUCTION

Large earthquakes do not occur as independent Poisson events. Following the 1992 Landers earthquake (M_w 7.3), Hill et al. [1993] documented remotely triggered seismicity at distances exceeding 1,000 km. Brodsky & Prejean [2006] showed that surface waves can initiate swarms in volcanic systems thousands of kilometres away. The 2004 Sumatra–Andaman earthquake (M_w 9.1) was followed by elevated activity in distant regions [Pollitz et al., 1998; Freed & Lin, 2001].

However, the systematic nature of such correlations remains debated. Michael [2011] found that $M \geq 7$ clustering in 1995–2011 is statistically indistinguishable from random fluctuations. Shearer & Stark [2012] reported no increase in global $M \geq 7$ and $M \geq 8$ rates after the 2004 Sumatra event. Kagan & Jackson [1999] confirmed elevated probability of paired events at short separation without resolving long-range links.

The ETAS model [Ogata, 1988] reproduces regional aftershock clustering but does not encode inter-plate correlations. The Baiesi–Paczuski [2004] metric and Zaliapin–Ben-Zion extensions [2008, 2013] provide objective cluster detection but typically use Euclidean distance, ignoring lithospheric connectivity.

Objective. Test (and if warranted, falsify) the hypothesis that physically meaningful multi-regional global series exist, with primary inference on the modern window (1973–2026), using complementary null tests (permutation vs ETAS) and explicit detector liberalness assessment. Scope. We combine nearest-neighbor clustering with heuristic tectonic hint and ETAS validation; this complements global rate tests (Michael 2011; Shearer & Stark 2012) with a different η -linkage statistic but does not supersede their conclusions.

2. DATA AND METHODS

2.1 Catalog compilation

Three catalogs were merged: USGS ComCat (1900–2026, primary instrumental); ISC Bulletin (relocated hypocenters); and NOAA NGDC (historical and paleoseismic records pre-1900, 47 events). Duplicates were merged using ± 30 days and ≤ 50 km tolerance; source priority: ISC > USGS > NOAA. After deduplication, the catalog contains 4,267 unique $M \geq 6.5$ events (from 4,418 CSV rows; ~ 151 $M < 6.5$ excluded from clustering). USGS ComCat: 2,088 raw (1973–2026) $\rightarrow$ 2,041 after merge/ISC. GK: 24 aftershocks (2,017/2,041); ZBZ: 1 dependent (2,040/2,041). `quality_score` is metadata, not an inclusion filter. Primary analysis set: events 1900–2026 (4,218); 47 pre-1900 records remain in CSV (provenance) but are excluded from the primary detector pipeline and ETAS calibration window (1973–2026 only) — see Appendix A. Final catalog: 4,267 $M \geq 6.5$ events (4,418 CSV rows).

2.1.1 Catalog completeness

Maximum-curvature analysis yields $M_c = 6.55$. Maximum-likelihood b -value from 1,688 events above M_c :

$$b = 0.911 \pm 0.018 \quad (n = 1,688 \text{ events})$$

2.2 Tectonic heuristic (deprecated)

The Bird (2003) tectonic-path heuristic is a deprecated diagnostic only (Appendix); primary analysis uses great-circle distance only.

$$\eta_{ij} = t_{ij} * r_{ij}^{1.6} * 10^{(-b * m_i)}$$

b=1.0 — deliberate Baiesi & Paczuski (2004) simplification; catalog $b=0.911 \pm 0.018$ for M_c /completeness and MC null only — not in the η formula. Zaliapin (2008): $D \approx b - \eta$, η_0 , cluster shifts not tested.

2.3 Analysis pipeline

Raw catalogs → dedup → exclude $M < 6.5$ (~151 NOAA) → GK declustering (primary) → η NN forest → sliding windows → merge overlapping → series criteria → MC/ETAS/FDR. GK: 24 aftershocks (2,017/2,041); ZBZ: 1 dependent (2,040/2,041) — different algorithms (GK: magnitude-dependent windows; ZBZ: η NN + KDE threshold). At global $M \geq 6.5$ scale ZBZ is permissive (sparse events → high η). GK is sole primary for inference; ZBZ sensitivity only, does not replace GK. If ZBZ were primary, $N=27$ likely unchanged (untested).

Why GK-ZBZ counts differ: not contradictory methods — GK removes short-range fore/aftershocks via fixed windows; ZBZ classifies only 1 event with exceptionally low η at global catalog scale. Under fixed gates GK/ZBZ/none all yield $N=27$ (sensitivity_declustering.json); different mainshock labels (24 vs 1) could matter for cluster analysis.

2.4 Clustering and detector criteria

1. GK declustering (primary) → mainshocks for η NN forest.
2. η NN forest: $b=1.0$, $r^{1.6}$; tectonic Bird 2003 ($1.5 \times$ GC fallback).
3. Sliding windows 1, 2, 5 yr (1-yr step); merge 142→47.
4. Detector gate: $N \geq 4$; $M \geq 6.5$; mean pairwise GC > 1500 km.
5. Flinn-Engdahl zone count — diagnostic only (not admission criterion).

2.5 Primary ETAS null

Primary ETAS null uses literature H&S 2003 parameters, decoupled from detector output. Catalog calibration scripts are Appendix reproducibility only.

2.6 Statistical validation

Permutation test: $n = 10,000$, $p \leq 0.0001$, $z = -6.17$ (modern). ETAS validation (primary): literature H&S 2003: mean ≈ 15.4 , $p_{\text{ETAS}} \leq 0.001$ — $N_{\text{obs}} = 27$ exceeds local aftershock expectation; not teleseismic proof. GK mainshocks only: $N = 27$

unchanged. BH post-hoc on $N = 47$ — not discovery.

3. RESULTS

3.1 Identified series

Full historical analysis yields 47 detector candidates (not validated global series): 27 modern ($p \leq 0.0001$), 15 early instrumental ($p = 0.007$; pre-1960 incompleteness caveat), 5 historical candidates (not significant, $p = 0.46$; 47 $M \geq 6.5$ events pre-1900). 142 cluster candidates before filtering.

ID	N	Reg.	M max	Period	lat	lon	Notes
1905-1910	193	43	8.8	1905-1910	-60..72	-180..180	Early instrumental; incomplete before ~1960
S047	53	5	8.0	1982-2024	-21.5..18.9	-175.5..155.2	Western Pacific
S170	46	12	9.1	2002-2023	-14.3..33.8	-113.1..167.3	Sumatra 2004 (M 9.1)
S095	25	4	7.9	1989-2017	-8.1..16.5	120.4..146.4	Western Pacific arc
S116	22	5	8.2	1993-2021	-31.7..10.8	-179.5..179.4	South Pacific

Table 1. Top-5 detector candidates (not ETAS-validated physical series).

3.2 Spatial-temporal distribution

Elevated detector candidate frequency (not validated physical series) occurs in 1952–1965 and 2002–2016 (post-Sumatra period). Spatially, clusters concentrate along the circum-Pacific belt (Kamchatka, Kuril Islands, Japan, Tonga, Indonesia). Tectonic diagnostic: median $\Delta \log_{10} \eta = +0.28$ on random pairs (mostly $1.5\times$ GC fallback).

4. DISCUSSION

The detector is liberal: under literature ETAS (mean ≈ 15.4 , $p_{\text{ETAS}} \leq 0.001$) N_{obs} exceeds local-clustering null. No physical mechanism; tectonic metric failed (98% GC fallback). Compatible with Michael (2011) and Shearer & Stark (2012).

5. CONCLUSIONS

The detector is liberal: literature ETAS yields mean ≈ 15.4 , $p_{\text{ETAS}} \leq 0.001$. Global-series hypothesis not supported (Sec. 5.4–5.5). Permutation rejects Poisson times only (Ogata 1988).

1. Heuristic with tectonic hint: 98% GC fallback — failed hypothesis test.

2. 47 detector candidates indistinguishable from ETAS null; Δ CFS/dynamic stress — future work only.

Future work / supplement: Static Δ CFS and dynamic stress for S170, S047, S095; full ETAS MLE and multi-seed robustness.

APPENDIX A. PRE-1900 NOAA RECORDS

Forty-seven fragmentary paleoseismic/historical $M \geq 6.5$ records from NOAA NGDC are retained in `data/processed/unified_catalog_full.csv` for provenance (not removed). These 47 events are excluded from the primary detector pipeline and ETAS calibration window: `pipeline_v2.py` and `calibrate_etas.py` use the modern catalog 1973–2026 only ($N=2,041$). Epoch-stratified counts include pre-1900 descriptively via `run_full_historical_analysis.py` but do not enter primary significance claims.

DATA AND CODE AVAILABILITY

Data and code: github.com/marshalkin-ux/paleoseismic-clustering (`docs/data_availability.md`). External DOI deposition (Zenodo) deferred — GitHub only.

REFERENCES

1. Baiesi M., Paczuski M. (2004). Scale-free networks of earthquakes and aftershocks. *Phys. Rev. E*, 69, 066106.
2. Zaliapin I. et al. (2008). Clustering analysis of seismicity and aftershock identification. *Phys. Rev. Lett.*, 101, 018501.
3. Zaliapin I., Ben-Zion Y. (2013). Earthquake clusters in southern California I. *J. Geophys. Res.*, 118, 2847–2864.
4. Bird P. (2003). An updated digital model of plate boundaries. *Geochem. Geophys. Geosyst.*, 4(3), 1027.
5. Hill D.P. et al. (1993). Seismicity remotely triggered by the magnitude 7.3 Landers earthquake. *Science*, 260, 1617–1623.
6. Brodsky E.E., Prejean S.G. (2006). New constraints on mechanisms of remotely triggered seismicity. *J. Geophys. Res.*, 111, B06312.
7. Pollitz F.F. et al. (1998). Viscosity of oceanic asthenosphere inferred from remote triggering. *Science*, 280, 1245–1249.
8. King G.C.P. et al. (1994). Static stress changes and the triggering of earthquakes. *BSSA*, 84, 935–953.
9. Stein R.S. (1999). The role of stress transfer in earthquake occurrence. *Nature*, 402, 605–609.
10. Michael A.J. (2011). Random variability explains apparent global clustering of large earthquakes. *Geophys. Res. Lett.*, 38, L21301.
11. Shearer P.M., Stark P.B. (2012). Global risk of big earthquakes has not recently increased. *PNAS*, 109(3), 717–721.
12. Ogata Y. (1988). Statistical models for earthquake occurrences and residual analysis. *J. Amer. Stat. Assoc.*, 83, 9–27.
13. Benjamini Y., Hochberg Y. (1995). Controlling the false discovery rate. *J. Roy. Stat. Soc. B*, 57(1), 289–300.

14. Gardner J.K., Knopoff L. (1974). Is the sequence of earthquakes in Southern California Poissonian? BSSA, 64, 1363–1367.
15. Woessner J., Wiemer S. (2005). Assessing the quality of earthquake catalogues. BSSA, 95, 684–698.
16. Young J.B. et al. (1996). The Flinn–Engdahl regionalization scheme: the 1995 revision. Phys. Earth Planet. Int., 96, 223–297.

Acknowledgments. The author acknowledges USGS, NOAA NGDC, and the ISC for maintaining open seismic catalogs. This work received no targeted funding.