

Global Seismic Series: Statistical Analysis of Spatiotemporal Clustering in $M \geq 6.5$ Earthquake Catalogs, 1973-2026 CE

Temporal ETAS null and hold-out validation

© 2026 Yaroslav Marshalkin

Yaroslav Marshalkin

Corresponding author: marshalkin@gmail.com

Telegram: @MRSHLKN | GitHub: github.com/marshalkin-ux/paleoseismic-clustering

Abstract.

We analyze an **analysis catalog of 4,267 unique $M \geq 6.5$ events** (modern window 1973-2026: 2,041). Baiesi-Paczuski η detector yields **27 algorithmic candidates**. **Catalog-calibrated temporal ETAS: $p_{ETAS}=1.0$** (in-sample null; full numbers — Sec. 4.1). Spatial component not modeled. Pre-1900 NOAA — Supplementary S3.

Keywords: global seismicity; seismic series; earthquake clustering; heuristic metric with tectonic hint; Baiesi-Paczuski metric; ETAS validation; Monte Carlo; paleoseismology; remote triggering

1. INTRODUCTION

Large earthquakes do not occur as independent Poisson events. Following the 1992 Landers earthquake (M_w 7.3), Hill et al. [1993] documented remotely triggered seismicity at distances exceeding 1,000 km. Brodsky & Prejean [2006] showed that surface waves can initiate swarms in volcanic systems thousands of kilometres away. The 2004 Sumatra–Andaman earthquake (M_w 9.1) was followed by elevated activity in distant regions [Pollitz et al., 1998; Freed & Lin, 2001].

However, the systematic nature of such correlations remains debated. Michael [2011] found that $M \geq 7$ clustering in 1995–2011 is statistically indistinguishable from random fluctuations. Shearer & Stark [2012] reported no increase in global $M \geq 7$ and $M \geq 8$ rates after the 2004 Sumatra event. Kagan & Jackson [1999] confirmed elevated probability of paired events at short separation without resolving long-range links.

The ETAS model [Ogata, 1988] reproduces regional aftershock clustering but does not encode inter-plate correlations. The Baiesi–Paczuski [2004] metric and Zaliapin–Ben-Zion extensions [2008, 2013] provide objective cluster detection but typically use Euclidean distance, ignoring lithospheric connectivity.

Objective. Test (and if warranted, **falsify**) the hypothesis that physically meaningful multi-regional global series exist, with **primary inference on the modern window (1973–2026)**, using complementary null tests (permutation vs ETAS) and explicit detector liberalness assessment. **Scope.** We combine nearest-neighbor clustering with heuristic tectonic hint and ETAS validation; this complements global rate tests (Michael 2011; Shearer & Stark 2012) with a different η -linkage statistic but does not supersede their conclusions.

§1.1 Research question. Do physically meaningful multi-regional global series exist in $M \geq 6.5$ (1973–2026)? (a) Permutation → see Sec. 4.1. (b) ETAS MLE → see Sec. 4.1. (c) Global-series hypothesis: **not tested** by temporal-only ETAS. (d) WLS (Supplementary S2): coupling illustration, not primary null.

2. DATA AND METHODS

2.1 Catalog compilation

Magnitude notation. Catalog thresholds and detector gates use **$M \geq 6.5$** (USGS ComCat, predominantly M_w); individual events are cited with catalog type where relevant (e.g. M_w 7.3).

Three catalogs were merged: **USGS ComCat** (1900–2026, primary instrumental); **ISC Bulletin** (relocated hypocenters); and **NOAA NGDC** (historical and paleoseismic records pre-1900, 47 events). Duplicates were merged using ± 30 days and ≤ 50 km tolerance; source priority: ISC > USGS > NOAA. After deduplication, the catalog contains **4,267** unique $M \geq 6.5$ events (from **4,418** CSV rows; ~ 151 $M < 6.5$ excluded from clustering). USGS ComCat: **2,088** raw (1973–2026) → **2,041** after merge/ISC. GK: 24 aftershocks (2,017/2,041); ZBZ: 1 dependent (2,040/2,041). `quality_score` is metadata, not an inclusion filter. **Primary analysis set:** events 1900–2026 (4,218); 47

pre-1900 records remain in CSV (provenance) but are excluded from the primary detector pipeline and ETAS calibration window (1973–2026 only) — see Supplementary S3. **Final catalog:** 4,267 $M \geq 6.5$ events (4,418 CSV rows).

2.1.1 Catalog completeness

Maximum-curvature analysis yields $M_c = 6.55$. Maximum-likelihood b -value from 1,688 events above M_c :

$$b = 0.911 \pm 0.018 \text{ (n = 1,688 events)}$$

2.2 Tectonic heuristic (deprecated)

The Bird (2003) tectonic-path heuristic is **excluded from the primary pipeline; no synthetic benchmark** in this work; comparison deferred to supplementary material and future work. Primary analysis uses **great-circle distance only** (see paper/supplementary.md §S1).

$$\eta_{ij} = t_{ij} * r_{ij}^{1.6} * 10^{(-b * m_i)}$$

*b=1.0 — deliberate Baiesi & Paczuski (2004) simplification; catalog $b=0.911 \pm 0.018$ for M_c /completeness and MC null only — **not** in the η formula. Equal $N=27$ **does not prove** candidate identity (Jaccard=1.0); upstream clusters at $b=0.911$ **not re-run** (~9.8% label mismatch).*

2.3 Analysis pipeline

Raw catalogs → dedup → exclude $M < 6.5$ (~151 NOAA) → GK declustering (primary) → η NN forest → sliding windows → merge overlapping → series criteria → MC/ETAS/FDR. GK: 24 aftershocks (2,017/2,041); ZBZ: 1 dependent (2,040/2,041) — different algorithms (GK: magnitude-dependent windows; ZBZ: η NN + KDE threshold). At global $M \geq 6.5$ scale ZBZ is permissive (sparse events → high η). GK is sole primary for inference; ZBZ sensitivity only, does not replace GK. If ZBZ were primary, $N=27$ likely unchanged (untested).

Why GK-ZBZ counts differ: not contradictory methods — GK removes short-range fore/aftershocks via fixed windows; ZBZ classifies only 1 event with exceptionally low η at global catalog scale. Under fixed gates GK/ZBZ/none all yield $N=27$ (sensitivity_declustering.json); different mainshock labels (24 vs 1) could matter for cluster analysis.

Algorithm specification (§3.4.1). GK WINDOWS, $T(M)/R(M)$ interpolation, magnitude-descending; aftershocks $[0, T]$, foreshocks $[-T/2, 0]$; haversine. find_nearest_neighbor: causal argmin η , $b=1.0$, GC km. identify_clusters: Union-Find, η_0 KDE. global_series: used[] mask, mean GC > 1500 km; merge 142→47 across epochs.

2.4 Clustering and detector criteria

1. GK declustering (primary) → mainshocks for η NN forest.
2. η NN forest: $b=1.0$, $r^{1.6}$; tectonic Bird 2003 ($1.5\times$ GC fallback).
3. Sliding windows 1, 2, 5 yr (1-yr step); merge 142→47.
4. **Detector gate:** $N\geq 4$; $M\geq 6.5$; mean pairwise GC>1500 km.
5. Flinn-Engdahl zone count — diagnostic only (not admission criterion).

2.5 Primary ETAS null

Primary ETAS null uses catalog-calibrated temporal MLE on GK mainshocks (calibrate_etas_mle.py). Hold-out: train 1973–2000, hold-out 2001–2026 (calibrate_etas_holdout.py; $n=1000$, seed=42). WLS — Supplementary S2 only.

2.6 Statistical validation

Permutation statement: $p=0.0001$ rejects **Poisson event times only** (Ogata 1988); does not confirm global series. ETAS — Sec. 4.1. BH post-hoc — not discovery.

3. RESULTS

3.1 Identified series

Full historical analysis yields **47 detector candidates** (not validated global series): 27 modern ($p \leq 0.0001$), 15 early instrumental ($p = 0.007$; pre-1960 incompleteness caveat), 5 historical candidates (**not significant**, $p = 0.46$; 47 $M\geq 6.5$ events pre-1900). 142 cluster candidates before filtering. *Interpretation — Sec. 4.*

Serie s ID	Ev ent s	Re gio ns	M ma x	Years	Mean lat°	Mean lon°	Notes
1905-1910	193	43	8.8	1905-1910	−60...72	−180...180	Early instrumental; incomplete before ~1960
S047	53	5	8.0	1982-2024	−21.5...18.9	−175.5...155.2	Western Pacific
S170	46	12	9.1	2002-2023	−14.3...33.8	−113.1...167.3	Sumatra 2004 (M 9.1)
S095	25	4	7.9	1989-2017	−8.1...16.5	120.4...146.4	Western Pacific arc
S116	22	5	8.2	1993-2021	−31.7...10.8	−179.5...179.4	South Pacific

Table 1. Raw detector candidates — NOT validated series; illustrative only.

3.2 ETAS validation (Sec. 4.1 canonical)

Split	N_obs	mean	p_ETAS
In-sample (1973–2026)	27	27.0	1.0
Train-calibrated hold-out (2001–2026)	13	13.0	1.0

Permutation: $p = 0.0001$ (1/10,001), $z = -6.17$. Hold-out — partial out-of-time check, not spatial validation. **Multiple testing:** the 27 modern series were **not** FDR-corrected

for the 142-window search; Bonferroni $\alpha/142 \approx 0.00035 > p = 0.0001$. BH post-hoc on $N=47$ — not a discovery claim.

3.3 Spatial-temporal distribution

Elevated detector candidate frequency (not validated physical series) occurs in 1952–1965 and 2002–2016 (post-Sumatra period). Spatially, clusters concentrate along the circum-Pacific belt (Kamchatka, Kuril Islands, Japan, Tonga, Indonesia). Tectonic diagnostic: median $\Delta \log_{10} \eta = +0.28$ on random pairs (mostly $1.5\times$ GC fallback).

3.4 Consolidated sensitivity table (modern window)

Parameter	Setting	N_series
GC gate	1000 km	27
GC gate	1500 km (baseline)	27
GC gate	2000 km	27
Window	1 yr	53
Window	2 yr (baseline)	27
Window	5 yr	11
Window	10 yr	6
b in η	1.0 (BP 2004)	27
b in η	0.911 (catalog)	27
b overlap (series sets)	Jaccard=1.0; upstream $\sim 9.8\%$	27
Declustering	GK / ZBZ / none	27 / 27 / 27
min_events (strict)	5 / 6 / 8	27 / 27 / 27
Catalog	GK mainshocks only	27
$\eta_0 \pm 20\%$	not_applied	—

Table 2. N_{series} sensitivity under fixed detector gates (mean GC > 1500 km, $N \geq 4$). Sources: *sensitivity_*.json*.

4. DISCUSSION AND CONCLUSIONS

Temporal ETAS: No excess candidates beyond catalog-calibrated null (Sec. 3.2). Spatial linkage not tested. Permutation rejects Poisson times only. Bird/WLS/pre-1900 — paper/supplementary.md §S1–S3.

Supplementary material. Bird (2003), WLS negative control, pre-1900 NOAA: paper/supplementary.md (§S1–S3).

Limitation	Affected step	Impact on main conclusion
η_0 unverified at global scale	GK/ZBZ identify_clusters	KDE valley $\approx 7.1 \times 10^{-6}$; global_series skips $\eta_0 \rightarrow N=27$; mis-specified η_0 shifts GK/ZBZ labels only
$b = 1.0$ vs 0.911 in η	η upstream	$N_{\text{series}} = 27$ both; cluster labels not re-run
No spatial Ogata MLE	ETAS null	Temporal MLE primary; spatial MLE future work
142 windows + merge	Detector	Main source of liberalness
Calibrated $p_{\text{ETAS}} = 1.0$	ETAS test	Consistent with null; supports negative conclusion

Table 3. Limitation $\rightarrow$ affected step $\rightarrow$ impact on main conclusion (Sec. 5.6).

Impact analysis. $N = 27$ is computed by global_series() without η_0 filtering; $b = 1.0$ vs 0.911 leaves $N_{\text{series}} = 27$ under fixed gates, but upstream clusters at $b = 0.911$ were not re-verified. 142 sliding windows dominate detector liberalness.

DATA AND CODE AVAILABILITY

Data and code: github.com/marshalkin-ux/paleoseismic-clustering (docs/data_availability.md). External DOI deposition (Zenodo) deferred — GitHub only.

REFERENCES

1. Baiesi M., Paczuski M. (2004). Scale-free networks of earthquakes and aftershocks. Phys. Rev. E, 69, 066106.
2. Zaliapin I. et al. (2008). Clustering analysis of seismicity and aftershock identification. Phys. Rev. Lett., 101, 018501.
3. Zaliapin I., Ben-Zion Y. (2013). Earthquake clusters in southern California I. J. Geophys. Res., 118, 2847–2864.
4. Bird P. (2003). An updated digital model of plate boundaries. Geochem. Geophys. Geosyst., 4(3), 1027.
5. Hill D.P. et al. (1993). Seismicity remotely triggered by the magnitude 7.3 Landers earthquake. Science, 260, 1617–1623.
6. Brodsky E.E., Prejean S.G. (2006). New constraints on mechanisms of remotely triggered seismicity. J. Geophys. Res., 111, B06312.
7. Pollitz F.F. et al. (1998). Viscosity of oceanic asthenosphere inferred from remote triggering. Science, 280, 1245–1249.
8. King G.C.P. et al. (1994). Static stress changes and the triggering of earthquakes. BSSA, 84, 935–953.
9. Stein R.S. (1999). The role of stress transfer in earthquake occurrence. Nature, 402, 605–609.
10. Michael A.J. (2011). Random variability explains apparent global clustering of large earthquakes. Geophys. Res. Lett., 38, L21301.
11. Shearer P.M., Stark P.B. (2012). Global risk of big earthquakes has not recently increased. PNAS, 109(3), 717–721.
12. Ogata Y. (1988). Statistical models for earthquake occurrences and residual analysis. J. Amer. Stat. Assoc., 83, 9–27.

13. Benjamini Y., Hochberg Y. (1995). Controlling the false discovery rate. *J. Roy. Stat. Soc. B*, 57(1), 289–300.
14. Gardner J.K., Knopoff L. (1974). Is the sequence of earthquakes in Southern California Poissonian? *BSSA*, 64, 1363–1367.
15. Woessner J., Wiemer S. (2005). Assessing the quality of earthquake catalogues. *BSSA*, 95, 684–698.
16. Young J.B. et al. (1996). The Flinn-Engdahl regionalization scheme: the 1995 revision. *Phys. Earth Planet. Int.*, 96, 223–297.

Acknowledgments. The author acknowledges USGS, NOAA NGDC, and the ISC for maintaining open seismic catalogs. This work received no targeted funding.