

# Global Seismic Series: Statistical Analysis of Spatiotemporal Clustering in $M \geq 6.5$ Earthquake Catalogs, 1973–2026 CE

© 2026 Yaroslav Marshalkin

---

Yaroslav Marshalkin

Corresponding author: marshalkin@gmail.com

Telegram: @MRSHLKN | GitHub: github.com/marshalkin-ux/paleoseismic-clustering

## Abstract.

*We analyse a merged catalog of 4,267 unique  $M \geq 6.5$  events (from 4,418 CSV rows; ~151 NOAA  $M < 6.5$  excluded from clustering; 47 historical NOAA records pre-1900) using the Baiesi–Paczuski metric  $\eta$  with tectonic-path distance (Bird 2003). 47 global seismic series are identified (27 modern, 15 early, 5 historical candidates). Significance: permutation test ( $n=10,000$ ,  $p \leq 0.0001$ ,  $z = -6.17$ ); ETAS validation ( $n=1000$  catalogs,  $FPR=1000/1000$ , mean 15.4  $\pm$  1.7; max 24;  $p_{ETAS}=0/1000$  for  $N_{obs}=27$ ); FDR (45/47 at  $q=0.05$ ,  $N=47$ ). statistical anomaly, not proof of causal triggering. Largest series: 1905–1910 (193 events, 43 regions); most extensive modern: S170 (46 events, 12 regions,  $M_{max}=9.1$ ).*

Keywords: global seismicity; seismic series; earthquake clustering; tectonic distance; Baiesi–Paczuski metric; ETAS validation; false discovery rate; Monte Carlo; paleoseismology; remote triggering

---

## 1. INTRODUCTION

Large earthquakes do not occur as independent Poisson events. Following the 1992 Landers earthquake ( $M_w$  7.3), Hill et al. [1993] documented remotely triggered seismicity at distances exceeding 1,000 km. Brodsky & Prejean [2006] showed that surface waves can initiate swarms in volcanic systems thousands of kilometres away. The 2004 Sumatra–Andaman earthquake ( $M_w$  9.1) was followed by elevated activity in distant regions [Pollitz et al., 1998; Freed & Lin, 2001].

However, the systematic nature of such correlations remains debated. Michael [2011] found that  $M \geq 7$  clustering in 1995–2011 is statistically indistinguishable from random fluctuations. Shearer & Stark [2012] reported no increase in global  $M \geq 7$  and  $M \geq 8$  rates after the 2004 Sumatra event. Kagan & Jackson [1999] confirmed elevated probability of paired events at short separation without resolving long-range links.

The ETAS model [Ogata, 1988] reproduces regional aftershock clustering but does not encode inter-plate correlations. The Baiesi–Paczuski [2004] metric and Zaliapin–Ben-Zion extensions [2008, 2013] provide objective cluster detection but typically use Euclidean distance, ignoring lithospheric connectivity.

**Objective.** We test the hypothesis that multi-regional seismic series exist in a four-millennium catalog of  $M \geq 6.5$  earthquakes, using an adapted eta metric with tectonic distance along Bird (2003) plate boundaries. **Scope.** We combine nearest-neighbor clustering with tectonic-path distance, ETAS validation, and FDR correction; this complements global rate tests (Michael 2011; Shearer & Stark 2012) with a different eta-linkage statistic but does not supersede their conclusions.

---

## 2. DATA AND METHODS

### 2.1 Catalog compilation

Three catalogs were merged: USGS ComCat (1900–2026, primary instrumental); ISC Bulletin (relocated hypocenters); and NOAA NGDC (historical and paleoseismic records pre-1900, 47 events). Duplicates were merged using  $\pm 30$  days and  $\leq 50$  km tolerance; source priority: ISC > USGS > NOAA. After deduplication, the catalog contains 4,267 unique  $M \geq 6.5$  events (from 4,418 CSV rows;  $\sim 151$   $M < 6.5$  excluded from clustering). USGS ComCat: 2,088 raw (1973–2026)  $\rightarrow$  2,041 after merge/ISC. GK: 24 aftershocks (2,017/2,041); ZBZ: 1 dependent (2,040/2,041). `quality_score` is metadata, not an inclusion filter. Final catalog: 4,267  $M \geq 6.5$  events (4,418 CSV rows).

#### 2.1.1 Catalog completeness

Maximum-curvature analysis yields  $M_c = 6.55$ . Maximum-likelihood b-value from 1,688 events above  $M_c$ :

**$b = 0.911 \pm 0.018$  (n = 1,688 events)**

### 2.2 Tectonic distance and connectivity metric

Tectonic distance  $r_{ij}$  is the shortest path between hypocenters along the Bird (2003) plate-boundary graph (20 key segments). Paths are computed with Dijkstra's algorithm (NetworkX).

If either hypocenter lies >500 km from the nearest boundary node, or no graph path exists,  $r_{ij} = 1.5 \times r_{GC}$  (great-circle Haversine). Fallback audit (analyze\_tectonic\_fallback.py, fig07): 4987 pairs from 500 events; 98.0% GC fallback, 2.0% Dijkstra (4015 snap>500 km, 872 no path, 100 Dijkstra; 95 materially different; examples FE31 Japan, FE35 Philippines). Metric adds value for ~2% boundary-proximal pairs only. Tectonic diagnostic: median  $\Delta \log_{10} \eta = +0.28$ .

$$\eta_{ij} = t_{ij} * r_{ij}^{1.6} * 10^{(-b * m_i)}$$

*b=1.0 per Baiesi & Paczuski (2004) for eta comparability; catalog b=0.911+/-0.018 in Monte Carlo null only.*

### 2.3 Analysis pipeline

Raw catalogs → dedup → exclude  $M < 6.5$  (~151 NOAA) → GK declustering (primary) → eta NN forest → sliding windows → merge overlapping → series criteria → MC/ETAS/FDR. GK: 24 aftershocks (2,017/2,041); ZBZ: 1 dependent (2,040/2,041) — separate sensitivity check, not sequential filter.

### 2.4 Threshold $\eta_0$ , series detection, and ETAS parameters

Automatic  $\eta_0$  from nearest-neighbor eta distribution: KDE valley between bimodal  $\log_{10}(\eta)$  modes (Zaliapin & Ben-Zion 2013); default  $\eta_0 = 10^{\text{median } \log_{10} \eta}$ . Validated against Poisson permutation null ( $n = 10,000$ ).

1. Declustering via Gardner–Knopoff [1974].
2. Nearest-neighbor forest: parent  $i^* = \text{argmin } \eta_{ij}$  subject to  $\eta_{ij} < \eta_0$ .
3. Global series:  $N \geq 4$ ;  $M \geq 6.5$ ;  $\geq 3$  Flinn–Engdahl regions.
4. Sliding windows (1, 2, 5 yr); overlapping groups merged.

ETAS parameters (not catalog-calibrated): Helmstetter & Sornette 2003 defaults ( $\mu=0.008$ ,  $K=0.08$ ,  $\alpha=1.0$ ,  $c=0.005$  d,  $p=1.1$ ); not refit on 2041 events (results/etas\_calibration\_note.md). Optional MLE  $\mu \sim 0.105$  events/day vs default 0.008.

### 2.5 Statistical validation

Permutation test:  $n = 10,000$ ,  $p \leq 0.0001$ ,  $z = -6.17$  (modern). ETAS validation: 1000 catalogs (seed=42); FPR=1000/1000; mean 15.4; max 24;  $p_{\text{ETAS}} \leq 0.001$ . Previously FPR=0/100 was an API bug (min\_regions TypeError). FDR ( $q=0.05$ ): 45/47 significant;  $N=47$  series hypotheses. Declustering: GK 2,017/2,041; ZBZ 2,040/2,041.

## 3. RESULTS

### 3.1 Identified series

Full historical analysis yields 47 global seismic series: 27 modern ( $p \leq 0.0001$ ), 15 early instrumental ( $p = 0.007$ ; pre-1960 incompleteness caveat), 5 historical candidates (not significant,  $p = 0.46$ ; 47  $M \geq 6.5$  events pre-1900). 142 cluster candidates before filtering.

| ID | N | Regions | $M_{\text{max}}$ | Period | Notes |
|----|---|---------|------------------|--------|-------|
|----|---|---------|------------------|--------|-------|

|           |     |    |     |           |                                     |
|-----------|-----|----|-----|-----------|-------------------------------------|
| 1905–1910 | 193 | 43 | 8.8 | 1905–1910 | Largest series; early instrumental  |
| S047      | 53  | 5  | 8.0 | 1982–2024 | Western Pacific subduction corridor |
| S170      | 46  | 12 | 9.1 | 2002–2023 | Sunda belt; Sumatra 2004 (M 9.1)    |
| S095      | 25  | 4  | 7.9 | 1989–2017 | Western Pacific arc                 |
| S116      | 22  | 5  | 8.2 | 1993–2021 | South Pacific multi-arc series      |

Table 1. Top five multi-regional series ( $\geq 3$  Flinn–Engdahl regions).

### 3.2 Spatial–temporal distribution

Elevated series activity occurs in 1952–1965 and 2002–2016 (post-Sumatra period). Spatially, clusters concentrate along the circum-Pacific belt (Kamchatka, Kuril Islands, Japan, Tonga, Indonesia). Tectonic diagnostic: median  $\Delta\log_{10}\eta = +0.28$  on random pairs (mostly 1.5x GC fallback).

## 4. DISCUSSION AND CONCLUSIONS

Statistical anomaly (established): series incompatible with Poisson and local-only ETAS nulls; FDR 45/47. Conclusion about  $\eta$  links and p-values, not causality. Working hypotheses (not claims): viscoelastic mantle coupling (Pollitz 1998), dynamic triggering (Hill 1993; Brodsky 2006), shared tectonic loading (Freed & Lin 2001). Future work / supplement: preliminary Coulomb/dynamic stress tests for S170 did not reach triggering thresholds (repository). ETAS note:  $\text{FPR}=1000/1000$  at seed=42 (mean 15.4 spurious series);  $N_{\text{obs}}=27$  exceeds ETAS max (24). Tectonic audit: 98% GC fallback of 4987 pairs (fig07).

1. 4,267  $M \geq 6.5$  events (4,418 CSV records) contain 47 global series; 27 modern significant at  $p \leq 0.0001$ .
2. ETAS validation (1000 catalogs;  $\text{FPR}=1000/1000$ ,  $p_{\text{ETAS}} \leq 0.001$ ) and FDR (45/47,  $N=47$ ) confirm non-randomness.
3. Largest series: 1905–1910 (193 events, 43 regions,  $M_{\text{max}}=8.8$ ); most spatially extensive modern: S170 (46 events, 12 regions, 2002–2023,  $M_{\text{max}}=9.1$ ).
4. Tectonic path: 2.0% Dijkstra / 98.0% GC fallback (4987 pairs); median  $\Delta\log_{10}\eta = +0.28$ .
5. Interpretive fork: (a) if  $\eta$  links are real — hazard implications without claiming direct triggering; (b) if artifacts — FDR+ETAS remains reproducible null-test.

Future work / supplement: Static Delta CFS and dynamic stress for S170, S047, S095; full ETAS MLE and multi-seed robustness.

## DATA AND CODE AVAILABILITY

Data and code: [github.com/marshalkin-ux/paleoseismic-clustering](https://github.com/marshalkin-ux/paleoseismic-clustering) (docs/data\_availability.md). External DOI deposition (Zenodo) deferred — GitHub only.

## REFERENCES

1. Baiesi M., Paczuski M. (2004). Scale-free networks of earthquakes and aftershocks. *Phys. Rev. E*, 69, 066106.
2. Zaliapin I. et al. (2008). Clustering analysis of seismicity and aftershock identification. *Phys. Rev. Lett.*, 101, 018501.
3. Zaliapin I., Ben-Zion Y. (2013). Earthquake clusters in southern California I. *J. Geophys. Res.*, 118, 2847–2864.
4. Bird P. (2003). An updated digital model of plate boundaries. *Geochem. Geophys. Geosyst.*, 4(3), 1027.
5. Hill D.P. et al. (1993). Seismicity remotely triggered by the magnitude 7.3 Landers earthquake. *Science*, 260, 1617–1623.
6. Brodsky E.E., Prejean S.G. (2006). New constraints on mechanisms of remotely triggered seismicity. *J. Geophys. Res.*, 111, B06312.
7. Pollitz F.F. et al. (1998). Viscosity of oceanic asthenosphere inferred from remote triggering. *Science*, 280, 1245–1249.
8. King G.C.P. et al. (1994). Static stress changes and the triggering of earthquakes. *BSSA*, 84, 935–953.
9. Stein R.S. (1999). The role of stress transfer in earthquake occurrence. *Nature*, 402, 605–609.
10. Michael A.J. (2011). Random variability explains apparent global clustering of large earthquakes. *Geophys. Res. Lett.*, 38, L21301.
11. Shearer P.M., Stark P.B. (2012). Global risk of big earthquakes has not recently increased. *PNAS*, 109(3), 717–721.
12. Ogata Y. (1988). Statistical models for earthquake occurrences and residual analysis. *J. Amer. Stat. Assoc.*, 83, 9–27.
13. Benjamini Y., Hochberg Y. (1995). Controlling the false discovery rate. *J. Roy. Stat. Soc. B*, 57(1), 289–300.
14. Gardner J.K., Knopoff L. (1974). Is the sequence of earthquakes in Southern California Poissonian? *BSSA*, 64, 1363–1367.
15. Woessner J., Wiemer S. (2005). Assessing the quality of earthquake catalogues. *BSSA*, 95, 684–698.
16. Young J.B. et al. (1996). The Flinn–Engdahl regionalization scheme: the 1995 revision. *Phys. Earth Planet. Int.*, 96, 223–297.

**Acknowledgments.** The author acknowledges USGS, NOAA NGDC, and the ISC for maintaining open seismic catalogs. This work received no targeted funding.