

Parking Functions with a Fixed Set of Lucky Cars

Lucy Martinez



Joint work with Pamela E. Harris

Outline


1 Background

2 Motivation

3 Results


Background

Parking Functions




- n cars and a one-way street with n parking lots numbered $1, 2, \dots, n$.

Parking Functions




- n cars and a one-way street with n parking lots numbered $1, 2, \dots, n$.
- Each car i enters the street (one at a time) and has a preferred parking spot a_i .

Parking Functions




- n cars and a one-way street with n parking lots numbered $1, 2, \dots, n$.
- Each car i enters the street (one at a time) and has a preferred parking spot a_i .
- Parking Rule:

Parking Functions



- n cars and a one-way street with n parking lots numbered $1, 2, \dots, n$.
- Each car i enters the street (one at a time) and has a preferred parking spot a_i .
- Parking Rule:
 - Car i tries to park in their favorite spot a_i .

Parking Functions




- n cars and a one-way street with n parking lots numbered $1, 2, \dots, n$.
- Each car i enters the street (one at a time) and has a preferred parking spot a_i .
- Parking Rule:
 - Car i tries to park in their favorite spot a_i .
 - If occupied: the car keeps driving and parks at the next available spot (if any).

Example 1

- Is $\alpha = (3, 1, 1)$ a parking function?


Example 1

- Is $\alpha = (3, 1, 1)$ a parking function?




Example 1

- Is $\alpha = (3, 1, 1)$ a parking function?




Example 1

- Is $\alpha = (3, 1, 1)$ a parking function?



Example 1

- Is $\alpha = (3, 1, 1)$ a parking function?




Example 2

- Is $\alpha = (3, 2, 2)$ a parking function?


Example 2

- Is $\alpha = (3, 2, 2)$ a parking function?




Example 2

- Is $\alpha = (3, 2, 2)$ a parking function?




Example 2

- Is $\alpha = (3, 2, 2)$ a parking function?



Example 2

- Is $\alpha = (3, 2, 2)$ a parking function?



Parking Functions Size

How many parking functions of length n are there? Let $PF(n)$ be the set of parking functions of length n .

Parking Functions Size

How many parking functions of length n are there? Let $PF(n)$ be the set of parking functions of length n .

Permutations on n elements $\{1, \dots, n\}$ are certainly parking functions.

Parking Functions Size

How many parking functions of length n are there? Let $PF(n)$ be the set of parking functions of length n .

Permutations on n elements $\{1, \dots, n\}$ are certainly parking functions.

Let's list some:

- $n = 2$: 11, 21, 12

Parking Functions Size

How many parking functions of length n are there? Let $PF(n)$ be the set of parking functions of length n .

Permutations on n elements $\{1, \dots, n\}$ are certainly parking functions.

Let's list some:

- $n = 2$: 11, 21, 12
- $n = 3$: 111, 211, 121, 112, 122, 221, 212, 113, 131, 311, 231, 123, 132, 213, 312, 321

Parking Functions Size

How many parking functions of length n are there? Let $PF(n)$ be the set of parking functions of length n .

Permutations on n elements $\{1, \dots, n\}$ are certainly parking functions.

Let's list some:

- $n = 2$: 11, 21, 12
- $n = 3$: 111, 211, 121, 112, 122, 221, 212, 113, 131, 311, 231, 123, 132, 213, 312, 321

Theorem (Konheim and Weiss, 1966)

$$|PF(n)| = (n + 1)^{(n-1)}.$$

Outcomes of Parking Functions

Definition

If \mathfrak{S}_n denotes the set of permutations of $[n]$ written in one-line notation, then the outcome of α is

$$\mathcal{O}(\alpha) = \pi_1 \pi_2 \cdots \pi_n,$$

where π_i denotes that car π_i parked in spot i on the street.

Outcomes of Parking Functions

Definition


If \mathfrak{S}_n denotes the set of permutations of $[n]$ written in one-line notation, then the outcome of α is

$$\mathcal{O}(\alpha) = \pi_1 \pi_2 \cdots \pi_n,$$

where π_i denotes that car π_i parked in spot i on the street.

Example

From our previous example: If $\alpha = (3, 1, 1)$ then $\pi = 231$.



More Examples

Example

- If $\alpha = (1, 1, 1, 1)$ then $\pi = 1234$

More Examples

Example

- If $\alpha = (1, 1, 1, 1)$ then $\pi = 1234$
- If $\alpha = (4, 1, 3, 2)$ then $\pi = 2431$

More Examples

Example

- If $\alpha = (1, 1, 1, 1)$ then $\pi = 1234$
- If $\alpha = (4, 1, 3, 2)$ then $\pi = 2431$
 - Observation: If all entries are distinct in α then $\pi = \alpha^{-1}$

Lucky Cars

Definition

A car is said to be *lucky* if it parks on its preferred parking spot.

Lucky Cars

Definition

A car is said to be *lucky* if it parks on its preferred parking spot.

Example

- If $\alpha = (1, 1, 1, 1)$ then $\pi = 1234$

Lucky Cars

Definition

A car is said to be *lucky* if it parks on its preferred parking spot.

Example

- If $\alpha = (1, 1, 1, 1)$ then $\pi = \textcolor{blue}{1234}$
- If $\alpha = (4, 1, 3, 2)$ then $\pi = \textcolor{blue}{2431}$

Lucky Cars

Definition

A car is said to be *lucky* if it parks on its preferred parking spot.

Example

- If $\alpha = (1, 1, 1, 1)$ then $\pi = \textcolor{blue}{1234}$
- If $\alpha = (4, 1, 3, 2)$ then $\pi = \textcolor{blue}{2431}$
 - Observation: If the entries in a parking function are distinct, then all cars are lucky.

Previous Work

Theorem (Gessel and Seo)

The generating function for the number of parking functions based on the number of lucky cars is given by

$$L_n(q) = \sum_{\alpha \in PF_n} q^{\text{lucky}(\alpha)} = q \prod_{i=1}^{n-1} (i + (n - i + 1)q).$$

Previous Work

Theorem (Gessel and Seo)

The generating function for the number of parking functions based on the number of lucky cars is given by

$$L_n(q) = \sum_{\alpha \in PFn} q^{\text{lucky}(\alpha)} = q \prod_{i=1}^{n-1} (i + (n - i + 1)q).$$

Example

Let $n = 3$ then the generating function is $2q + 8q^2 + 6q^3$. Recall:

Previous Work

Theorem (Gessel and Seo)

The generating function for the number of parking functions based on the number of lucky cars is given by

$$L_n(q) = \sum_{\alpha \in PFn} q^{\text{lucky}(\alpha)} = q \prod_{i=1}^{n-1} (i + (n - i + 1)q).$$

Example

Let $n = 3$ then the generating function is $2q + 8q^2 + 6q^3$. Recall:

| | | | | |
|-----|--|--------------------|--|---------------|
| 111 | | 211, 121, 122, 221 | | 231, 123, 132 |
| 112 | | 212, 113, 131, 311 | | 213, 312, 321 |

Previous Work

Theorem (Gessel and Seo)

The generating function for the number of parking functions based on the number of lucky cars is given by

$$L_n(q) = \sum_{\alpha \in PFn} q^{\text{lucky}(\alpha)} = q \prod_{i=1}^{n-1} (i + (n - i + 1)q).$$

Note: This does not tell you which cars are lucky!

Notation

Definition

Let $\text{Lucky}(\alpha)$ denote the set of lucky cars of $\alpha \in PF_n$.
We will often denote $I = \text{Lucky}(\alpha)$.

Notation

Definition

Let $\text{Lucky}(\alpha)$ denote the set of lucky cars of $\alpha \in PF_n$.
We will often denote $I = \text{Lucky}(\alpha)$.

Example

- $\text{Lucky}((1, 1, 1, 1)) = \{1\}$

Notation

Definition

Let $\text{Lucky}(\alpha)$ denote the set of lucky cars of $\alpha \in PF_n$.
We will often denote $I = \text{Lucky}(\alpha)$.

Example

- $\text{Lucky}((1, 1, 1, 1)) = \{1\}$
- $\text{Lucky}((1, 1, 3, 2)) = \{1, 3\}$

Notation

Definition

Let $\text{Lucky}(\alpha)$ denote the set of lucky cars of $\alpha \in PF_n$.
We will often denote $I = \text{Lucky}(\alpha)$.

Example

- $\text{Lucky}((1, 1, 1, 1)) = \{1\}$
- $\text{Lucky}((1, 1, 3, 2)) = \{1, 3\}$
- $\text{Lucky}(\alpha) = \{1, 2, \dots, n\}$ whenever $\alpha = (\pi_1, \pi_2, \dots, \pi_n)$ and $\pi = \pi_1 \pi_2 \cdots \pi_n \in \mathfrak{S}_n$.

Motivation

Motivation

Definition

For any subset $I \subseteq [n]$, we define

$$\text{LuckyPF}_n(I) := \{\alpha \in PF_n : \text{Lucky}(\alpha) = I\},$$

which is the set of parking functions of length n with lucky cars being the cars in the set I and unlucky cars being the cars in the set $[n] \setminus I$.

Motivation

Definition

For any subset $I \subseteq [n]$, we define

$$\text{LuckyPF}_n(I) := \{\alpha \in PF_n : \text{Lucky}(\alpha) = I\},$$

which is the set of parking functions of length n with lucky cars being the cars in the set I and unlucky cars being the cars in the set $[n] \setminus I$.

Remarks

- For $I \subseteq [n]$, if $\text{LuckyPF}_n(I) \neq \emptyset$, then I is said to be a **lucky set** of PF_n .

Motivation

Definition

For any subset $I \subseteq [n]$, we define

$$\text{LuckyPF}_n(I) := \{\alpha \in PF_n : \text{Lucky}(\alpha) = I\},$$

which is the set of parking functions of length n with lucky cars being the cars in the set I and unlucky cars being the cars in the set $[n] \setminus I$.

Remarks

- For $I \subseteq [n]$, if $\text{LuckyPF}_n(I) \neq \emptyset$, then I is said to be a **lucky set** of PF_n .
- Since the first car is always a lucky car regardless of the preference of the other cars, a set I not containing 1 can never be a lucky set of PF_n for any $n \geq 1$.

Motivation

Definition

For any subset $I \subseteq [n]$, we define

$$\text{LuckyPF}_n(I) := \{\alpha \in PF_n : \text{Lucky}(\alpha) = I\},$$

which is the set of parking functions of length n with lucky cars being the cars in the set I and unlucky cars being the cars in the set $[n] \setminus I$.

Remarks

- For $I \subseteq [n]$, if $\text{LuckyPF}_n(I) \neq \emptyset$, then I is said to be a **lucky set** of PF_n .
- Since the first car is always a lucky car regardless of the preference of the other cars, a set I not containing 1 can never be a lucky set of PF_n for any $n \geq 1$.
- We stress that lucky sets contain the lucky cars rather than the position at which the lucky cars park.

Motivation (2)

- Not every permutation can be an outcome of a parking function with a fixed set of lucky cars!

Motivation (2)

- Not every permutation can be an outcome of a parking function with a fixed set of lucky cars!

We begin with an example:

Motivation (2)

- Not every permutation can be an outcome of a parking function with a fixed set of lucky cars!

We begin with an example:

- Let $\pi = 312 \in \mathfrak{S}_3$, then π cannot be the outcome of a parking function with lucky set $\{1\}$.

Motivation (2)

- Not every permutation can be an outcome of a parking function with a fixed set of lucky cars!

We begin with an example:

- Let $\pi = 312 \in \mathfrak{S}_3$, then π cannot be the outcome of a parking function with lucky set $\{1\}$.
- Why?

Motivation (2)

- Not every permutation can be an outcome of a parking function with a fixed set of lucky cars!

We begin with an example:

- Let $\pi = 312 \in \mathfrak{S}_3$, then π cannot be the outcome of a parking function with lucky set $\{1\}$.
- Why? Car $3 = \pi_1$ parks in spot 1. Moreover, it was the first car to prefer spot 1 since it was empty when it entered the street.

Motivation (2)

- Not every permutation can be an outcome of a parking function with a fixed set of lucky cars!

We begin with an example:

- Let $\pi = 312 \in \mathfrak{S}_3$, then π cannot be the outcome of a parking function with lucky set $\{1\}$.
- Why?
- This means that parking in spot 1 makes car 3 a lucky car, but $I = \{1\}$ so we reach a contradiction.

Motivation (2)

- Not every permutation can be an outcome of a parking function with a fixed set of lucky cars!

We begin with an example:

- Let $\pi = 312 \in \mathfrak{S}_3$, then π cannot be the outcome of a parking function with lucky set $\{1\}$.
- Why?
- This means that parking in spot 1 makes car 3 a lucky car, but $I = \{1\}$ so we reach a contradiction.

Question

For a fixed subset $I \subseteq [n]$ (containing 1), what permutations in \mathfrak{S}_n arise as the outcome of parking functions with lucky set I ?

Results

Lucky Sets

Theorem (Harris, M.)

A subset $I \subseteq [n]$ is a lucky set of PF_n if and only if $1 \in I$.

Lucky Sets

Theorem (Harris, M.)

A subset $I \subseteq [n]$ is a lucky set of PF_n if and only if $1 \in I$.

Proof.

(\Rightarrow) If I is a lucky set of PF_n , then there exists a parking function $\alpha \in PF_n$ such that $\text{Lucky}(\alpha) = I$.



Lucky Sets

Theorem (Harris, M.)

A subset $I \subseteq [n]$ is a lucky set of PF_n if and only if $1 \in I$.

Proof.

(\Rightarrow) If I is a lucky set of PF_n , then there exists a parking function $\alpha \in PF_n$ such that $\text{Lucky}(\alpha) = I$. First car is always a lucky car for any $\alpha \in PF_n$, this implies that $1 \in I$.



Lucky Sets

Theorem (Harris, M.)

A subset $I \subseteq [n]$ is a lucky set of PF_n if and only if $1 \in I$.

Proof.

(\Rightarrow) If I is a lucky set of PF_n , then there exists a parking function $\alpha \in PF_n$ such that $\text{Lucky}(\alpha) = I$. First car is always a lucky car for any $\alpha \in PF_n$, this implies that $1 \in I$.

(\Leftarrow) Assume $I = \{i_1 = 1, i_2, i_3, \dots, i_k\}$, with $k \in [n]$.



Lucky Sets

Theorem (Harris, M.)

A subset $I \subseteq [n]$ is a lucky set of PF_n if and only if $1 \in I$.

Proof.

(\Rightarrow) If I is a lucky set of PF_n , then there exists a parking function $\alpha \in PF_n$ such that $\text{Lucky}(\alpha) = I$. First car is always a lucky car for any $\alpha \in PF_n$, this implies that $1 \in I$.

(\Leftarrow) Assume $I = \{i_1 = 1, i_2, i_3, \dots, i_k\}$, with $k \in [n]$.

Without loss of generality assume that $i_1 = 1 < i_2 < i_3 < \dots < i_k$.



Lucky Sets

Theorem (Harris, M.)

A subset $I \subseteq [n]$ is a lucky set of PF_n if and only if $1 \in I$.

Proof.

(\Rightarrow) If I is a lucky set of PF_n , then there exists a parking function $\alpha \in PF_n$ such that $\text{Lucky}(\alpha) = I$. First car is always a lucky car for any $\alpha \in PF_n$, this implies that $1 \in I$.

(\Leftarrow) Assume $I = \{i_1 = 1, i_2, i_3, \dots, i_k\}$, with $k \in [n]$.

Without loss of generality assume that $i_1 = 1 < i_2 < i_3 < \dots < i_k$.

We want to construct α such that $\alpha \in PF_n$ and $\text{Lucky}(\alpha) = I$. □

Lucky Sets

Theorem (Harris, M.)

A subset $I \subseteq [n]$ is a lucky set of PF_n if and only if $1 \in I$.

Proof.

(\Rightarrow) If I is a lucky set of PF_n , then there exists a parking function $\alpha \in PF_n$ such that $\text{Lucky}(\alpha) = I$.

(\Leftarrow) Assume $I = \{i_1 = 1, i_2, i_3, \dots, i_k\}$, with $k \in [n]$.

Without loss of generality assume that $i_1 = 1 < i_2 < i_3 < \dots < i_k$.

We want to construct α such that $\alpha \in PF_n$ and $\text{Lucky}(\alpha) = I$. Let $\alpha = (a_1, a_2, \dots, a_n)$ where

$$a_x = \begin{cases} x & \text{if } x \in I \\ i_\ell & \text{if } i_\ell < x < i_{\ell+1} . \end{cases}$$



Permutations

We begin with an example that gives insight into the types of outcome permutations that arise from a fixed lucky set I .

Permutations

We begin with an example that gives insight into the types of outcome permutations that arise from a fixed lucky set I .

Example

Let $\alpha \in PF_4$ and suppose that $\text{Lucky}(\alpha) = \{1, 4\}$.

Permutations

We begin with an example that gives insight into the types of outcome permutations that arise from a fixed lucky set I .

Example

Let $\alpha \in PF_4$ and suppose that $\text{Lucky}(\alpha) = \{1, 4\}$.

Notice that $\pi = 1423 \in \mathfrak{S}_4$ is not a possible outcome. Why?

Permutations

We begin with an example that gives insight into the types of outcome permutations that arise from a fixed lucky set I .

Example

Let $\alpha \in PF_4$ and suppose that $\text{Lucky}(\alpha) = \{1, 4\}$.

Notice that $\pi = 1423 \in S_4$ is not a possible outcome. Why?

Car 2 must be in the lucky set.

Permutations

We begin with an example that gives insight into the types of outcome permutations that arise from a fixed lucky set I .

Example

Let $\alpha \in PF_4$ and suppose that $\text{Lucky}(\alpha) = \{1, 4\}$.

Notice that $\pi = 1423 \in \mathfrak{S}_4$ is not a possible outcome. Why?

Car 2 must be in the lucky set.

Note: the descents in a permutation force certain cars to be lucky!

Descents

Definition

Given a permutation $\pi = \pi_1\pi_2 \cdots \pi_n \in \mathfrak{S}_n$, an index $1 < i \leq n$ is a *descent* of π if $\pi_{i-1} > \pi_i$, and the value π_i is called a *descent bottom* of π .

Descents

Definition

Given a permutation $\pi = \pi_1 \pi_2 \cdots \pi_n \in \mathfrak{S}_n$, an index $1 < i \leq n$ is a *descent* of π if $\pi_{i-1} > \pi_i$, and the value π_i is called a *descent bottom* of π . For convenience, we say $i = 1$ is always a descent and π_1 is always a descent bottom.

Descents

Definition

Given a permutation $\pi = \pi_1\pi_2 \cdots \pi_n \in \mathfrak{S}_n$, an index $1 < i \leq n$ is a *descent* of π if $\pi_{i-1} > \pi_i$, and the value π_i is called a *descent bottom* of π . For convenience, we say $i = 1$ is always a descent and π_1 is always a descent bottom.

Example

- ① If $\pi = 1423$ then the descent bottoms are at $i = 1, 3$.

Descents

Definition

Given a permutation $\pi = \pi_1\pi_2 \cdots \pi_n \in \mathfrak{S}_n$, an index $1 < i \leq n$ is a *descent* of π if $\pi_{i-1} > \pi_i$, and the value π_i is called a *descent bottom* of π . For convenience, we say $i = 1$ is always a descent and π_1 is always a descent bottom.

Example

- ① If $\pi = 1423$ then the descent bottoms are at $i = 1, 3$.
- ② If $\pi = 45312$ then the descent bottoms are at $i = 1, 3, 4$.

Descent Bottoms and Lucky Sets

Lemma (Harris, M.)

Fix $\pi = \pi_1\pi_2 \cdots \pi_n \in \mathfrak{S}_n$. For any $\alpha \in PF_n$ with outcome $\mathcal{O}(\alpha) = \pi$, if $i \in \text{Des}(\pi)$, then $\pi_i \in \text{Lucky}(\alpha)$. In other words,

$$\text{DBottom}(\pi) \subseteq \text{Lucky}(\alpha).$$

Descent Bottoms and Lucky Sets

Lemma (Harris, M.)

Fix $\pi = \pi_1\pi_2 \cdots \pi_n \in \mathfrak{S}_n$. For any $\alpha \in PF_n$ with outcome $\mathcal{O}(\alpha) = \pi$, if $i \in \text{Des}(\pi)$, then $\pi_i \in \text{Lucky}(\alpha)$. In other words,

$$\text{DBottom}(\pi) \subseteq \text{Lucky}(\alpha).$$

If $\pi = 58164237$ then cars 1, 2, 4 and 5 must be in the lucky set.

Descent Bottoms and Lucky Sets

Lemma (Harris, M.)

Fix $\pi = \pi_1\pi_2 \cdots \pi_n \in \mathfrak{S}_n$. For any $\alpha \in PF_n$ with outcome $\mathcal{O}(\alpha) = \pi$, if $i \in \text{Des}(\pi)$, then $\pi_i \in \text{Lucky}(\alpha)$. In other words,

$$\text{DBottom}(\pi) \subseteq \text{Lucky}(\alpha).$$

If $\pi = 58164237$ then cars 1, 2, 4 and 5 must be in the lucky set.

Remark: Descent bottoms are lucky cars but not all lucky cars have to be descent bottoms.

Descent Bottoms and Lucky Sets

Lemma (Harris, M.)

Fix $\pi = \pi_1\pi_2\cdots\pi_n \in \mathfrak{S}_n$. For any $\alpha \in PF_n$ with outcome $\mathcal{O}(\alpha) = \pi$, if $i \in \text{Des}(\pi)$, then $\pi_i \in \text{Lucky}(\alpha)$. In other words,

$$\text{DBottom}(\pi) \subseteq \text{Lucky}(\alpha).$$

If $\pi = 58164237$ then cars 1, 2, 4 and 5 must be in the lucky set.

Remark: Descent bottoms are lucky cars but not all lucky cars have to be descent bottoms.

For example, if $I = \{1, 2, 3, 4\}$ and $\pi = 1234$. The only descent bottom is at $i = 1$.

Characterization

Theorem (Harris, M.)

Fix a lucky set $I \subseteq [n]$ of PF_n . Then $\pi = \pi_1 \pi_2 \cdots \pi_n \in \mathfrak{S}_n$ is the outcome an $\alpha \in \text{LuckyPF}_n(I)$ if and only if

- ① $\pi_1 \in I$,
- ② if $\pi_i \in I$ and $\pi_{i+1} \notin I$, then $\pi_i < \pi_{i+1}$, and
- ③ if $\pi_{i-1} > \pi_i$, then $\pi_i \in I$.

Note: Condition 2 follows from condition 3 via the converse.

Characterization Example

Example

Suppose we want to construct all outcomes of length $n = 5$ such that $I = \{1, 4\}$.

Characterization Example

Example

Suppose we want to construct all outcomes of length $n = 5$ such that $I = \{1, 4\}$.

By the characterization theorem, $\pi = \pi_1\pi_2\pi_3\pi_4\pi_5 \in \mathcal{O}_5(\{1, 4\})$ if and only if

- ① $\pi_1 = 1$ or $\pi_1 = 4$
- ② if $\pi_i \in I$ and $\pi_{i+1} \notin I$, then $\pi_i < \pi_{i+1}$, and
- ③ if $\pi_{i-1} > \pi_i$, then $\pi_i \in I$.

Characterization Example

Example

Suppose we want to construct all outcomes of length $n = 5$ such that $I = \{1, 4\}$.

By the characterization theorem, $\pi = \pi_1\pi_2\pi_3\pi_4\pi_5 \in \mathcal{O}_5(\{1, 4\})$ if and only if

- ① $\pi_1 = 1$ or $\pi_1 = 4$
- ② if $\pi_i \in I$ and $\pi_{i+1} \notin I$, then $\pi_i < \pi_{i+1}$, and
- ③ if $\pi_{i-1} > \pi_i$, then $\pi_i \in I$.

We know $\text{DBottom}(\pi) \subseteq I$. There are $2 \cdot 4! = 48$ permutations of length $n = 5$ satisfying Condition (1), as $\pi = 1\sigma$ with $\sigma \in \mathfrak{S}_{\{2,3,4,5\}}$ or $\pi = 4\sigma$ with $\sigma \in \mathfrak{S}_{\{1,2,3,5\}}$.

Characterization Example

Example

Suppose we want to construct all outcomes of length $n = 5$ such that $I = \{1, 4\}$.

We know $\text{DBottom}(\pi) \subseteq I$. There are $2 \cdot 4! = 48$ permutations of length $n = 5$ satisfying Condition (1), as $\pi = 1\sigma$ with $\sigma \in \mathfrak{S}_{\{2,3,4,5\}}$ or $\pi = 4\sigma$ with $\sigma \in \mathfrak{S}_{\{1,2,3,5\}}$.

- If $\pi = 1\sigma$ with $\sigma = \sigma_1\sigma_2\sigma_3\sigma_4 \in \mathfrak{S}_{\{2,3,4,5\}}$, then $2 \leq \sigma_1 \leq 5$. Then, σ can either have no descents, or a descent bottom $\sigma_i = 4$ for some i . Hence π can only be 12345 or 12354.

Characterization Example

Example

Suppose we want to construct all outcomes of length $n = 5$ such that $I = \{1, 4\}$.

We know $\text{DBottom}(\pi) \subseteq I$. There are $2 \cdot 4! = 48$ permutations of length $n = 5$ satisfying Condition (1), as $\pi = 1\sigma$ with $\sigma \in \mathfrak{S}_{\{2,3,4,5\}}$ or $\pi = 4\sigma$ with $\sigma \in \mathfrak{S}_{\{1,2,3,5\}}$.

- If $\pi = 1\sigma$ with $\sigma = \sigma_1\sigma_2\sigma_3\sigma_4 \in \mathfrak{S}_{\{2,3,4,5\}}$, then $2 \leq \sigma_1 \leq 5$. Then, σ can either have no descents, or a descent bottom $\sigma_i = 4$ for some i . Hence π can only be 12345 or 12354.
- If $\pi = 4\sigma$ with $\sigma = \sigma_1\sigma_2\sigma_3\sigma_4 \in \mathfrak{S}_{\{1,2,4,5\}}$, then σ must have only a descent bottom $\sigma_i = 1$ for some i . Hence π can only be 41235 or 45123.

Characterization Example

Example

Suppose we want to construct all outcomes of length $n = 5$ such that $I = \{1, 4\}$.

We know $\text{DBottom}(\pi) \subseteq I$. There are $2 \cdot 4! = 48$ permutations of length $n = 5$ satisfying Condition (1), as $\pi = 1\sigma$ with $\sigma \in \mathfrak{S}_{\{2,3,4,5\}}$ or $\pi = 4\sigma$ with $\sigma \in \mathfrak{S}_{\{1,2,3,5\}}$.

- If $\pi = 1\sigma$ with $\sigma = \sigma_1\sigma_2\sigma_3\sigma_4 \in \mathfrak{S}_{\{2,3,4,5\}}$, then $2 \leq \sigma_1 \leq 5$. Then, σ can either have no descents, or a descent bottom $\sigma_i = 4$ for some i . Hence π can only be 12345 or 12354.
- If $\pi = 4\sigma$ with $\sigma = \sigma_1\sigma_2\sigma_3\sigma_4 \in \mathfrak{S}_{\{1,2,4,5\}}$, then σ must have only a descent bottom $\sigma_i = 1$ for some i . Hence π can only be 41235 or 45123.

Thus $\mathcal{O}_5(\{1, 4\}) = \{12345, 12354, 41235, 45123\}$.

Enumeration of Outcomes

Given a lucky set $I \subseteq [n]$, can we count the number of outcomes with that lucky set?

Enumeration of Outcomes

Given a lucky set $I \subseteq [n]$, can we count the number of outcomes with that lucky set?

Theorem (Harris, M.)

Let $I = \{1, 2, 3, \dots, k\} \subseteq [n]$ be a lucky set of PF_n . Then

$$|\mathcal{O}_n(I)| = \sum_{J \subseteq [n]} k! \binom{n - k}{j_2 - j_1 - 1, j_3 - j_2 - 1, \dots, j_k - j_{k-1} - 1, n - j_k}$$

where $J = \{j_1 = 1, j_2, j_3, \dots, j_k\}$.

Enumeration of Outcomes

Given a lucky set $I \subseteq [n]$, can we count the number of outcomes with that lucky set?

Theorem (Harris, M.)

Let $I = \{1, 2, 3, \dots, k\} \subseteq [n]$ be a lucky set of PF_n . Then

$$|\mathcal{O}_n(I)| = \sum_{J \subseteq [n]} k! \binom{n-k}{j_2 - j_1 - 1, j_3 - j_2 - 1, \dots, j_k - j_{k-1} - 1, n - j_k}$$

where $J = \{j_1 = 1, j_2, j_3, \dots, j_k\}$.

Note: We have since found and proven a closed formula for this case and the general case!

Enumeration of Parking Functions

What about given a lucky set $I \subseteq [n]$, can we count the number of parking functions with that lucky set?

Enumeration of Parking Functions

What about given a lucky set $I \subseteq [n]$, can we count the number of parking functions with that lucky set?

We should focus on weakly increasing parking function, i.e.

$\alpha = (a_1, a_2 \dots, a_n)$ where $a_1 \leq \dots \leq a_n$.

Enumeration of Parking Functions

What about given a lucky set $I \subseteq [n]$, can we count the number of parking functions with that lucky set?

We should focus on weakly increasing parking function, i.e.

$\alpha = (a_1, a_2 \dots, a_n)$ where $a_1 \leq \dots \leq a_n$.

Theorem (Harris, M.)

Fix a lucky set $I = \{i_1, i_2, i_3, \dots, i_k\} \subseteq [n]$ of PF_n^\uparrow and assume $1 = i_1 < i_2 < \dots < i_k \leq n$. Then

$$|\text{LuckyPF}_n^\uparrow(I)| = \prod_{j=1}^k \text{Cat}_{x_j}, \quad (1)$$

where $x_j = i_{j+1} - i_j - 1$ for each $j \in [k-1]$ and $x_k = n - i_k$.

Enumeration of Parking Functions (2)

Example

Let $n = 9$ and $I = \{1, 6, 8\}$. We want to count all the weakly increasing parking functions of length 9 whose lucky cars are 1, 6 and 8.

Enumeration of Parking Functions (2)

Example

Let $n = 9$ and $I = \{1, 6, 8\}$. We want to count all the weakly increasing parking functions of length 9 whose lucky cars are 1, 6 and 8.

We have the following picture:

Enumeration of Parking Functions (2)

Example

Let $n = 9$ and $I = \{1, 6, 8\}$. We want to count all the weakly increasing parking functions of length 9 whose lucky cars are 1, 6 and 8.

We have the following picture:

$$\frac{1}{1} \quad \frac{2}{2} \quad \frac{3}{3} \quad \frac{4}{4} \quad \frac{5}{5} \quad \frac{6}{6} \quad \frac{7}{7} \quad \frac{8}{8} \quad \frac{9}{9}$$

Enumeration of Parking Functions (2)

Example

Let $n = 9$ and $I = \{1, 6, 8\}$. We want to count all the weakly increasing parking functions of length 9 whose lucky cars are 1, 6 and 8.

We have the following picture:

$$\begin{array}{ccccccccc} \frac{1}{1} & \frac{2}{2} & \frac{3}{3} & \frac{4}{4} & \frac{5}{5} & \frac{6}{6} & \frac{7}{7} & \frac{8}{8} & \frac{9}{9} \\ & \underbrace{\phantom{\frac{3}{3}}}_{1 \leq a_2 \leq 1} & \underbrace{\phantom{\frac{3}{3}}}_{1 \leq a_3 \leq 2} & \underbrace{\phantom{\frac{3}{3}}}_{1 \leq a_4 \leq 3} & \underbrace{\phantom{\frac{3}{3}}}_{1 \leq a_5 \leq 4} & & \underbrace{\phantom{\frac{7}{7}}}_{6 \leq a_7 \leq 6} & & \underbrace{\phantom{\frac{9}{9}}}_{8 \leq a_9 \leq 8} \end{array}$$

All tuples of the form $(1, a_2, a_3, a_4, a_5, 6, a_7, 8, a_9)$

Enumeration of Parking Functions (2)

Example

Let $n = 9$ and $I = \{1, 6, 8\}$. We want to count all the weakly increasing parking functions of length 9 whose lucky cars are 1, 6 and 8.

We have the following picture:

$$\begin{array}{ccccccccc} \frac{1}{1} & \frac{2}{2} & \frac{3}{3} & \frac{4}{4} & \frac{5}{5} & \frac{6}{6} & \frac{7}{7} & \frac{8}{8} & \frac{9}{9} \\[10pt] 1 \leq a_2 \leq 1 & & & & & 6 \leq a_7 \leq 6 & & 8 \leq a_9 \leq 8 \\ 1 \leq a_3 \leq 2 & & & & & & & \\ 1 \leq a_4 \leq 3 & & & & & & & \\ 1 \leq a_5 \leq 4 & & & & & & & \\ \underbrace{\hspace{10em}}_{\text{Cat}_4 = \text{Cat}_{6-1-1}} & & & & & \underbrace{\hspace{1.5em}}_{\text{Cat}_1} & & \underbrace{\hspace{1.5em}}_{\text{Cat}_1 = \text{Cat}_{9-8}} \end{array}$$

All tuples of the form $(1, a_2, a_3, a_4, a_5, 6, a_7, 8, a_9)$

Enumeration of Parking Functions (2)

Example

Let $n = 9$ and $I = \{1, 6, 8\}$. We want to count all the weakly increasing parking functions of length 9 whose lucky cars are 1, 6 and 8.

We have the following picture:

$$\begin{array}{ccccccccc} \frac{1}{1} & \frac{2}{2} & \frac{3}{3} & \frac{4}{4} & \frac{5}{5} & \frac{6}{6} & \frac{7}{7} & \frac{8}{8} & \frac{9}{9} \\ & & & & & & \underbrace{}_{6 \leq a_7 \leq 6} & & \underbrace{}_{8 \leq a_9 \leq 8} \\ 1 \leq a_2 \leq 1 & 1 \leq a_3 \leq 2 & 1 \leq a_4 \leq 3 & 1 \leq a_5 \leq 4 & & & & & \\ & \underbrace{}_{\text{Cat}_4 = \text{Cat}_{6-1-1}} & & & & \underbrace{}_{\text{Cat}_1} & & \underbrace{}_{\text{Cat}_1 = \text{Cat}_{9-8}} & \end{array}$$

All tuples of the form $(1, a_2, a_3, a_4, a_5, 6, a_7, 8, a_9)$

This implies that the number of weakly increasing parking functions with lucky set $I = \{1, 6, 8\}$ is $\text{Cat}_4 \cdot \text{Cat}_1 \cdot \text{Cat}_1 = 14 \cdot 1 \cdot 1 = 14$.

Summary

- We characterized the types of permutations that are outcomes of parking functions with a fixed set of lucky cars I .

Summary

- We characterized the types of permutations that are outcomes of parking functions with a fixed set of lucky cars I .
- We provided a sum formula for the number of *outcomes* where the first k cars are lucky, i.e. $I = \{1, 2, \dots, k\}$ (which is closed now).

Summary

- We characterized the types of permutations that are outcomes of parking functions with a fixed set of lucky cars I .
- We provided a sum formula for the number of *outcomes* where the first k cars are lucky, i.e. $I = \{1, 2, \dots, k\}$ (which is closed now).
- We provided a formula for the number of *weakly increasing parking functions* with a fixed set of lucky cars I .

Summary

- We characterized the types of permutations that are outcomes of parking functions with a fixed set of lucky cars I .
- We provided a sum formula for the number of *outcomes* where the first k cars are lucky, i.e. $I = \{1, 2, \dots, k\}$ (which is closed now).
- We provided a formula for the number of *weakly increasing parking functions* with a fixed set of lucky cars I .
- More results we did not discuss: formula for the number of *parking functions* with a fixed set of lucky cars I .

Summary

- We characterized the types of permutations that are outcomes of parking functions with a fixed set of lucky cars I .
- We provided a sum formula for the number of *outcomes* where the first k cars are lucky, i.e. $I = \{1, 2, \dots, k\}$ (which is closed now).
- We provided a formula for the number of *weakly increasing parking functions* with a fixed set of lucky cars I .
- More results we did not discuss: formula for the number of *parking functions* with a fixed set of lucky cars I .

What happens to all the previous results when there are more spots than cars?

Thank You!



<https://marti310.github.io/research.html>