

How many coin tosses would you need until you get
 n Heads or m Tails?

Lucy Martinez

Rutgers University

Experimental Math Seminar
–Joint work with Svante Janson
and Doron Zeilberger–

Background and History

Problem of points

In 1654, Fermat and Pascal were interested in the following game.

Problem of points

In 1654, Fermat and Pascal were interested in the following game.

Two players compete in a fair game such as “first to win ten rounds takes the prize”. Suppose that the game stops early, for example at a score of 3 to 4. How should the players decide how to split the prize?

Problem of points

In 1654, Fermat and Pascal were interested in the following game.

Two players compete in a fair game such as “first to win ten rounds takes the prize”. Suppose that the game stops early, for example at a score of 3 to 4. How should the players decide how to split the prize?

The correspondence (snail mail) between Fermat and Pascal laid the foundation to what is now known as probability theory.

Problem of points (2)

We can rephrase the problem of points as follows. Suppose we toss a fair coin.

- Instead of rounds, we can think of the game as tossing a coin until we get to n Heads OR m Tails.

Problem of points (2)

We can rephrase the problem of points as follows. Suppose we toss a fair coin.

- Instead of rounds, we can think of the game as tossing a coin until we get to n Heads OR m Tails.
- Why?

Problem of points (2)

We can rephrase the problem of points as follows. Suppose we toss a fair coin.

- Instead of rounds, we can think of the game as tossing a coin until we get to n Heads OR m Tails.
- Why?
- If player 1 has n rounds left to win and player 2 has m rounds to win, then we can translate that into tossing a fair coin until we get the corresponding face values.

Problem of points (2)

We can rephrase the problem of points as follows. Suppose we toss a fair coin.

- Instead of rounds, we can think of the game as tossing a coin until we get to n Heads OR m Tails.
- Why?
- If player 1 has n rounds left to win and player 2 has m rounds to win, then we can translate that into tossing a fair coin until we get the corresponding face values.

Convention: For the rest of the talk, p represents the probability of getting Heads and $q := 1 - p$ represents the probability of getting Tails.

Probability generating function until reaching n Heads

Consider calculating the probability that we reach n Heads. Let $q = 1 - p$, then

$$\sum_{t=0}^{\infty} \binom{n+t-1}{n-1} (px)^n (qx)^t$$

Probability generating function until reaching n Heads

Consider calculating the probability that we reach n Heads. Let $q = 1 - p$, then

$$\sum_{t=0}^{\infty} \binom{n+t-1}{n-1} (px)^n (qx)^t = \binom{n-1}{n-1} (px)^n (qx)^0 + \binom{n}{n-1} (px)^n (qx)^1 + \dots$$

Probability generating function until reaching n Heads

Consider calculating the probability that we reach n Heads. Let $q = 1 - p$, then

$$\sum_{t=0}^{\infty} \binom{n+t-1}{n-1} (px)^n (qx)^t = \binom{n-1}{n-1} (px)^n (qx)^0 + \binom{n}{n-1} (px)^n (qx)^1 + \dots$$

$n + t - 1$ is the maximum number of coin tosses and the binomial coefficient represents the number of ways to get $n - 1$ Heads out of $n + t - 1$ coin tosses.

Probability generating function until reaching n Heads

Consider calculating the probability that we reach n Heads. Let $q = 1 - p$, then

$$\begin{aligned}\sum_{t=0}^{\infty} \binom{n+t-1}{n-1} (px)^n (qx)^t &= \binom{n-1}{n-1} (px)^n (qx)^0 + \binom{n}{n-1} (px)^n (qx)^1 + \cdots \\ &= \left(\frac{px}{1-qx} \right)^n.\end{aligned}$$

$n + t - 1$ is the maximum number of coin tosses and the binomial coefficient represents the number of ways to get $n - 1$ Heads out of $n + t - 1$ coin tosses.

Probability generating function until reaching n Heads

Consider calculating the probability that we reach n Heads. Let $q = 1 - p$, then

$$\begin{aligned}\sum_{t=0}^{\infty} \binom{n+t-1}{n-1} (px)^n (qx)^t &= \binom{n-1}{n-1} (px)^n (qx)^0 + \binom{n}{n-1} (px)^n (qx)^1 + \dots \\ &= \left(\frac{px}{1-qx} \right)^n.\end{aligned}$$

$n + t - 1$ is the maximum number of coin tosses and the binomial coefficient represents the number of ways to get $n - 1$ Heads out of $n + t - 1$ coin tosses.

Remark: Fermat and Pascal were only interested in the probability of getting n Heads vs. getting m Tails.

Computations

Probability generating function (n Heads OR m Tails)

$X := X(n, m; p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Probability generating function (n Heads OR m Tails)

$X := X(n, m; p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Let $F(n, m; p)$ be the probability generating function of X in terms of x , then

Probability generating function (n Heads OR m Tails)

$X := X(n, m; p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Let $F(n, m; p)$ be the probability generating function of X in terms of x , then

$$\begin{aligned} F(n, m; p)(x) &= (px)^n \sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (qx)^t \\ &\quad + (qx)^m \sum_{h=0}^{n-1} \binom{m+h-1}{m-1} (px)^h. \end{aligned}$$

Probability generating function (n Heads OR m Tails)

$X := X(n, m; p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Let $F(n, m; p)$ be the probability generating function of X in terms of x , then

$$F(n, m; p)(x) = (px)^n \sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (qx)^t \\ + (qx)^m \sum_{h=0}^{n-1} \binom{m+h-1}{m-1} (px)^h.$$

In the first sum: Each term $p^n \binom{n+t-1}{n-1} q^t$ represents the probability of getting n Heads and t Tails.

Probability generating function (n Heads OR m Tails)

$X := X(n, m; p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Let $F(n, m; p)$ be the probability generating function of X in terms of x , then

$$\begin{aligned} F(n, m; p)(x) &= (px)^n \sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (qx)^t \\ &\quad + (qx)^m \sum_{h=0}^{n-1} \binom{m+h-1}{m-1} (px)^h. \end{aligned}$$

Remark: $F(n, m; p)$ does not have a closed-form but *does* have a linear recurrence with polynomial coefficients. We do not show it here for sake of space.

Computing expected coin tosses (n Heads OR m Tails)

To compute *some* data, we can take the derivative of the previous sum $F(n, m; p)(x)$ with respect to x and then plug in $x = 1$,

Computing expected coin tosses (n Heads OR m Tails)

To compute *some* data, we can take the derivative of the previous sum $F(n, m; p)(x)$ with respect to x and then plug in $x = 1$,

$$p^n \sum_{t=0}^{m-1} (n+t) \binom{n-1+t}{n-1} q^t + q^m \sum_{h=0}^{n-1} (m+h) \binom{m-1+h}{m-1} p^h, \quad q = 1-p.$$

Computing expected coin tosses (n Heads OR m Tails)

To compute *some* data, we can take the derivative of the previous sum $F(n, m; p)(x)$ with respect to x and then plug in $x = 1$,

$$p^n \sum_{t=0}^{m-1} (n+t) \binom{n-1+t}{n-1} q^t + q^m \sum_{h=0}^{n-1} (m+h) \binom{m-1+h}{m-1} p^h, \quad q = 1-p.$$

Remark: This becomes computationally slow. Using Wilf-Zeilberger algorithmic proof theory, we will obtain a third-order linear recurrence!

Results (1)

Recurrence using Wilf-Zeilberger theory

Let $L(n, m; p)$ be the expectation of $X := X(n, m; p)$, which is the number of tosses until reaching (for the first time) either n Heads OR m Tails. Then,

Recurrence using Wilf-Zeilberger theory

Let $L(n, m; p)$ be the expectation of $X := X(n, m; p)$, which is the number of tosses until reaching (for the first time) either n Heads OR m Tails. Then,

$$\begin{aligned} L(n, m; p) = & \frac{(pn + pm - 2p + 2n - 2)}{n - 1} \cdot L(n - 1, m) \\ & - \frac{(2pn + 2pm - 4p + n - 1)}{n - 1} \cdot L(n - 2, m) \\ & + \frac{p(m - 2 + n)}{n - 1} \cdot L(n - 3, m), \end{aligned}$$

Recurrence using Wilf-Zeilberger theory

Let $L(n, m; p)$ be the expectation of $X := X(n, m; p)$, which is the number of tosses until reaching (for the first time) either n Heads OR m Tails. Then,

$$\begin{aligned} L(n, m; p) = & \frac{(pn + pm - 2p + 2n - 2)}{n - 1} \cdot L(n - 1, m) \\ & - \frac{(2pn + 2pm - 4p + n - 1)}{n - 1} \cdot L(n - 2, m) \\ & + \frac{p(m - 2 + n)}{n - 1} \cdot L(n - 3, m), \end{aligned}$$

with initial conditions:

$$\begin{aligned} L(1, 1) &= 1, L(1, 2) = 2 - p, L(1, 3) = p^2 - 3p + 3, L(2, 1) = p + 1, \\ L(2, 2) &= -2p^2 + 2p + 2, L(2, 3) = 3p^3 - 7p^2 + 3p + 3 \\ L(3, 1) &= p^2 + p + 1, L(3, 2) = -3p^3 + 2p^2 + 2p + 2, \\ L(3, 3) &= 6p^4 - 12p^3 + 3p^2 + 3p + 3. \end{aligned}$$

Example

Example

Using the recurrence, we can compute the expected coin tosses to get $2n$ Heads OR $3n$ Tails for $1 \leq n \leq 10$,

Example

Example

Using the recurrence, we can compute the expected coin tosses to get $2n$ Heads OR $3n$ Tails for $1 \leq n \leq 10$,

	Expected coin tosses
$n = 1$	3.272000000
$n = 2$	7.491773440
$n = 3$	11.90103592
$n = 4$	16.40588424
$n = 5$	20.97105153
$n = 6$	25.57874312
$n = 7$	30.21849553
$n = 8$	34.88353902
$n = 9$	39.56919552
$n = 10$	44.27207236

Table: The probability of getting Heads is $p = 2/5$.

Explicit Expression (an Heads OR bn Tails)

Fix positive integers a, b . Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then

Explicit Expression (an Heads OR bn Tails)

Fix positive integers a, b . Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then

$$L(an, bn, \frac{a}{a+b}) = (a+b)n \left(1 - \frac{((a+b)n)!}{(an)!(bn)!} \cdot \left(\frac{a^a b^b}{(a+b)^{a+b}} \right)^n \right).$$

Explicit Expression (an Heads OR bn Tails)

Fix positive integers a, b . Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then

$$L(an, bn, \frac{a}{a+b}) = (a+b)n \left(1 - \frac{((a+b)n)!}{(an)!(bn)!} \cdot \left(\frac{a^a b^b}{(a+b)^{a+b}} \right)^n \right).$$

Of course,

$$\frac{((a+b)n)!}{(an)!(bn)!} = \binom{(a+b)n}{an}.$$

Explicit Expression (an Heads OR bn Tails)

Fix positive integers a, b . Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then

$$L(an, bn, \frac{a}{a+b}) = (a+b)n \left(1 - \frac{((a+b)n)!}{(an)!(bn)!} \cdot \left(\frac{a^a b^b}{(a+b)^{a+b}} \right)^n \right).$$

Of course,

$$\frac{((a+b)n)!}{(an)!(bn)!} = \binom{(a+b)n}{an}.$$

Example

Let $a = 2, b = 3$ then $p = \frac{2}{5}$. If your goal is to flip the loaded coin until you reach 20,000 Heads OR 30,000 Tails, then

Explicit Expression (an Heads OR bn Tails)

Fix positive integers a, b . Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then

$$L(an, bn, \frac{a}{a+b}) = (a+b)n \left(1 - \frac{((a+b)n)!}{(an)!(bn)!} \cdot \left(\frac{a^a b^b}{(a+b)^{a+b}} \right)^n \right).$$

Of course,

$$\frac{((a+b)n)!}{(an)!(bn)!} = \binom{(a+b)n}{an}.$$

Example

Let $a = 2, b = 3$ then $p = \frac{2}{5}$. If your goal is to flip the loaded coin until you reach 20,000 Heads OR 30,000 Tails, then
the expected number of coin tosses is 50,182.09044.

Explicit Expression (an Heads OR bn Tails)

Fix positive integers a, b . Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then

$$L(an, bn, \frac{a}{a+b}) = (a+b)n \left(1 - \frac{((a+b)n)!}{(an)!(bn)!} \cdot \left(\frac{a^a b^b}{(a+b)^{a+b}} \right)^n \right).$$

Of course,

$$\frac{((a+b)n)!}{(an)!(bn)!} = \binom{(a+b)n}{an}.$$

One might be interested in the **asymptotics** of $L(an, bn, \frac{a}{a+b})$.

Asymptotics (an Heads OR bn Tails)

Fix positive integers a, b . Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then

Asymptotics (an Heads OR bn Tails)

Fix positive integers a, b . Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then $L(an, bn, \frac{a}{a+b})$ is asymptotically ($n \rightarrow \infty$),

$$(a+b)n \left(1 - \sqrt{\frac{a+b}{2ab\pi}} \cdot \frac{1}{\sqrt{n}} \right).$$

Pause

Summary so far!

Let p be the probability of getting Heads. If $X(n, m; p)$ is the number of tosses until reaching n Heads OR m Tails:

Summary so far!

Let p be the probability of getting Heads. If $X(n, m; p)$ is the number of tosses until reaching n Heads OR m Tails:

- ① Established (but not displayed) that the probability generating function $F(n, m; p)$ of $X(n, m; p)$ satisfies a linear recurrence with polynomial coefficients.

Summary so far!

Let p be the probability of getting Heads. If $X(n, m; p)$ is the number of tosses until reaching n Heads OR m Tails:

- 1 Established (but not displayed) that the probability generating function $F(n, m; p)$ of $X(n, m; p)$ satisfies a linear recurrence with polynomial coefficients.
- 2 The expectation denoted by $L(n, m; p)$ of $X(n, m; p)$ satisfies a third-order linear recurrence.

Summary so far!

Let p be the probability of getting Heads. If $X(n, m; p)$ is the number of tosses until reaching n Heads OR m Tails:

- 1 Established (but not displayed) that the probability generating function $F(n, m; p)$ of $X(n, m; p)$ satisfies a linear recurrence with polynomial coefficients.
- 2 The expectation denoted by $L(n, m; p)$ of $X(n, m; p)$ satisfies a third-order linear recurrence.
- 3 If $p = \frac{a}{a+b}$, then we have given an explicit expression for $L(an, bn; p)$ for positive integers a, b .

Summary so far!

Let p be the probability of getting Heads. If $X(n, m; p)$ is the number of tosses until reaching n Heads OR m Tails:

- 1 Established (but not displayed) that the probability generating function $F(n, m; p)$ of $X(n, m; p)$ satisfies a linear recurrence with polynomial coefficients.
- 2 The expectation denoted by $L(n, m; p)$ of $X(n, m; p)$ satisfies a third-order linear recurrence.
- 3 If $p = \frac{a}{a+b}$, then we have given an explicit expression for $L(an, bn; p)$ for positive integers a, b .
- 4 Given asymptotics for $L(an, bn; p)$ with $p = \frac{a}{a+b}$.

Connection to other work

Connection to work by Volkov and Wiktorsson

Volkov and Wiktorsson study the following game:

Connection to work by Volkov and Wiktorsson

Volkov and Wiktorsson study the following game:

Suppose two players play a game: In any given round, Player 1 wins with probability p and Player 2 wins with probability $q = 1 - p$.

The *first* player to reach n wins is the winner.

Connection to work by Volkov and Wiktorsson

Volkov and Wiktorsson study the following game:

Suppose two players play a game: In any given round, Player 1 wins with probability p and Player 2 wins with probability $q = 1 - p$.

The *first* player to reach n wins is the winner.

The prize? difference between their number of wins.

Connection to work by Volkov and Wiktorsson

Volkov and Wiktorsson study the following game:

Suppose two players play a game: In any given round, Player 1 wins with probability p and Player 2 wins with probability $q = 1 - p$.

The *first* player to reach n wins is the winner.

The prize? difference between their number of wins.

Back to our language: You toss a coin until you get n Heads OR n Tails where with probability p you get Heads.

Connection to work by Volkov and Wiktorsson (2)

Volkov and Wiktorsson prove the following:

Connection to work by Volkov and Wiktorsson (2)

Volkov and Wiktorsson prove the following:

Theorem (Volkov and Wiktorsson, 2025)

Let $\mathbb{E}_{n,p}$ be the expectation of (the number of Heads - the number of Tails) after $X(n, n; p)$ tosses i.e. once we reach either n Heads OR n Tails. Then,

Connection to work by Volkov and Wiktorsson (2)

Volkov and Wiktorsson prove the following:

Theorem (Volkov and Wiktorsson, 2025)

Let $\mathbb{E}_{n,p}$ be the expectation of (the number of Heads - the number of Tails) after $X(n, n; p)$ tosses i.e. once we reach either n Heads OR n Tails. Then,

$$\mathbb{E}_{n,p} = n(p - q) \sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ is the j th Catalan number.

Connection to work by Volkov and Wiktorsson (2)

Volkov and Wiktorsson prove the following:

Theorem (Volkov and Wiktorsson, 2025)

Let $\mathbb{E}_{n,p}$ be the expectation of (the number of Heads - the number of Tails) after $X(n,n; p)$ tosses i.e. once we reach either n Heads OR n Tails. Then,

$$\mathbb{E}_{n,p} = n(p - q) \sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ is the j th Catalan number.

Their result is equivalent to our result in the special case of reaching n Heads OR n Tails.

Wald's identity

Theorem (Wald's Identity)

Let X_1, X_2, \dots be independent, identically distributed random variables with finite mean μ and let $S_n (n \geq 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

Wald's identity

Theorem (Wald's Identity)

Let X_1, X_2, \dots be independent, identically distributed random variables with finite mean μ and let $S_n (n \geq 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \mathbb{E}[\tau].$$

Wald's identity

Theorem (Wald's Identity)

Let X_1, X_2, \dots be independent, identically distributed random variables with finite mean μ and let $S_n (n \geq 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n; p)$, and S_τ is the total profit after $X(n, n; p)$ coin tosses for a player betting on Heads

Wald's identity

Theorem (Wald's Identity)

Let X_1, X_2, \dots be independent, identically distributed random variables with finite mean μ and let $S_n (n \geq 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n; p)$, and S_τ is the total profit after $X(n, n; p)$ coin tosses for a player betting on Heads (let $+1$ represent Heads and -1 represent Tails).

Wald's identity

Theorem (Wald's Identity)

Let X_1, X_2, \dots be independent, identically distributed random variables with finite mean μ and let $S_n (n \geq 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n; p)$, and S_τ is the total profit after $X(n, n; p)$ coin tosses for a player betting on Heads (let $+1$ represent Heads and -1 represent Tails).

In addition, $\mu = p - q$ is the expected gain from 1 coin toss. Hence,

Wald's identity

Theorem (Wald's Identity)

Let X_1, X_2, \dots be independent, identically distributed random variables with finite mean μ and let $S_n (n \geq 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n; p)$, and S_τ is the total profit after $X(n, n; p)$ coin tosses for a player betting on Heads (let $+1$ represent Heads and -1 represent Tails).

In addition, $\mu = p - q$ is the expected gain from 1 coin toss. Hence,

$$\mathbb{E}[S_\tau] = \mu \mathbb{E}[\tau] = (p - q)L(n, n; p)$$

Wald's identity

Theorem (Wald's Identity)

Let X_1, X_2, \dots be independent, identically distributed random variables with finite mean μ and let $S_n (n \geq 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n; p)$, and S_τ is the total profit after $X(n, n; p)$ coin tosses for a player betting on Heads (let $+1$ represent Heads and -1 represent Tails).

In addition, $\mu = p - q$ is the expected gain from 1 coin toss. Hence,

$$\mathbb{E}[S_\tau] = \mu \mathbb{E}[\tau] = (p - q)L(n, n; p)$$

Takeaway:

total average profit = (average gain per toss)(average number of tosses)

Theorem (Volkov and Wiktorsson, 2025)

$$(p - q)L(n, n; p) = \mathbb{E}_{n,p} = n(p - q) \sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ is the j th Catalan number.

Back to Volkov and Wiktorsson

Theorem (Volkov and Wiktorsson, 2025)

$$(p - q)L(n, n; p) = \mathbb{E}_{n,p} = n(p - q) \sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ is the j th Catalan number.

This is equivalent to

$$L(n, n; p) = n \sum_{j=0}^{n-1} C_j (pq)^j.$$

Back to Volkov and Wiktorsson

Theorem (Volkov and Wiktorsson, 2025)

$$(p - q)L(n, n; p) = \mathbb{E}_{n,p} = n(p - q) \sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ is the j th Catalan number.

This is equivalent to

$$L(n, n; p) = n \sum_{j=0}^{n-1} C_j (pq)^j.$$

In the fair case, $p = \frac{1}{2}$, we can verify that with $a = b = 1$, we get

$$L(n, n; \tfrac{1}{2}) = 2n \left(1 - \frac{(2n)!}{n!^2} 4^{-n} \right).$$

Results (2)


Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.

Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.


If your goal is to reach either 200 Heads OR 10 Tails,



Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.


If your goal is to reach either 200 Heads OR 50 Tails,



Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.


If your goal is to reach either 200 Heads OR 100 Tails,



Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.


If your goal is to reach either 200 Heads OR 150 Tails,



Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.

If your goal is to reach either 200 Heads OR 200 Tails,



The r th factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

The r th factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$\left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} \left(\frac{1}{2}\right)^h x^{h+n}.$$

The r th factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$\left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} \left(\frac{1}{2}\right)^h x^{h+n}.$$

Now, we introduce the r th factorial moment of a random variable X ,

The r th factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$\left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} \left(\frac{1}{2}\right)^h x^{h+n}.$$

Now, we introduce the r th factorial moment of a random variable X ,

$$\mathbb{E}[X(X-1)\dots(X-r+1)] = r! \mathbb{E}\left[\binom{X}{r}\right].$$

The r th factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$\left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} \left(\frac{1}{2}\right)^h x^{h+n}.$$

Now, we introduce the r th factorial moment of a random variable X ,

$$\mathbb{E}[X(X-1)\dots(X-r+1)] = r! \mathbb{E} \left[\binom{X}{r} \right].$$

Let $A(n, r)$ be the r th factorial moment of our random variable $X = X(n, n; \frac{1}{2})$. Then,

The r th factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$\left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} \left(\frac{1}{2}\right)^h x^{h+n}.$$

Now, we introduce the r th factorial moment of a random variable X ,

$$\mathbb{E}[X(X-1)\dots(X-r+1)] = r! \mathbb{E}\left[\binom{X}{r}\right].$$

Let $A(n, r)$ be the r th factorial moment of our random variable $X = X(n, n; \frac{1}{2})$. Then,

$$A(n, r) = \left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} \left(\frac{1}{2}\right)^h$$

The r th factorial moment (n Heads OR n Tails) (2)

Let $A(n, r)$ be the r th factorial moment of our random variable $X = X(n, n; \frac{1}{2})$. Then,

$$A(n, r) = \left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} \left(\frac{1}{2}\right)^h.$$

The r th factorial moment (n Heads OR n Tails) (2)

Let $A(n, r)$ be the r th factorial moment of our random variable $X = X(n, n; \frac{1}{2})$. Then,

$$A(n, r) = \left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} \left(\frac{1}{2}\right)^h.$$

Remark: By fixing (small) r , we can use Maple to get a closed-form expression in n .

The r th factorial moment (n Heads OR n Tails) (2)

Let $A(n, r)$ be the r th factorial moment of our random variable $X = X(n, n; \frac{1}{2})$. Then,

$$A(n, r) = \left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} \left(\frac{1}{2}\right)^h.$$

Remark: By fixing (small) r , we can use Maple to get a closed-form expression in n .

Remark (2): There is no closed-form expression in terms of r .

The r th factorial moment (n Heads OR n Tails) (2)

Let $A(n, r)$ be the r th factorial moment of our random variable $X = X(n, n; \frac{1}{2})$. Then,

$$A(n, r) = \left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} \left(\frac{1}{2}\right)^h.$$

Remark: By fixing (small) r , we can use Maple to get a closed-form expression in n .

Remark (2): There is no closed-form expression in terms of r . Once again, we rely on linear recurrences. We use the Zeilberger algorithm!

The r th factorial moment (n Heads OR n Tails) (2)

Let $A(n, r)$ be the r th factorial moment of our random variable $X = X(n, n; \frac{1}{2})$. Then,

$$A(n, r) = \left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} \left(\frac{1}{2}\right)^h.$$

So,

$$\begin{aligned} A(n, r) &= 2nA(n, r-1) \\ &\quad + (r-1)(r-2)A(n, r-2) - 4n \binom{2n-1}{r-2} (r-2)! B_n \end{aligned}$$

The r th factorial moment (n Heads OR n Tails) (2)

Let $A(n, r)$ be the r th factorial moment of our random variable $X = X(n, n; \frac{1}{2})$. Then,

$$A(n, r) = \left(\frac{1}{2}\right)^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} \left(\frac{1}{2}\right)^h.$$

So,

$$\begin{aligned} A(n, r) &= 2nA(n, r-1) \\ &\quad + (r-1)(r-2)A(n, r-2) - 4n \binom{2n-1}{r-2} (r-2)! B_n \end{aligned}$$

where

$$A(n, 1) = 2n - 2B_n, \quad A(n, 2) = 4n^2 - 8nB_n, \quad B_n := \frac{n}{4^n} \binom{2n}{n}.$$

Central moments and scaled central moments

Recall that $A(n, 1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n} \binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

Central moments and scaled central moments

Recall that $A(n, 1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n} \binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

$$\mathbb{E}[(X - \mu)^r] = \sum_{i=0}^r \binom{r}{i} (-\mu)^{r-i} \mathbb{E}[X^i].$$

Central moments and scaled central moments

Recall that $A(n, 1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n} \binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

$$\mathbb{E}[(X - \mu)^r] = \sum_{i=0}^r \binom{r}{i} (-\mu)^{r-i} \mathbb{E}[X^i].$$

The **scaled central moments** (scaling by σ^r) is given by

Central moments and scaled central moments

Recall that $A(n, 1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n} \binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

$$\mathbb{E}[(X - \mu)^r] = \sum_{i=0}^r \binom{r}{i} (-\mu)^{r-i} \mathbb{E}[X^i].$$

The **scaled central moments** (scaling by σ^r) is given by

$$\frac{\mathbb{E}[(X - \mu)^r]}{\sigma^r} = \frac{\mathbb{E}[(X - (2n - 2\frac{n}{4^n} \binom{2n}{n}))^r]}{\sigma^r},$$

where $\sigma^2 = \mathbb{E}[(X - \mu)^2]$ is the variance.

Central moments and scaled central moments

Recall that $A(n, 1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n} \binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

$$\mathbb{E}[(X - \mu)^r] = \sum_{i=0}^r \binom{r}{i} (-\mu)^{r-i} \mathbb{E}[X^i].$$

The **scaled central moments** (scaling by σ^r) is given by

$$\frac{\mathbb{E}[(X - \mu)^r]}{\sigma^r} = \frac{\mathbb{E}[(X - (2n - 2\frac{n}{4^n} \binom{2n}{n}))^r]}{\sigma^r},$$

where $\sigma^2 = \mathbb{E}[(X - \mu)^2]$ is the variance.

Taking $n \rightarrow \infty$, we obtain the same scaled central moments of the **negative binomial distribution!**

Summary

- 1 We provided linear recurrences for the expected number of coin tosses until reaching n Heads OR m Tails.

Summary

- ① We provided linear recurrences for the expected number of coin tosses until reaching n Heads OR m Tails.
- ② Some special cases included an Heads OR bn Tails.

Summary

- ① We provided linear recurrences for the expected number of coin tosses until reaching n Heads OR m Tails.
- ② Some special cases included an Heads OR bn Tails.
- ③ For the case n Heads OR n Tails, we show that there is convergence of moments!

Summary

- ① We provided linear recurrences for the expected number of coin tosses until reaching n Heads OR m Tails.
- ② Some special cases included an Heads OR bn Tails.
- ③ For the case n Heads OR n Tails, we show that there is convergence of moments!
- ④ In our paper, we also discuss the case for n Heads AND m Tails.

Summary

- ① We provided linear recurrences for the expected number of coin tosses until reaching n Heads OR m Tails.
- ② Some special cases included an Heads OR bn Tails.
- ③ For the case n Heads OR n Tails, we show that there is convergence of moments!
- ④ In our paper, we also discuss the case for n Heads AND m Tails.

If you would like to read the paper:

<https://marti310.github.io/research.html>

Thank you!