How many coin tosses would you need until you get n Heads or m Tails?

Lucy Martinez

Rutgers University

Graduate Student Combinatorics Conference 2026

–Joint work with Svante Janson and Doron Zeilberger–

Background and History

Problem of points

In 1654, Fermat and Pascal were interested in the following game.

Problem of points

In 1654, Fermat and Pascal were interested in the following game.

Two players compete in a fair game such as "first to win ten rounds takes the prize". Suppose that the game stops early, for example at a score of 3 to 4. How should the players decide how to split the prize?

Problem of points

In 1654, Fermat and Pascal were interested in the following game.

Two players compete in a fair game such as "first to win ten rounds takes the prize". Suppose that the game stops early, for example at a score of 3 to 4. How should the players decide how to split the prize?

The mail correspondence between Fermat and Pascal laid the foundation for what is now known as probability theory.

Problem of points (2)

We can rephrase the problem of points as follows.

Problem of points (2)

We can rephrase the problem of points as follows.

- Suppose that player 1 has n rounds left to win and player 2 has m rounds to win

Problem of points (2)

We can rephrase the problem of points as follows.

- Suppose that player 1 has n rounds left to win and player 2 has m rounds to win
- Thus, we can translate the problem of points into the following scenario

Problem of points (2)

We can rephrase the problem of points as follows.

- Suppose that player 1 has n rounds left to win and player 2 has m rounds to win
- Thus, we can translate the problem of points into the following scenario
- Suppose we toss a fair coin. What is the probability that we reach n Heads OR m Tails?

Problem of points (2)

We can rephrase the problem of points as follows.

- Suppose that player 1 has n rounds left to win and player 2 has m rounds to win
- Thus, we can translate the problem of points into the following scenario
- Suppose we toss a fair coin. What is the probability that we reach n Heads OR m Tails?

Convention: For the rest of the talk, p represents the probability of getting Heads and $q := 1 - p$ represents the probability of getting Tails.

Probability generating function until reaching n Heads

Consider tossing a coin *until* we reach n Heads.

Probability generating function until reaching n Heads

Consider tossing a coin *until* we reach n Heads. The maximum number of possible more coin tosses is $n + m - 1$.

Probability generating function until reaching n Heads

Consider tossing a coin *until* we reach n Heads. The maximum number of possible more coin tosses is $n + m - 1$. Let $q = 1 - p$, then the probability generating function is

Probability generating function until reaching n Heads

Consider tossing a coin *until* we reach n Heads. The maximum number of possible more coin tosses is $n + m - 1$. Let $q = 1 - p$, then the probability generating function is

$$\sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (px)^n (qx)^t$$

Probability generating function until reaching n Heads

Consider tossing a coin *until* we reach n Heads. The maximum number of possible more coin tosses is $n+m-1$. Let $q = 1-p$, then the probability generating function is

$$\sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (px)^n (qx)^t = \binom{n-1}{n-1} (px)^n (qx)^0 + \binom{n}{n-1} (px)^n (qx)^1 + \cdots + \binom{n+m-2}{n-1} (px)^n (qx)^{m-1}$$

Probability generating function until reaching n Heads

Consider tossing a coin *until* we reach n Heads. The maximum number of possible more coin tosses is $n+m-1$. Let $q = 1-p$, then the probability generating function is

$$\begin{aligned}\sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (px)^n (qx)^t &= \binom{n-1}{n-1} (px)^n (qx)^0 + \binom{n}{n-1} (px)^n (qx)^1 \\ &\quad + \cdots + \binom{n+m-2}{n-1} (px)^n (qx)^{m-1} \\ &= A(n,m,x)\end{aligned}$$

One of the (many) solutions provided by Fermat and Pascal was to split the prize with Player 1 receiving the above expression, and Player 2 receiving $1 - A(n,m,1)$.

Probability generating function until reaching n Heads

Consider tossing a coin *until* we reach n Heads. The maximum number of possible more coin tosses is $n+m-1$. Let $q = 1-p$, then the probability generating function is

$$\sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (px)^n (qx)^t = \binom{n-1}{n-1} (px)^n (qx)^0 + \binom{n}{n-1} (px)^n (qx)^1 + \cdots + \binom{n+m-2}{n-1} (px)^n (qx)^{m-1}$$

Remark: Fermat and Pascal were only interested in the probability of getting n Heads vs. getting m Tails.

Probability generating function until reaching n Heads

Consider tossing a coin *until* we reach n Heads. The maximum number of possible more coin tosses is $n+m-1$. Let $q = 1-p$, then the probability generating function is

$$\sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (px)^n (qx)^t = \binom{n-1}{n-1} (px)^n (qx)^0 + \binom{n}{n-1} (px)^n (qx)^1 + \cdots + \binom{n+m-2}{n-1} (px)^n (qx)^{m-1}$$

Remark: If we extend the above expression to infinity, then it has the following *closed-form*: $\left(\frac{px}{1-qx}\right)^n$.

Computations

Probability generating function (n Heads OR m Tails)

$X := X(n, m;\ p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Probability generating function (n Heads OR m Tails)

$X := X(n, m;\ p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Let $F(n, m;\ p)$ be the probability generating function of the random variable X in terms of x, then

Probability generating function (n Heads OR m Tails)

$X := X(n, m;\ p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Let $F(n, m;\ p)$ be the probability generating function of the random variable X in terms of x, then

$$F(n, m;\ p)(x) = (px)^n \sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (qx)^t + (qx)^m \sum_{h=0}^{n-1} \binom{m+h-1}{m-1} (px)^h.$$

Probability generating function (n Heads OR m Tails)

$X := X(n, m;\ p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Let $F(n, m;\ p)$ be the probability generating function of the random variable X in terms of x, then

$$F(n, m;\ p)(x) = (px)^n \sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (qx)^t + (qx)^m \sum_{h=0}^{n-1} \binom{m+h-1}{m-1} (px)^h.$$

Remark: $F(n, m;\ p)$ does not have a closed-form but it *does* have a linear recurrence with polynomial coefficients.

Probability generating function (n Heads OR m Tails)

$X := X(n, m;\ p)$: The number of tosses until reaching (for the first time) either n Heads OR m Tails, where with probability p you get Heads.

Let $F(n, m;\ p)$ be the probability generating function of the random variable X in terms of x, then

$$F(n, m;\ p)(x) = (px)^n \sum_{t=0}^{m-1} \binom{n+t-1}{n-1} (qx)^t + (qx)^m \sum_{h=0}^{n-1} \binom{m+h-1}{m-1} (px)^h.$$

Remark: $F(n, m;\ p)$ does not have a closed-form but it *does* have a linear recurrence with polynomial coefficients. We do not show it here for sake of space.

Computing expected coin tosses (n Heads OR m Tails)

To compute *some* data, we can take the derivative of the previous sum $F(n, m;\ p)(x)$ with respect to x and then plug in $x = 1$,

Computing expected coin tosses (n Heads OR m Tails)

To compute *some* data, we can take the derivative of the previous sum $F(n, m;\ p)(x)$ with respect to x and then plug in $x = 1$,

$$p^n \sum_{t=0}^{m-1} (n+t) \binom{n-1+t}{n-1} q^t + q^m \sum_{h=0}^{n-1} (m+h) \binom{m-1+h}{m-1} p^h, \quad q = 1-p.$$

Computing expected coin tosses (n Heads OR m Tails)

To compute *some* data, we can take the derivative of the previous sum $F(n, m;\ p)(x)$ with respect to x and then plug in $x = 1$,

$$p^n \sum_{t=0}^{m-1} (n+t) \binom{n-1+t}{n-1} q^t + q^m \sum_{h=0}^{n-1} (m+h) \binom{m-1+h}{m-1} p^h, \quad q = 1-p.$$

Remark: This becomes computationally slow. Using Wilf-Zeilberger algorithmic proof theory, we will obtain a third-order linear recurrence!

Results (1)

Recurrence using Wilf-Zeilberger theory

Let $L(n, m;\ p)$ be the expectation of $X := X(n, m;\ p)$, which is the number of tosses until reaching (for the first time) either n Heads OR m Tails. Then,

Recurrence using Wilf-Zeilberger theory

Let $L(n, m;\ p)$ be the expectation of $X := X(n, m;\ p)$, which is the number of tosses until reaching (for the first time) either n Heads OR m Tails. Then,

$$\begin{aligned} L(n, m;\ p) = & \frac{(pn + pm - 2p + 2n - 2)}{n-1} \cdot L\left(n-1, m\right) \\ & - \frac{(2pn + 2pm - 4p + n - 1)}{n-1} \cdot L\left(n-2, m\right) \\ & + \frac{p\left(m - 2 + n\right)}{n-1} \cdot L\left(n-3, m\right), \end{aligned}$$

Recurrence using Wilf-Zeilberger theory

Let $L(n, m;\ p)$ be the expectation of $X := X(n, m;\ p)$, which is the number of tosses until reaching (for the first time) either n Heads OR m Tails. Then,

$$\begin{aligned} L(n, m;\ p) = & \frac{(pn + pm - 2p + 2n - 2)}{n-1} \cdot L(n-1, m) \\ & - \frac{(2pn + 2pm - 4p + n - 1)}{n-1} \cdot L(n-2, m) \\ & + \frac{p(m - 2 + n)}{n-1} \cdot L(n-3, m), \end{aligned}$$

with initial conditions:

$$\begin{aligned} & L(1,1) = 1, L(1,2) = 2 - p, L(1,3) = p^2 - 3p + 3, L(2,1) = p + 1, \\ & L(2,2) = -2p^2 + 2p + 2, L(2,3) = 3p^3 - 7p^2 + 3p + 3 \\ & L(3,1) = p^2 + p + 1, L(3,2) = -3p^3 + 2p^2 + 2p + 2, \\ & L(3,3) = 6p^4 - 12p^3 + 3p^2 + 3p + 3. \end{aligned}$$

Example

Example

Using the recurrence, we can compute the expected coin tosses to get $2n$ Heads OR $3n$ Tails for $1 \leqslant n \leqslant 10$,

Example

Example

Using the recurrence, we can compute the expected coin tosses to get $2n$ Heads OR $3n$ Tails for $1 \leqslant n \leqslant 10$,

	Expected coin tosses
$n = 1$	3.272000000
$n = 2$	7.491773440
$n = 3$	11.90103592
$n = 4$	16.40588424
$n = 5$	20.97105153
$n = 6$	25.57874312
$n = 7$	30.21849553
$n = 8$	34.88353902
$n = 9$	39.56919552
$n = 10$	44.27207236

Table: The probability of getting Heads is $p = 2/5$.

Explicit Expression (an Heads OR bn Tails)

Fix positive integers a, b. Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then the expected duration is given by

Explicit Expression (an Heads OR bn Tails)

Fix positive integers a, b. Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then the expected duration is given by

$$L(an, bn, \ \tfrac{a}{a+b}) = (a+b)n \left(1 - \binom{(a+b)n}{an} \cdot \left(\frac{a^a b^b}{(a+b)^{a+b}}\right)^n\right).$$

Explicit Expression (an Heads OR bn Tails)

Fix positive integers a, b. Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then the expected duration is given by

$$L(an, bn, \tfrac{a}{a+b}) = (a+b)n\left(1 - \binom{(a+b)n}{an} \cdot \left(\frac{a^a b^b}{(a+b)^{a+b}}\right)^n\right).$$

One might be interested in the **asymptotics** of $L(an, bn, \frac{a}{a+b})$.

Asymptotics (an Heads OR bn Tails)

Fix positive integers a, b. Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then we have the following approximation:

Asymptotics (an Heads OR bn Tails)

Fix positive integers a, b. Let $p = \frac{a}{a+b}$ be the probability of getting Heads. If your goal is to reach either an Heads OR bn Tails, then we have the following approximation:

$$L(an, bn, \ \tfrac{a}{a+b}) \sim (a+b)n \left(1 - \sqrt{\frac{a+b}{2ab\pi}} \cdot \frac{1}{\sqrt{n}}\right), \qquad n \to \infty.$$

Connection to other work

Connection to work by Volkov and Wiktorsson

Volkov and Wiktorsson study the following game:

Connection to work by Volkov and Wiktorsson

Volkov and Wiktorsson study the following game:

Suppose two players play a game: In any given round, Player 1 wins with probability p and Player 2 wins with probability $q = 1 - p$.

Goal: The *first* player to reach n wins is the winner.

Connection to work by Volkov and Wiktorsson

Volkov and Wiktorsson study the following game:

Suppose two players play a game: In any given round, Player 1 wins with probability p and Player 2 wins with probability $q = 1 - p$.

Goal: The *first* player to reach n wins is the winner.

The prize? Difference between their number of wins.

Connection to work by Volkov and Wiktorsson

Volkov and Wiktorsson study the following game:

Suppose two players play a game: In any given round, Player 1 wins with probability p and Player 2 wins with probability $q = 1 - p$.

Goal: The *first* player to reach n wins is the winner.

The prize? Difference between their number of wins.

Back to our language: You toss a coin until you get n Heads OR n Tails, where with probability p you get Heads.

Connection to work by Volkov and Wiktorsson (2)

Volkov and Wiktorsson prove the following:

Connection to work by Volkov and Wiktorsson (2)

Volkov and Wiktorsson prove the following:

Theorem (Volkov and Wiktorsson, 2025)

Let $\mathbb{E}_{n,p}$ be the expectation of (the number of Heads - the number of Tails) after $X(n, n;\ p)$ tosses i.e. once we reach either n Heads OR n Tails. Then,

Connection to work by Volkov and Wiktorsson (2)

Volkov and Wiktorsson prove the following:

Theorem (Volkov and Wiktorsson, 2025)

Let $\mathbb{E}_{n,p}$ be the expectation of (the number of Heads - the number of Tails) after $X(n, n;\ p)$ tosses i.e. once we reach either n Heads OR n Tails. Then,

$$\mathbb{E}_{n,p} = n(p-q)\sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ is the jth Catalan number.

Connection to work by Volkov and Wiktorsson (2)

Volkov and Wiktorsson prove the following:

Theorem (Volkov and Wiktorsson, 2025)

Let $\mathbb{E}_{n,p}$ be the expectation of (the number of Heads - the number of Tails) after $X(n, n;\ p)$ tosses i.e. once we reach either n Heads OR n Tails. Then,

$$\mathbb{E}_{n,p} = n(p-q)\sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ is the jth Catalan number.

Their result is equivalent to our result in the special case of reaching n Heads OR n Tails.

Wald's identity

Theorem (Wald's Identity)

Let $X_1, X_2, \ldots$ be independent, identically distributed random variables with finite mean μ and let $S_n (n \geqslant 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

Wald's identity

Theorem (Wald's Identity)

Let $X_1, X_2, \ldots$ be independent, identically distributed random variables with finite mean μ and let $S_n (n \geqslant 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \, \mathbb{E}[\tau].$$

Wald's identity

Theorem (Wald's Identity)

Let $X_1, X_2, \ldots$ be independent, identically distributed random variables with finite mean μ and let $S_n (n \geqslant 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \, \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n;\ p)$, and S_τ is the total profit after $X = X(n, n;\ p)$ coin tosses for a player betting on Heads

Wald's identity

Theorem (Wald's Identity)

Let $X_1, X_2, \ldots$ be independent, identically distributed random variables with finite mean μ and let $S_n (n \geqslant 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \, \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n;\ p)$, and S_τ is the total profit after $X = X(n, n;\ p)$ coin tosses for a player betting on Heads (let $+1$ represent Heads and -1 represent Tails).

Wald's identity

Theorem (Wald's Identity)

Let $X_1, X_2, \ldots$ be independent, identically distributed random variables with finite mean μ and let $S_n (n \geqslant 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \, \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n;\ p)$, and S_τ is the total profit after $X = X(n, n;\ p)$ coin tosses for a player betting on Heads (let $+1$ represent Heads and -1 represent Tails).
In addition, $\mu = p - q$ is the expected gain from 1 coin toss. Hence,

Wald's identity

Theorem (Wald's Identity)

Let $X_1, X_2, \ldots$ be independent, identically distributed random variables with finite mean μ and let $S_n (n \geqslant 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu\, \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n;\ p)$, and S_τ is the total profit after $X = X(n, n;\ p)$ coin tosses for a player betting on Heads (let $+1$ represent Heads and -1 represent Tails).
In addition, $\mu = p - q$ is the expected gain from 1 coin toss. Hence,

$$\mathbb{E}[S_\tau] = \mu\, \mathbb{E}[\tau] = (p - q) L(n, n;\ p)$$

Wald's identity

Theorem (Wald's Identity)

Let $X_1, X_2, \ldots$ be independent, identically distributed random variables with finite mean μ and let $S_n (n \geqslant 1)$, denote the partial sums. Suppose that τ is a stopping time. If $\mathbb{E}[\tau] < \infty$, then

$$\mathbb{E}[S_\tau] = \mu \, \mathbb{E}[\tau].$$

In our case: $\tau = X(n, n;\ p)$, and S_τ is the total profit after $X = X(n, n;\ p)$ coin tosses for a player betting on Heads (let $+1$ represent Heads and -1 represent Tails).
In addition, $\mu = p - q$ is the expected gain from 1 coin toss. Hence,

$$\mathbb{E}[S_\tau] = \mu \, \mathbb{E}[\tau] = (p - q) L(n, n;\ p)$$

Takeaway:
total average profit= (average gain per toss)(average number of tosses)

Back to Volkov and Wiktorsson

From Wald's identity, $\mathbb{E}_{n,p} = (p-q)L(n,n;\ p)$.

Back to Volkov and Wiktorsson

From Wald's identity, $\mathbb{E}_{n,p} = (p-q)L(n,n;\ p)$.

Theorem (Volkov and Wiktorsson, 2025)

$$(p-q)L(n,n;\ p) = \mathbb{E}_{n,p} = n(p-q)\sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ *is the* j*th Catalan number.*

Back to Volkov and Wiktorsson

Theorem (Volkov and Wiktorsson, 2025)

$$(p-q)L(n,n;\ p) = \mathbb{E}_{n,p} = n(p-q)\sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ *is the* j*th Catalan number.*

This is equivalent to

$$L(n,n;\ p) = n\sum_{j=0}^{n-1} C_j (pq)^j.$$

Back to Volkov and Wiktorsson

Theorem (Volkov and Wiktorsson, 2025)

$$(p-q)L(n,n;\ p) = \mathbb{E}_{n,p} = n(p-q)\sum_{j=0}^{n-1} C_j (pq)^j,$$

where $C_j = \frac{(2j)!}{j!(j+1)!}$ *is the* j*th Catalan number.*

This is equivalent to

$$L(n,n;\ p) = n\sum_{j=0}^{n-1} C_j (pq)^j.$$

In the fair case, $p = \frac{1}{2}$, we can verify that with $a = b = 1$, we get

$$L(n,n;\ \tfrac{1}{2}) = 2n\left(1 - \frac{(2n)!}{n!^2}4^{-n}\right).$$

Summary

1. We provided linear recurrences for the probability and expected number of coin tosses until reaching n Heads OR m Tails.

Summary

1. We provided linear recurrences for the probability and expected number of coin tosses until reaching n Heads OR m Tails.
2. Some special cases included an Heads OR bn Tails.

Summary

1. We provided linear recurrences for the probability and expected number of coin tosses until reaching n Heads OR m Tails.
2. Some special cases included an Heads OR bn Tails.
3. Other results: We show that there is convergence of moments for the case n Heads OR n Tails.

Summary

1. We provided linear recurrences for the probability and expected number of coin tosses until reaching n Heads OR m Tails.
2. Some special cases included an Heads OR bn Tails.
3. Other results: We show that there is convergence of moments for the case n Heads OR n Tails.
4. In our paper, we **also** prove similar results for the case of reaching n Heads AND m Tails.

Summary

1. We provided linear recurrences for the probability and expected number of coin tosses until reaching n Heads OR m Tails.
2. Some special cases included an Heads OR bn Tails.
3. Other results: We show that there is convergence of moments for the case n Heads OR n Tails.
4. In our paper, we **also** prove similar results for the case of reaching n Heads AND m Tails.

If you would like to read the paper:

`https://marti310.github.io/research.html`

Thank you!

Results (2)

Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.

Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.

If your goal is to reach either 200 Heads OR 10 Tails,



Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.

If your goal is to reach either 200 Heads OR 50 Tails,



Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.

If your goal is to reach either 200 Heads OR 100 Tails,



Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.

If your goal is to reach either 200 Heads OR 150 Tails,



Expected coin tosses figures for $n = m$

Using the recurrence for the expectation, we can compute the following with various p values.

If your goal is to reach either 200 Heads OR 200 Tails,



The rth factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

The rth factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$(\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} (\tfrac{1}{2})^h x^{h+n}.$$

The rth factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$(\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} (\tfrac{1}{2})^h x^{h+n}.$$

Now, we introduce the rth factorial moment of a random variable X,

The rth factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$(\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} (\tfrac{1}{2})^h x^{h+n}.$$

Now, we introduce the rth factorial moment of a random variable X,

$$\mathbb{E}\left[X(X-1)\ldots(X-r+1)\right] = r!\,\mathbb{E}\left[\binom{X}{r}\right].$$

The rth factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$(\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} (\tfrac{1}{2})^h x^{h+n}.$$

Now, we introduce the rth factorial moment of a random variable X,

$$\mathbb{E}\left[X(X-1)\ldots(X-r+1)\right] = r!\,\mathbb{E}\left[\binom{X}{r}\right].$$

Let $A(n,r)$ be the rth factorial moment of our random variable $X = X(n,n;\ \frac{1}{2})$. Then,

The rth factorial moment (n Heads OR n Tails)

The probability generating function for the number of tosses until a fair coin reaches n Heads OR n Tails is

$$(\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} (\tfrac{1}{2})^h x^{h+n}.$$

Now, we introduce the rth factorial moment of a random variable X,

$$\mathbb{E}\left[X(X-1)\ldots(X-r+1)\right] = r!\,\mathbb{E}\left[\binom{X}{r}\right].$$

Let $A(n,r)$ be the rth factorial moment of our random variable $X = X(n,n;\ \frac{1}{2})$. Then,

$$A(n,r) = (\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} (\tfrac{1}{2})^h$$

The rth factorial moment (n Heads OR n Tails) (2)

Let $A(n,r)$ be the rth factorial moment of our random variable $X = X(n,n;\ \frac{1}{2})$. Then,

$$A(n,r) = (\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} (\tfrac{1}{2})^h.$$

The rth factorial moment (n Heads OR n Tails) (2)

Let $A(n,r)$ be the rth factorial moment of our random variable $X = X(n,n;\ \frac{1}{2})$. Then,

$$A(n,r) = (\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} (\tfrac{1}{2})^h.$$

Remark: By fixing (small) r, we can use Maple to get a closed-form expression in n.

The rth factorial moment (n Heads OR n Tails) (2)

Let $A(n,r)$ be the rth factorial moment of our random variable $X = X(n,n;\ \frac{1}{2})$. Then,

$$A(n,r) = (\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} (\tfrac{1}{2})^h.$$

Remark: By fixing (small) r, we can use Maple to get a closed-form expression in n.
Remark (2): There is no closed-form expression in terms of r.

The rth factorial moment (n Heads OR n Tails) (2)

Let $A(n,r)$ be the rth factorial moment of our random variable $X = X(n,n;\ \frac{1}{2})$. Then,

$$A(n,r) = (\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} (\tfrac{1}{2})^h.$$

Remark: By fixing (small) r, we can use Maple to get a closed-form expression in n.
Remark (2): There is no closed-form expression in terms of r. Once again, we rely on linear recurrences. We use the Zeilberger algorithm!

The rth factorial moment (n Heads OR n Tails) (2)

Let $A(n,r)$ be the rth factorial moment of our random variable $X = X(n,n;\ \frac{1}{2})$. Then,

$$A(n,r) = (\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} (\tfrac{1}{2})^h.$$

So,

$$\begin{aligned} A(n,r) &= 2nA(n,r-1) \\ &\qquad + (r-1)(r-2)A(n,r-2) - 4n\binom{2n-1}{r-2}(r-2)!\,B_n \end{aligned}$$

The rth factorial moment (n Heads OR n Tails) (2)

Let $A(n,r)$ be the rth factorial moment of our random variable $X = X(n,n;\ \frac{1}{2})$. Then,

$$A(n,r) = (\tfrac{1}{2})^{n-1} \sum_{h=0}^{n-1} \binom{h+n-1}{n-1} r! \binom{h+n}{r} (\tfrac{1}{2})^h.$$

So,

$$\begin{aligned} A(n,r) &= 2nA(n,r-1) \\ &\qquad + (r-1)(r-2)A(n,r-2) - 4n\binom{2n-1}{r-2}(r-2)!\, B_n \end{aligned}$$

where

$$A(n,1) = 2n - 2B_n, \quad A(n,2) = 4n^2 - 8nB_n, \quad B_n := \frac{n}{4^n}\binom{2n}{n}.$$

Central moments and scaled central moments

Recall that $A(n,1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n}\binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

Central moments and scaled central moments

Recall that $A(n,1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n}\binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

$$\mathbb{E}[(X-\mu)^r] = \sum_{i=0}^{r} \binom{r}{i} (-\mu)^{r-i}\, \mathbb{E}[X^i].$$

Central moments and scaled central moments

Recall that $A(n,1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n}\binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

$$\mathbb{E}[(X-\mu)^r] = \sum_{i=0}^{r} \binom{r}{i} (-\mu)^{r-i}\, \mathbb{E}[X^i].$$

The **scaled central moments** (scaling by σ^r) is given by

Central moments and scaled central moments

Recall that $A(n,1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n}\binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

$$\mathbb{E}[(X-\mu)^r] = \sum_{i=0}^{r} \binom{r}{i} (-\mu)^{r-i}\, \mathbb{E}[X^i].$$

The **scaled central moments** (scaling by σ^r) is given by

$$\frac{\mathbb{E}[(X-\mu)^r]}{\sigma^r} = \frac{\mathbb{E}\left[\left(X - (2n - 2\frac{n}{4^n}\binom{2n}{n})\right)^r\right]}{\sigma^r},$$

where $\sigma^2 = \mathbb{E}[(X-\mu)^2]$ is the variance.

Central moments and scaled central moments

Recall that $A(n,1) = \mathbb{E}[X] = 2n - 2\frac{n}{4^n}\binom{2n}{n}$. Let $\mu = \mathbb{E}[X]$, then the **central moment** is given by

$$\mathbb{E}[(X-\mu)^r] = \sum_{i=0}^{r} \binom{r}{i} (-\mu)^{r-i}\, \mathbb{E}[X^i].$$

The **scaled central moments** (scaling by σ^r) is given by

$$\frac{\mathbb{E}[(X-\mu)^r]}{\sigma^r} = \frac{\mathbb{E}\left[\left(X - (2n - 2\frac{n}{4^n}\binom{2n}{n})\right)^r\right]}{\sigma^r},$$

where $\sigma^2 = \mathbb{E}[(X-\mu)^2]$ is the variance.
Taking $n \to \infty$, we obtain the same scaled central moments of the **negative binomial distribution!**