

Computational Methods Assignment 2 - Interpolation

Martik Aghajanian

1 Question 1

The results outputted by the program to interpolate the following x from the given data are:

- (i) $y(x = 0.4) = 0.804498$
- (ii) $y(x = -0.128) = 0.967812$
- (iii) $y(x = -2.0) = 0.0290775$
- (iv) $y(x = 3.2) = 0.000239416$

2 Question2

For natural splines, the second derivatives at the boundaries, denoted y_1'' and y_n'' , are set to zero. This means that they are known and do not need to be solved for, reducing the $N \times N$ matrix to an $N - 2 \times N - 2$ matrix. However, if we wish to retain the dimensions of the matrix such that they match the square of the number of known data points we have to set the boundary equations such that $F_1 = F_n = c_1 = a_n = 0$ and $b_1 \neq 0$, $b_n \neq 0$ (these can be set to any non-zero constant):

$$b_1 y_1'' = 0, \quad b_n y_n'' = 0 \quad (1)$$

which satisfy the natural boundary conditions.

$$\frac{x_3 - x_1}{3} y_2'' + \frac{x_3 - x_2}{6} y_3'' = \frac{y_3 - y_2}{x_3 - x_2} - \frac{y_2 - y_1}{x_2 - x_1} \quad (2)$$

$$\frac{x_{n-1} - x_{n-2}}{6} y_{n-2}'' + \frac{x_n - x_{n-1}}{3} y_{n-1}'' = \frac{y_n - y_{n-1}}{x_n - x_{n-1}} - \frac{y_{n-1} - y_{n-2}}{x_{n-1} - x_{n-2}} \quad (3)$$

(ii) In the case of estimating the first derivatives at the boundary, we can no longer impose that $y_1'' = y_n'' = 0$ and the system of linear equation includes 2 more unknowns. The 2 extra equations needed for this are provided by specification of the first derivatives. The fundamental master equation is obtain by imposing the continuity of the first derivative at the interface of adjacent segments between known points. At the x_1 and x_n points, this continuity constraint can be written as (for $j = 1$):

$$\lim_{x \rightarrow x_1^+} \frac{dy}{dx} = \lim_{x \rightarrow x_1^-} \frac{dy}{dx} \quad (4)$$

Which is evaluated as:

$$\frac{y_2 - y_1}{x_2 - x_1} + \frac{1}{6}(x_2 - x_1)((3B_1^2 - 1)y_2'' - (3A_1^2 - 1)y_1'') = \delta_1; \quad (5)$$

As $x \rightarrow x_1^+$, the coefficients are $B_1 \rightarrow 0$ and $A_1 \rightarrow 1$, meaning that:

$$\frac{y_2 - y_1}{x_2 - x_1} + \frac{1}{6}(x_2 - x_1)((-1)y_2'' - 2y_1'') = \delta_1 \quad (6)$$

$$\frac{y_2 - y_1}{x_2 - x_1} - \delta_1 = \frac{(x_2 - x_1)}{3} y_1'' + \frac{(x_2 - x_1)}{6} y_2'' \quad (7)$$

as required. Similar treatment can be given to obtaining a second equation to relate the second derivative of y given the first derivative at the boundary ($j = n$) as δ_n :

$$\lim_{x \rightarrow x_n^+} \frac{dy}{dx} = \lim_{x \rightarrow x_n^-} \frac{dy}{dx} \quad (8)$$

$$\delta_n = \frac{y_n - y_{n-1}}{x_n - x_{n-1}} + \frac{1}{6}(x_n - x_{n-1}) ((3B_{n-1}^2 - 1)y_n'' - (3A_1^{n-1} - 1)y_{n-1}'') \quad (9)$$

where the coefficient A and B are labelled by the j th interval, of which there are $n - 1$ between the n points. Since $\lim_{x \rightarrow x_n^-} A_{n-1} = 0$ $\lim_{x \rightarrow x_n^-} B_{n-1} = 1$ this becomes:

$$\delta_n = \frac{y_n - y_{n-1}}{x_n - x_{n-1}} + \frac{1}{6}(x_n - x_{n-1}) (2y_n'' - (-1)y_{n-1}'') \quad (10)$$

$$\delta_n - \frac{y_n - y_{n-1}}{x_n - x_{n-1}} = \frac{(x_n - x_{n-1})}{6} y_{n-1}'' + \frac{(x_n - x_{n-1})}{3} y_n'' \quad (11)$$

$$\frac{y_n - y_{n-1}}{x_n - x_{n-1}} - \delta_n = \frac{(x_{n-1} - x_n)}{6} y_{n-1}'' + \frac{(x_{n-1} - x_n)}{3} y_n'' \quad (12)$$

3 Question 3

For this program, the routine **tridag** from Numerical Recipes was studied and used for diagonalising the matrix necessary to obtain the second derivatives. In the case of natural boundary conditions, the elements b_1 (the first element of b) and b_n (last element of b) set to 1 and $a_2 = c_1 = a_n = c_{n-1} = 0$. This means an $N \times N$ matrix is still solved but with 2 of the decoupled equations corresponding to $y_1'' = y_n'' = 0$ included as part of the matrix. For first-derivative specified boundary conditions, the values of the first derivative are used in the above equations to include in the matrix, solving the $N \times N$ problem.

4 Question 4

The results from the cubic spline interpolation for both natural splines and zero-first-derivative boundary conditions for the given data are as follow:

- (i) Natural spline: $y(x = 0.4) = 0.85085$, Zero first derivative: $y(x = 0.4) = 0.85085$
- (ii) Natural spline: $y(x = -0.128) = 0.982731$, Zero first derivative: $y(x = -0.128) = 0.982751$
- (iii) Natural spline: $y(x = -2) = 0.0206424$, Zero first derivative: $y(x = -2) = 0.0145888$
- (iv) Natural spline: $y(x = 3.2) = 0.00476971$, Zero first derivative: $y(x = 3.2) = 0.00308786$

5 Question 5

See Figure 1.

6 Question 6

The cubic spline interpolation appears to give a better representation of the function, being smooth in both value and first derivative. The reason why is would be a better choice over the linear interpolation is that the gradient smoothly varies, and is not piecewise, meaning it is more likely to be a true representation of a function in nature.

If this function were known to be a probability function, then it would be better to use the linear interpolation scheme. This is because for cubic spline interpolation, the rapid change in gradient towards the right hand side of the graph as the data smooths to zero, results in "ringing" to occur, in which the interpolated function overshoots and as a result there is a portion of the graph which has negative y values. A probability function cannot have negative values as this is unphysical, and hence the linear interpolation, albeit less accurate, will be a better estimate of the function to use, as we can tabulate it for all x within the range of the data without running into negative probabilities.




Figure 1: Linear interpolation, and cubic interpolation with both zero first derivative and natural boundary conditions implemented.

7 Question 7