```
Advent of Code [About] [Events] [Shop] [Settings] [Log Out] Martin Graham 26* /*2019*/ [Calendar] [AoC++] [Sponsors] [Leaderboard] [Stats]
```

--- Day 14: Space Stoichiometry ---

As you approach the rings of Saturn, your ship's low fuel indicator turns on. There isn't any fuel here, but the rings have plenty of raw material. Perhaps your ship's Inter-Stellar Refinery Union brand nanofactory can turn these raw materials into fuel.

You ask the nanofactory to produce a list of the reactions it can perform that are relevant to this process (your puzzle input). Every reaction turns some quantities of specific input chemicals into some quantity of an output chemical. Almost every chemical is produced by exactly one reaction; the only exception, $\overline{\text{ORE}}$, is the raw material input to the entire process and is not produced by a reaction.

You just need to know how much ORE you'll need to collect before you can produce one unit of FUEL.

Each reaction gives specific quantities for its inputs and output; reactions cannot be partially run, so only whole integer multiples of these quantities can be used. (It's okay to have leftover chemicals when you're done, though.) For example, the reaction $\boxed{1\ A,\ 2\ B,\ 3\ C} \Rightarrow 2\ D$ means that exactly 2 units of chemical $\boxed{0}$ can be produced by consuming exactly $1\ \boxed{A},\ 2\ \boxed{B}$ and $3\ \boxed{C}$. You can run the full reaction as many times as necessary; for example, you could produce $10\ \boxed{0}$ by consuming $5\ \boxed{A},\ 10\ \boxed{B},\$ and $15\ \boxed{C}$.

Suppose your nanofactory produces the following list of reactions:

```
10 ORE => 10 A

1 ORE => 1 B

7 A, 1 B => 1 C

7 A, 1 C => 1 D

7 A, 1 D => 1 E

7 A, 1 E => 1 FUEL
```

The first two reactions use only \overline{ORE} as inputs; they indicate that you can produce as much of chemical \overline{A} as you want (in increments of 10 units, each 10 costing 10 \overline{ORE}) and as much of chemical \overline{B} as you want (each costing 1 \overline{ORE}). To produce 1 \overline{FUEL} , a total of 31 \overline{ORE} is required: 1 \overline{ORE} to produce 1 \overline{B} , then 30 more \overline{ORE} to produce the 7 + 7 + 7 + 7 = 28 \overline{A} (with 2 extra \overline{A} wasted) required in the reactions to convert the \overline{B} into \overline{C} , \overline{C} into \overline{D} , \overline{D} into \overline{E} , and finally \overline{E} into \overline{FUEL} . (30 \overline{A} is produced because its reaction requires that it is created in increments of 10.)

Or, suppose you have the following list of reactions:

```
9 ORE => 2 A
8 ORE => 3 B
7 ORE => 5 C
3 A, 4 B => 1 AB
5 B, 7 C => 1 BC
4 C, 1 A => 1 CA
2 AB, 3 BC, 4 CA => 1 FUEL
```

The above list of reactions requires 165 ORE to produce 1 FUEL:

```
- Consume 45 ORE to produce 10 A.
- Consume 64 ORE to produce 24 B.
- Consume 56 ORE to produce 40 C.
- Consume 6 A, 8 B to produce 2 AB.
- Consume 15 B, 21 C to produce 3 BC.
- Consume 16 C, 4 A to produce 4 CA.
- Consume 2 AB, 3 BC, 4 CA to produce 1 FUEL.
```

Our sponsors help make Advent of Code possible:

Reaktor -Tracking Santa from space since 2018. Always looking for more talented Santa's little helpers. Here are some larger examples:

- 13312 ORE for 1 FUEL:

```
157 ORE => 5 NZVS

165 ORE => 6 DCFZ

44 XJWVT, 5 KHKGT, 1 QDVJ, 29 NZVS, 9 GPVTF, 48 HKGWZ => 1 FUEL

12 HKGWZ, 1 GPVTF, 8 PSHF => 9 QDVJ

179 ORE => 7 PSHF

177 ORE => 5 HKGWZ

7 DCFZ, 7 PSHF => 2 XJWVT

165 ORE => 2 GPVTF

3 DCFZ, 7 NZVS, 5 HKGWZ, 10 PSHF => 8 KHKGT
```

- 180697 ORE for 1 FUEL:

```
2 VPVL, 7 FWMGM, 2 CXFTF, 11 MNCFX => 1 STKFG
17 NVRVD, 3 JNWZP => 8 VPVL
53 STKFG, 6 MNCFX, 46 VJHF, 81 HVMC, 68 CXFTF, 25 GNMV => 1 FUEL
22 VJHF, 37 MNCFX => 5 FWMGM
139 ORE => 4 NVRVD
144 ORE => 7 JNWZP
5 MNCFX, 7 RFSQX, 2 FWMGM, 2 VPVL, 19 CXFTF => 3 HVMC
5 VJHF, 7 MNCFX, 9 VPVL, 37 CXFTF => 6 GNMV
145 ORE => 6 MNCFX
1 NVRVD => 8 CXFTF
1 VJHF, 6 MNCFX => 4 RFSQX
176 ORE => 6 VJHF
```

- 2210736 ORE for 1 FUEL:

```
171 ORE => 8 CNZTR
7 ZLQW, 3 BMBT, 9 XCVML, 26 XMNCP, 1 WPTQ, 2 MZWV, 1 RJRHP => 4 PLWSL
114 ORE => 4 BHXH
14 VRPVC => 6 BMBT
6 BHXH, 18 KTJDG, 12 WPTQ, 7 PLWSL, 31 FHTLT, 37 ZDVW => 1 FUEL
6 WPTO, 2 BMBT, 8 ZLOW, 18 KTJDG, 1 XMNCP, 6 MZWV, 1 RJRHP => 6 FHTLT
15 XDBXC, 2 LTCX, 1 VRPVC => 6 ZLQW
13 WPTQ, 10 LTCX, 3 RJRHP, 14 XMNCP, 2 MZWV, 1 ZLQW => 1 ZDVW
5 BMBT => 4 WPTQ
189 ORE => 9 KTJDG
1 MZWV, 17 XDBXC, 3 XCVML => 2 XMNCP
12 VRPVC, 27 CNZTR => 2 XDBXC
15 KTJDG, 12 BHXH => 5 XCVML
3 BHXH, 2 VRPVC => 7 MZWV
121 ORE => 7 VRPVC
7 XCVML => 6 RJRHP
5 BHXH, 4 VRPVC => 5 LTCX
```

Given the list of reactions in your puzzle input, what is the minimum amount of $\overline{\text{ORE}}$ required to produce exactly 1 $\overline{\text{FUEL}}$?

To I	begir	n, ge	t yo	ur pi	uzzle	inp	ut.
Ansı	wer:						[Submit]
You	can	also	ΓSh	arel	this	puz.	zle.