

Spatially optimal targeting of interventions to reduce air pollution

Martin Kosík

Motivation

- Air pollution (PM2.5, PM10) shown to have adverse effects on health (Deryugina et al., 2019), productivity (Chang et al., 2016), and academic achievement (Gilraine and Zheng, 2022)
- Crop residue burning is an important contributor air pollution in Indian cities during the late fall (Liu et al., 2018)
- Various interventions proposed to reduce residue burning such as conditional payments to farmers
- **This paper:** Given limited resourced, which places should be targeted for interventions to reduce air pollution?

- **Goal:** Target the interventions into places where the greatest impact can be achieved
- Modeling of two main aspects:
 1. Harm
 - On average, how much harm would additional emissions from a given location cause?
 - Depends on the weather patterns (wind direction, strength, etc.) and spatial distribution of the population
 - I will use an air pollution transport model (HYSPLIT) to estimate the overall impact
 2. Costs
 - How much we would have to spend to reduce the pollution in a given location

Modeling air pollution transport


- HYSPLIT dispersion model
 - Hybrid Single-Particle Lagrangian Integrated Trajectory model
 - One of the most extensively used atmospheric transport and dispersion models in the atmospheric sciences
 - Applications include tracking and forecasting the release of wildfire smoke, wind-blown dust, volcanic ash, and crop residue burning (Stein et al., 2015)
- Main output of interest
 - Source-receptor matrix: SRM_{ij}
 - Fraction of emissions from source i that are transported into j
 - Average over 8 days after release

test_conc

Concentration (/m3) averaged between 0 m and 100 m

Integrated from 1200 01 Oct to 0000 02 Oct 16 (UTC)

TEST Release started at 1200 01 Oct 16 (UTC)




NGM METEOROLOGICAL DATA

test_conc

Concentration (/m3) averaged between 0 m and 100 m

Integrated from 0000 02 Oct to 1200 02 Oct 16 (UTC)

TEST Release started at 1200 01 Oct 16 (UTC)




test_conc

Concentration (/m3) averaged between 0 m and 100 m

Integrated from 1200 02 Oct to 0000 03 Oct 16 (UTC)

TEST Release started at 1200 01 Oct 16 (UTC)




NGM METEOROLOGICAL DATA

test_conc

Concentration (/m3) averaged between 0 m and 100 m

Integrated from 0000 03 Oct to 1200 03 Oct 16 (UTC)

TEST Release started at 1200 01 Oct 16 (UTC)




test_conc

Concentration (/m3) averaged between 0 m and 100 m

Integrated from 1200 03 Oct to 0000 04 Oct 16 (UTC)

TEST Release started at 1200 01 Oct 16 (UTC)



NGM METEOROLOGICAL DATA

Measuring the impact - definitions

- SRM_{ij} ... fraction of emissions from source i that are transported into j
- E_i ... total air pollution emitted from location i
- $P_j = \sum_i SRM_{ij} E_i$... total air pollution in j
- $L_j = f(P_j)$... loss (harm) to a single person from being exposed to air pollution in j
- N_j ... total population in j
- $TL = \sum_j L_j \cdot N_j$... total population-weighted loss caused by air pollution across all locations

Measuring the impact

- The impact of small change emissions from i on total loss

$$\frac{\partial TL}{\partial E_i} = \sum_j \frac{\partial L_j}{\partial E_i} N_j = \sum_j \frac{\partial f(P_j)}{\partial P_j} \frac{\partial P_j}{\partial E_i} N_j = \sum_j \frac{\partial f(P_j)}{\partial P_j} SRM_{ij} N_j$$

- if $f(P_j) = a + b \cdot P_j$, this simplifies to


$$\frac{\partial TL}{\partial E_i} = b \sum_j SRM_{ij} N_j := b \cdot \alpha_j$$

- Clearly, it is optimal to target locations with the highest $\frac{\partial TL}{\partial E_i}$ since that will lead to greatest reductions in loss
- In case of linear $f(P_j)$, this means locations with highest α_j
 - Some evidence to support linear effect of PM2.5 concentrations on infant mortality (Heft-Neal et al., 2020)


Preliminary results

- I focus on North-west of India where the air pollution is very severe and crop residue burning is common
- I run simulations based on weather data for the beginning of October for 10 different years
- Regular grid of 121 source location
 - α_j computed for each location separately, then interpolated across across them on a finer grid
 - Further, I restricted the locations to those with likely winter cropping (per Jain et al., 2016 data) and with the total fire radiating power of at least 5 (using VIIRS fire data)


Winter cropped area




Total fire radiation power - October 2019




α_j in space




α_j - interpolated



α_j - interpolated - only crop residue burning locations



α_j - histogram



Conclusion

- The very preliminary results suggest that might be larger gains from optimal spatial targeting of interventions to reduce air pollution
- This is especially case for small-scale projects
 - Assuming linear loss and spatially uniform costs, targeting 10% crop-residue burning locations with the highest α_j results in around 55% greater reduction in loss compared to targeting 10% location with mean α_j
- More work to do on the cost side




References



Chang, Tom, Joshua Graff Zivin, Tal Gross, and Matthew Neidell (2016). *The Effect of Pollution on Worker Productivity: Evidence from Call-Center Workers in China*. Working Paper 22328. Series: Working Paper Series. National Bureau of Economic Research. DOI: [10.3386/w22328](https://doi.org/10.3386/w22328).



Deryugina, Tatyana, Garth Heutel, Nolan H. Miller, David Molitor, and Julian Reif (2019). “The Mortality and Medical Costs of Air Pollution: Evidence from Changes in Wind Direction”. en. In: *American Economic Review* 109.12, pp. 4178–4219. DOI: [10.1257/aer.20180279](https://doi.org/10.1257/aer.20180279).

-  Gilraine, Michael and Angela Zheng (2022). *Air Pollution and Student Performance in the U.S.* Working Paper 30061. Series: Working Paper Series. National Bureau of Economic Research. DOI: [10.3386/w30061](https://doi.org/10.3386/w30061).
-  Heft-Neal, Sam, Jennifer Burney, Eran Bendavid, Kara K. Voss, and Marshall Burke (2020). “Dust pollution from the Sahara and African infant mortality”. en. In: *Nature Sustainability* 3.10, pp. 863–871. DOI: [10.1038/s41893-020-0562-1](https://doi.org/10.1038/s41893-020-0562-1).
-  Jain, M., P. Mondal, G.L. Galford, G. Fiske, and R.S. DeFries (2016). *India Annual Winter Cropped Area, 2001-2016*. Type: dataset. DOI: [10.7927/H47D2S3W](https://doi.org/10.7927/H47D2S3W).



Liu, Tianjia, Miriam E. Marlier, Ruth S. DeFries, Daniel M. Westervelt, Karen R. Xia, Arlene M. Fiore, Loretta J. Mickley, Daniel H. Cusworth, and George Milly (2018). “Seasonal impact of regional outdoor biomass burning on air pollution in three Indian cities: Delhi, Bengaluru, and Pune”. en. In: *Atmospheric Environment* 172, pp. 83–92. DOI: [10.1016/j.atmosenv.2017.10.024](https://doi.org/10.1016/j.atmosenv.2017.10.024).



Stein, A. F., R. R. Draxler, G. D. Rolph, B. J. B. Stunder, M. D. Cohen, and F. Ngan (2015). “NOAA’s HYSPLIT Atmospheric Transport and Dispersion Modeling System”. EN. In: *Bulletin of the American Meteorological Society* 96.12. Publisher: American Meteorological Society Section: Bulletin of the American Meteorological Society, pp. 2059–2077. DOI: [10.1175/BAMS-D-14-00110.1](https://doi.org/10.1175/BAMS-D-14-00110.1).

Thank you for your attention.