All you need to know about relational hyper-event modeling

Linear RHEMs - Tutorial Section: Solution

Martina Boschi, Melania Lembo, Juergen Lerner

2026-02-10

Part 1: Les Misérables

Setup

```
# install.packages("mgcv")
# install.packages("RColorBrewer")
# install.packages("survival")
# install.packages("dplyr")
library(mgcv)
library(RColorBrewer)
library(survival)
library(dplyr)
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

1. Les Misérables: Data and Covariate Computation

This tutorial is inspired by the analysis conducted in (Lerner et al., 2025), where, for illustrative purposes, RHEM is applied to the network of actor co-appearances in *Les Misérables*, compiled by Donald Knuth (Knuth, 1993). The data comprises 80 actors (co-)appearing in one or several of 288 chapters. The (binary[1]) gender of the actors is known.

a) **Given the information above, interpret this data as a relational hyper-event network. Define the components of the hyperevent sequence, the vertex set, its cardinality, and the available attributes.**

- **80 actors** $|V| = 80$ (co-)appearing in one or several of **288 chapters** ($n = 288$);
- $e_i = (S_i, t_i)$ is a **hyperevent** (a chapter) involving a set of co-appearing actors S_i. In this case, t_i is simply a counter for the chapter;
- female(i) is a **binary gender attribute** for each actor.

b) **Open the file `jean_events.csv`. How are the events reported? Does it contain only event information?**

The file `jean_events.csv` contains three types of entries (which can be distinguished using the column `type`), namely `add.actor`, `is.female`, and `chapter`. Each entry involves only one actor at a time. A chapter is therefore represented by as many rows as the number of actors involved.

[1] only for mathematical purposes

c) **Open the configuration file `jean.config.small.txt` in `eventnet` and inspect the different windows.**

See the tutorial notebook for more information on how the data is treated in `eventnet`.

d) **Load the input data `jean_events_EVENTS.csv` located in `01-Data`. Is this a case-1-control dataset? If not, how do you think it can be constructed?**

In`eventnet` we computed the necessary statistics to support the inference techniques that will be explored. Upon inspecting the dataset, you may notice the presence of missing values (NA) in some columns.

In particular, the `TARGET` column contains only missing values, while the `SOURCE` column includes multiple entries. This indicates that we are dealing with **undirected relational hyper-events**, where the sender set is a subset of the vertices in a relational hypergraph V, and the receiver set is empty.

Additionally, the `EVENT_INTERVAL_ID` column does not hold interpretable values, as in this application we just know the order of events. The `IS_OBSERVED` column is used to distinguish between actual observed events and non-events (i.e., events that could have occurred but did not).

```
data_original <- read.csv("01-Data/jean_events_EVENTS.csv")
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

```
head(data_original)
```

```
##   IS_OBSERVED     SOURCE TARGET    TYPE EVENT_INTERVAL_ID EVENT INTEGER_TIME
## 1           1 |MY|NP|MB|     NA chapter             1.1.1   109          108
## 2           0 |CL|GR|MT|     NA                             111          110
## 3           0 |MA|BB|CV|     NA                             111          110
## 4           0 |PG|GT|CM|     NA                             111          110
## 5           0 |TH|MM|HL|     NA                             111          110
## 6           0 |TH|LL|SS|     NA                             111          110
##   TIME_POINT TIME_UNIT EVENT_INTERVAL num.actors individual.activity
## 1        109       108              2          3                   0
## 2        111       110              2          3                   0
## 3        111       110              2          3                   0
## 4        111       110              2          3                   0
## 5        111       110              2          3                   0
## 6        111       110              2          3                   0
##   dyadic.activity closure female diff.female
## 1               0       0      1           2
## 2               0       0      2           2
## 3               0       0      0           0
## 4               0       0      0           0
## 5               0       0      1           2
## 6               0       0      2           2
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

In this application, the `num.actors` column is also unnecessary. This is because non-events have been sampled to match the cardinality of the observed events, meaning that the effect of event size cannot be meaningfully estimated.

To simplify visualization, we proceed by removing all unnecessary columns from the dataset.

```
data <- data_original[,setdiff(colnames(data_original),
                               c("TARGET", "TYPE", "EVENT_INTERVAL_ID", "EVENT",
                                 "INTEGER_TIME", "TIME_POINT",
```

```r
                                  "TIME_UNIT", "num.actors"))]
rm(data_original)
head(data)
```

```
##   IS_OBSERVED     SOURCE EVENT_INTERVAL individual.activity dyadic.activity
## 1           1 |MY|NP|MB|              2                   0               0
## 2           0 |CL|GR|MT|              2                   0               0
## 3           0 |MA|BB|CV|              2                   0               0
## 4           0 |PG|GT|CM|              2                   0               0
## 5           0 |TH|MM|HL|              2                   0               0
## 6           0 |TH|LL|SS|              2                   0               0
##   closure female diff.female
## 1       0      1           2
## 2       0      2           2
## 3       0      0           0
## 4       0      0           0
## 5       0      1           2
## 6       0      2           2
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

The object `data` is not a case-1-control dataset. To construct it, we first create two separate objects for the datasets containing events and non-events, respectively. We then merge these datasets using a set of variables that identify which event each non-event refers to. Among the non-events satisfying this condition, we sample only one.

```r
data_ev <- data %>% filter(IS_OBSERVED == 1)
data_nv <- data %>% filter(IS_OBSERVED == 0)
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

```r
# Add a new column to label each row in 'data_ev' and 'data_nv'
# as event data and non-event data respectively
data_ev_tagged <- data_ev %>%
  mutate(.row_type = "ev")
data_nv_tagged <- data_nv %>%
  mutate(.row_type = "nv")
head(data_ev_tagged)
```

```
##   IS_OBSERVED              SOURCE EVENT_INTERVAL individual.activity
## 1           1          |MY|NP|MB|              2                   0
## 2           1          |MY|ME|MB|              3                   2
## 3           1                |MY|              4                   2
## 4           1 |MY|ME|CL|GE|MC|MB|              5                   6
## 5           1          |MY|MB|ME|              6                   9
## 6           1             |ME|MY|              7                   8
##   dyadic.activity closure female diff.female .row_type
## 1               0       0      1           2        ev
## 2               1       2      2           2        ev
## 3               0       0      0           0        ev
## 4               4       8      3           9        ev
## 5               7      32      2           2        ev
## 6               3      12      1           1        ev
```

```r
# head(data_nv_tagged)
rm(data_ev)
rm(data_nv)
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

```r
set.seed(1234)
merge_id_cols <- c("EVENT_INTERVAL")

# For each group defined by the merge_id_cols, take one random non-event
data_nv_1_sampled <- data_nv_tagged %>%
  group_by(across(all_of(merge_id_cols))) %>%
  slice_sample(n = 1) %>%
  ungroup()

# Perform a left join:
# for each row in data_ev, attach the corresponding non-event row (if available)
# based on matching values in the merge_id_cols
# The suffixes "_ev" and "_nv" will be added to columns from data_ev and data_nv
dat_gam_1 <- data_ev_tagged %>%
  left_join(data_nv_1_sampled,
            by = merge_id_cols, suffix = c("_ev", "_nv"))
rm(data_nv_1_sampled)
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

2. Model fitting: conditional and degenerate logistic regression

a) **Based on the data structures you were given and those you constructed, which inference techniques can you apply?**

OPTION 1

Since we do not have access to timing information for the events and we have a case-control sampling with 20 non-events, we use the `clogit` function in R to fit our model (conditional logistic regression). When applied in this context, `clogit` internally calls the `coxph` routine.

```r
clogit_fit <- clogit(IS_OBSERVED ~
                        + diff.female
                        + female
                        + individual.activity
                        + dyadic.activity
                        + strata(EVENT_INTERVAL)
                        , data = data)
summary(clogit_fit)
```

```
## Call:
## coxph(formula = Surv(rep(1, 6048L), IS_OBSERVED) ~ +diff.female +
##     female + individual.activity + dyadic.activity + strata(EVENT_INTERVAL),
##     data = data, method = "exact")
##
##   n= 6048, number of events= 288
##
##                          coef exp(coef)  se(coef)      z Pr(>|z|)
```

```
## diff.female         -0.290310  0.748032  0.117345 -2.474   0.0134 *
## female              -0.293382  0.745737  0.144822 -2.026   0.0428 *
## individual.activity  0.042973  1.043909  0.003918 10.967   <2e-16 ***
## dyadic.activity      0.460395  1.584701  0.049289  9.341   <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
##                     exp(coef) exp(-coef) lower .95 upper .95
## diff.female            0.7480     1.3368    0.5943    0.9415
## female                 0.7457     1.3410    0.5615    0.9905
## individual.activity    1.0439     0.9579    1.0359    1.0520
## dyadic.activity        1.5847     0.6310    1.4388    1.7454
##
## Concordance= 0.896  (se = 0.013 )
## Likelihood ratio test= 819.9  on 4 df,   p=<2e-16
## Wald test            = 245.8  on 4 df,   p=<2e-16
## Score (logrank) test = 1418  on 4 df,   p=<2e-16
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

We find a positive effect of individual, dyadic. This suggests that prior (co-)presence in previous chapters - whether alone or in pairs - increases the rate of participating in events. A negative effect for diff.female suggests a positive effect for gender homophily, favouring co-appearence with other actors of the same gender.

OPTION 2

Using the case-1-control dataset, it is also possible to fit a degenerate logistic regression without an intercept, where the covariates are defined as the differences between the covariates for the events and those for the non-events, and the response variable is constant and equal to 1.

```r
# covariates defined as difference between
# covariate values for event and corresponding non-event
dat_gam_1$female <-
  dat_gam_1$female_ev - dat_gam_1$female_nv
dat_gam_1$diff_female <-
  dat_gam_1$diff.female_ev - dat_gam_1$diff.female_nv
dat_gam_1$individual_activity <-
  dat_gam_1$individual.activity_ev - dat_gam_1$individual.activity_nv
dat_gam_1$dyadic_activity <-
  dat_gam_1$dyadic.activity_ev - dat_gam_1$dyadic.activity_nv
dat_gam_1$closure <-
  dat_gam_1$closure_ev - dat_gam_1$closure_nv

# constant response equal to 1
dat_gam_1$y <- 1
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

```r
gam_fit <- glm(y ~
                + diff_female
                + female
                + individual_activity
                + dyadic_activity
                - 1 # no intercept
               , data = dat_gam_1,
```

```r
                family="binomial")
```

```
## Warning: glm.fit: fitted probabilities numerically 0 or 1 occurred
```

```r
summary(gam_fit)
```

```
##
## Call:
## glm(formula = y ~ +diff_female + female + individual_activity +
##     dyadic_activity - 1, family = "binomial", data = dat_gam_1)
##
## Deviance Residuals:
##     Min       1Q   Median       3Q      Max
## 0.00000  0.00002  0.05696  0.69837  2.94123
##
## Coefficients:
##                     Estimate Std. Error z value Pr(>|z|)
## diff_female         -0.67999    0.37963  -1.791 0.073260 .
## female              -0.71924    0.31532  -2.281 0.022550 *
## individual_activity  0.07649    0.01704   4.490 7.13e-06 ***
## dyadic_activity      1.81184    0.54181   3.344 0.000826 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## (Dispersion parameter for binomial family taken to be 1)
##
##     Null deviance: 399.25  on 288  degrees of freedom
## Residual deviance: 121.75  on 284  degrees of freedom
## AIC: 129.75
##
## Number of Fisher Scoring iterations: 11
```

/Users/martina.boschi/Dropbox/Martina/PhD/12-REM-tutorial/02-All-you-need-to-know-RHEM/01-Les-Miserables/01-AYNTK-RHEM-Les-Miserables.R

As before, we find a positive effect of individual, dyadic and a negative main effect for female gender and a positive effect for gender homophily.