

# Shallow water model

Martin Claus, Willi Rath

February 15, 2011

## 1 Equations

### 1.1 Basic Equations

In Cartesian coordinates, the system is described by

$$u_t - fv = -g\eta_x + F_u \quad (1)$$

$$v_t + fu = -g\eta_y + F_v \quad (2)$$

$$\eta_t + (Hu)_x + (Hv)_y = F_\eta \quad (3)$$

Spherical coordinates with  $a$  being the radius of the earth,  $\lambda$  the longitude, and  $\vartheta$  the latitude:

$$u_t - fv = -g \frac{1}{a \cos \vartheta} \frac{\partial \eta}{\partial \lambda} + F_u \quad (4)$$

$$v_t + fu = -g \frac{1}{a} \frac{\partial \eta}{\partial \vartheta} + F_v \quad (5)$$

$$\eta_t + \frac{1}{a \cos \vartheta} \frac{\partial Hu}{\partial \lambda} + \frac{1}{a \cos \vartheta} \frac{\partial \cos \vartheta H v}{\partial \vartheta} = F_\eta \quad (6)$$

### 1.2 Forcing and Damping

The forcing and damping terms  $F_u$  in (1) and  $F_v$  in (2) represent one or the sum of some of the following terms.

#### 1.2.1 Bottom Friction Term

The model can use both linear and quadratic bottom friction. The linear friction term is described by

$$F = -\frac{r}{H}(u, v) \quad (7)$$

Discretized this leads to

$$F_u|_u^{i,j} = -\frac{r}{H|_u^{i,j}} u|_u^{ijl} \quad (8)$$

in the u-equation and

$$F_v|_v^{i,j} = -\frac{r}{H|_v^{i,j}} v|_v^{ijl} \quad (9)$$

in the v-equation, where r is the linear bottom friction coefficient. The quadratic friction term is described by

$$F = -\frac{k}{H} \sqrt{u^2 + v^2} (u, v) \quad (10)$$

Discretisation leads to

$$F_u|_u^{i,j} = -\frac{k}{H|_u^{ij}} \sqrt{(u|_u^{ijl})^2 + \left( \frac{v|_v^{i,j+1,l} + v|_v^{ijl} + v|_v^{i-1,j+1,l} + v|_v^{i-1,j,l}}{4} \right)^2} u|_u^{ijl} \quad (11)$$

in the u-equation and

$$F_v|_v^{i,j} = -\frac{k}{H|_v^{ij}} \sqrt{\left( \frac{u|_u^{i+1,j,l+1} + u|_u^{i+1,j-1,l+1} + u|_u^{i,j,l+1} + u|_u^{i,j-1,l+1}}{4} \right)^2 + (v|_v^{ijl})^2} v|_v^{ijl} \quad (12)$$

in the v-equation, where k is the empirical quadratic bottom friction coefficient.

### 1.2.2 Lateral Mixing of Momentum

The lateral mixing of momentum is described by

$$F_u = A_h f^\lambda \quad (13)$$

in the u-equation and

$$F_v = A_h f^\vartheta \quad (14)$$

in the v-equation, where  $A_h$  is the horizontal eddy viscosity. Following Bryan (1969) the lateral mixing of momentum is parameterized in spherical coordinates with consideration of conservation of momentum by

$$f^\lambda = \frac{1}{a^2 \cos^2 \vartheta H} \frac{\partial^2 u H}{\partial \lambda^2} + \frac{1}{a^2 \cos \vartheta H} \frac{\partial}{\partial \vartheta} \left( \cos \vartheta \frac{\partial u H}{\partial \vartheta} \right) + \frac{1}{a^2 H} \left( (1 - \tan^2 \vartheta) u H - \frac{2 \tan \vartheta}{\cos \vartheta} \frac{\partial v H}{\partial \lambda} \right) \quad (15)$$

$$f^\vartheta = \frac{1}{a^2 \cos^2 \vartheta H} \frac{\partial^2 v H}{\partial \lambda^2} + \frac{1}{a^2 \cos \vartheta H} \frac{\partial}{\partial \vartheta} \left( \cos \vartheta \frac{\partial v H}{\partial \vartheta} \right) + \frac{1}{a^2 H} \left( (1 - \tan^2 \vartheta) v H - \frac{2 \tan \vartheta}{\cos \vartheta} \frac{\partial u H}{\partial \lambda} \right) \quad (16)$$

Discretisation leads to

$$\begin{aligned}
f^\lambda|_u^{i,j,l} = & \frac{1}{2\Delta\lambda a^2 \cos^2 \vartheta|_u^j H|_u^{i,j}} (u|_u^{i+1,j,l} H|_u^{i+1,j} - 2u|_u^{i,j,l} H|_u^{i,j} + u|_u^{i-1,j,l} H|_u^{i-1,j}) \\
& - \frac{\tan \vartheta|_u^j}{2\Delta\vartheta a^2 H|_u^{i,j}} (u|_u^{i,j+1,l} H|_u^{i,j+1} - u|_u^{i,j-1,l} H|_u^{i,j-1}) \\
& + \frac{1}{2\Delta\vartheta a^2 H|_u^{i,j}} (u|_u^{i,j+1,l} H|_u^{i,j+1} - 2u|_u^{i,j,l} H|_u^{i,j} + u|_u^{i,j-1,l} H|_u^{i,j-1}) \\
& + \frac{1 - \tan^2 \vartheta|_u^j}{a^2 H|_u^{i,j}} u|_u^{i,j,l} H|_u^{i,j} \\
& - \frac{\tan \vartheta|_u^j}{a^2 \cos \vartheta|_u^j \Delta\lambda H|_u^{i,j}} (v|_v^{i,j+1,l} H|_v^{i,j+1} + v|_v^{i,j,l} H|_v^{i,j}) \\
& + \frac{\tan \vartheta|_u^j}{a^2 \cos \vartheta|_u^j \Delta\lambda H|_u^{i,j}} (v|_v^{i-1,j+1,l} H|_v^{i-1,j+1} + v|_v^{i-1,j,l} H|_v^{i-1,j})
\end{aligned} \tag{17}$$

and

$$\begin{aligned}
f^\vartheta|_v^{i,j,l} = & \frac{1}{2\Delta\lambda a^2 \cos^2 \vartheta|_v^j H|_v^{i,j}} (v|_v^{i+1,j,l} H|_v^{i+1,j} - 2v|_v^{i,j,l} H|_v^{i,j} + v|_v^{i-1,j,l} H|_v^{i-1,j}) \\
& - \frac{\tan \vartheta|_v^j}{2\Delta\vartheta a^2 H|_v^{i,j}} (v|_v^{i,j+1,l} H|_v^{i,j+1} - v|_v^{i,j-1,l} H|_v^{i,j-1}) \\
& + \frac{1}{2\Delta\vartheta a^2 H|_v^{i,j}} (v|_v^{i,j+1,l} H|_v^{i,j+1} - 2v|_v^{i,j,l} H|_v^{i,j} + v|_v^{i,j-1,l} H|_v^{i,j-1}) \\
& + \frac{1 - \tan^2 \vartheta|_v^j}{a^2 H|_v^{i,j}} v|_v^{i,j,l} H|_v^{i,j} \\
& - \frac{\tan \vartheta|_v^j}{a^2 \cos \vartheta|_v^j \Delta\lambda H|_v^{i,j}} (u|_u^{i+1,j,l} H|_u^{i+1,j} + u|_u^{i+1,j-1,l} H|_u^{i+1,j-1}) \\
& + \frac{\tan \vartheta|_v^j}{a^2 \cos \vartheta|_v^j \Delta\lambda H|_v^{i,j}} (u|_u^{i,j,l} H|_u^{i,j} + u|_u^{i,j-1,l} H|_u^{i,j-1})
\end{aligned} \tag{18}$$

Note that  $u|_u^l$  is used in the v-equation instead of  $u|_u^{l+1}$ .

### 1.2.3 Surface Stress Forcing

Wind stress  $\tau_s$  can be used to force the model. The forcing term in the u- and v-equation can be written as

$$F_u = \frac{\tau_s^\lambda}{\rho_0 H} \tag{19}$$

$$F_v = \frac{\tau_s^\vartheta}{\rho_0 H} \tag{20}$$

where  $\tau_s^\lambda$  is the zonal and  $\tau_s^\vartheta$  the meridional component of the wind stress. Since both components are provided on different grids, the zonal on the u-grid, the

meridional on the v-grid, discretization yields<sup>1</sup>

$$F_u|_u^{i,j} = \frac{\tau_s^\lambda|_u^{i,j}}{\rho_0 H|_u^{i,j}} \quad (21)$$

$$F_v|_v^{i,j} = \frac{\tau_s^\vartheta|_v^{i,j}}{\rho_0 H|_v^{i,j}} \quad (22)$$

### 1.2.4 Reynolds Stress Forcing

Reynolds stresses derived from observations can be used to force the model. These stresses are known at the surface and are supposed to decrease linearly with depth to zero. So the forcing terms in the u- and v-equation can be written as

$$\begin{aligned} F_u &= -\frac{1}{H} \int_{-H}^0 \frac{\partial}{\partial x} \left[ \overline{u'u'} \left( \frac{z+H}{H} \right) \right] dz - \frac{1}{H} \int_{-H}^0 \frac{\partial}{\partial y} \left[ \overline{u'v'} \left( \frac{z+H}{H} \right) \right] dz \\ &= -\frac{1}{2Ha \cos \vartheta} \frac{\partial}{\partial \lambda} (\overline{u'u'} H) - \frac{1}{2Ha \cos \vartheta} \frac{\partial}{\partial \vartheta} (\cos \vartheta \overline{u'v'} H) \end{aligned} \quad (23)$$

And analog

$$F_v = -\frac{1}{2Ha \cos \vartheta} \frac{\partial}{\partial \lambda} (\overline{u'v'} H) - \frac{1}{2Ha \cos \vartheta} \frac{\partial}{\partial \vartheta} (\cos \vartheta \overline{v'v'} H) \quad (24)$$

Discretization<sup>2</sup> leads to

$$\begin{aligned} F_u|_u^{i,j} &= -\frac{1}{2 H|_u^{i,j} a \cos \vartheta|_u^j} \left( \frac{\overline{u'u'}|_\eta^{i,j} H|_\eta^{i,j} - \overline{u'u'}|_\eta^{i-1,j} H|_\eta^{i-1,j}}{\Delta \lambda} \right. \\ &\quad \left. - \frac{\cos \vartheta|_H^{j+1} \overline{u'v'}|_H^{i,j+1} H|_H^{i,j+1} - \cos \vartheta|_H^j \overline{u'v'}|_H^{i,j} H|_H^{i,j}}{\Delta \vartheta} \right) \\ F_v|_v^{i,j} &= -\frac{1}{2 H|_v^{i,j} a \cos \vartheta|_v^j} \left( \frac{\cos \vartheta|_\eta^j \overline{v'v'}|_\eta^{i,j} H|_\eta^{i,j} - \cos \vartheta|_\eta^{j-1} \overline{v'v'}|_\eta^{i,j-1} H|_\eta^{i,j-1}}{\Delta \vartheta} \right. \\ &\quad \left. - \frac{\overline{u'v'}|_H^{i+1,j} H|_H^{i+1,j} - \overline{u'v'}|_H^{i,j} H|_H^{i,j}}{\Delta \lambda} \right) \end{aligned} \quad (25)$$

(26)

### 1.2.5 Newtonian Cooling

The  $F_\eta$  represents a Newtonian cooling term parameterized by

$$F_\eta = -\gamma \eta \quad (27)$$

Discretization leads to

$$F_\eta|_\eta^{i,j} = -\gamma \eta_\eta|_\eta^{i,j} \quad (28)$$

<sup>1</sup>There is an implicit two-point averaging of H to the u- and v-grid

<sup>2</sup>There is an implicit four point averaging from the H-grid to the  $\eta$ -grid since all stress terms and topography are provided on the H-grid

## 2 Grid

We want the grid to be as natural as possible (i.e., no averaging in the  $\eta$ -gradients and in the continuity equation):

$$\begin{array}{ccccc} H & v & H & v & H & v \\ u & \eta & u & \eta & u & \eta \\ H & v & H & v & H & v \end{array}$$

Figure 1: Grid arrangement. We use an Arakawa C-grid with the topography on the free grid points.

The fields will all carry integer indices which are to be interpreted according to

$$H|_H^{ij} = H(\Delta x \cdot i, \Delta y \cdot j) \quad (29)$$

$$\eta|_\eta^{ijl} = \eta(\Delta x \cdot (i + 1/2), \Delta y \cdot (j + 1/2), \Delta t \cdot l) \quad (30)$$

$$u|_u^{ijl} = u(\Delta x \cdot i, \Delta y \cdot (j + 1/2), \Delta t \cdot l) \quad (31)$$

$$v|_v^{ijl} = v(\Delta x \cdot (i + 1/2), \Delta y \cdot j, \Delta t \cdot l) \quad (32)$$

An expression like  $H|_u^{ij}$  means that the depth is interpolated to the  $u$ -grid by simple linear interpolation which is two-point averaging here. In the Coriolis term, we'll need four point averaging from the  $v$ -grid to the  $u$ -grid and from the  $u$ -grid to the  $v$ -grid. The  $x$ - and  $y$ -derivatives in the continuity equation will all need two point averaging from the  $H$ -grid to the  $u$ - and  $v$ -grid.

## 3 Boundary conditions

### 3.1 Coastlines / Land mask

We'll define land to be where  $H$  is zero. This implies that  $u$ - and  $v$ -grid points are over land if both the closest  $H$ -grid points are over land.  $\eta$ -grid points are over land, if all four surrounding  $H$ -grid points are over land. It will probably be computationally effective to define masks for all three dynamic variables.

### 3.2 Periodic boundary conditions

For a global model, we'll need to implement periodic boundary conditions in the zonal direction. In the meridional direction we might get away without as we won't be able to extend the model domain to the poles anyway (CFL will become a problem there).

## 4 Time stepping scheme

Use Heaps (1972) which basically amounts to<sup>3</sup>

$$\eta^{l+1} = \eta^l - \Delta t (H u^l)_x - \Delta t (H v^l)_y + \Delta t F_\eta \quad (33)$$

$$u^{l+1} = u^l + \Delta t f v^l - \Delta t g \eta_x^{l+1} + \Delta t F_u \quad (34)$$

$$v^{l+1} = v^l - \Delta t f u^{l+1} - \Delta t g \eta_y^{l+1} + \Delta t F_v \quad (35)$$

Spherical coordinates with full<sup>4</sup> glory:

$$\begin{aligned} \eta_{|_\eta}^{i,j,l+1} = & \eta_{|_\eta}^{ijl} + \Delta t F_{\eta|_\eta}^{i,j} \\ & - \frac{\Delta t}{a \cos \vartheta_{|_\eta}^j \Delta \lambda} \left( H_{|_u}^{i+1,j} u_{|_u}^{i+1,j,l} - H_{|_u}^{ij} u_{|_u}^{ijl} \right) \\ & - \frac{\Delta t}{a \cos \vartheta_{|_\eta}^j \Delta \vartheta} \left( H_{|_v}^{i,j+1} \cos \vartheta_{|_v}^{j+1} v_{|_v}^{i,j+1,l} - H_{|_v}^{ij} \cos \vartheta_{|_v}^j v_{|_v}^{ijl} \right) \end{aligned} \quad (36)$$

$$u_{|_u}^{i,j,l+1} = u_{|_u}^{ijl} + \Delta t f v_{|_u}^{ijl} - \frac{\Delta t g}{a \cos \vartheta_{|_u}^j \Delta \lambda} \left( \eta_{|_\eta}^{i,j,l+1} - \eta_{|_\eta}^{i-1,j,l+1} \right) + \Delta t F_{u|_u}^{i,j} \quad (37)$$

$$v_{|_v}^{i,j,l+1} = v_{|_v}^{ijl} - \Delta t f u_{|_v}^{i,j,l+1} - \frac{\Delta t g}{a \Delta \vartheta} \left( \eta_{|_\eta}^{i,j,l+1} - \eta_{|_\eta}^{i,j-1,l+1} \right) + \Delta t F_{v|_v}^{i,j} \quad (38)$$

In the model, we'll calculate  $H_{|_u}^{ij}$  and  $H_{|_v}^{ij}$  once in the beginning and then use it throughout the simulation. However, for defining which grid points are over land, it is still helpful to use the original  $H$ -grid.

## 5 Stability

Note that we will not consider optimization of roundoff errors at this point. We'll stick to SI-units and readable code as long as possible.

There are two criteria which may be of importance:  $|\Delta t \cdot f| < 1$  which comes from the Coriolis term, and the CFL criterion  $c \frac{\Delta t}{\Delta x} = \sqrt{gH} \frac{\Delta t}{\Delta x} < 1$ . The CFL criterion is by far the most dangerous. From the first criterion,  $\Delta t$  varies from infinity at the equator to half a day at the poles, while from CFL with high resolution, high latitudes and large  $H$  time steps of much less than 10 seconds can easily arise.

$$\Delta t < \frac{\Delta \lambda / \text{deg} \cdot 111 \text{ km} \cdot \cos \vartheta^{max}}{\sqrt{gH}} \quad (39)$$

For  $H = O(10 \text{ km})$ ,  $\Delta \lambda = 1/10^\circ$  and a model domain which extends up to  $89N$ , this results in as short a time steps as 0.6 seconds.

<sup>3</sup>Note that the superscrip of  $v$  in the second equation is mean to be an  $l$  while all the others are meant to be  $l + 1$ . In each line, you use the newest information that is available ...

<sup>4</sup>There still is an implicit two point averaging from the  $H$ -grid to the  $u$ - and  $v$ -grids as well as an implicit four point averaging in the Coriolis term.

## 6 Estimation of computational effort

In the end, we might want to simulate an almost global model (say  $\lambda = 0 \dots 360^\circ$ ,  $\vartheta = -89 \dots 89^\circ$ ) with a resolution of up to  $1/10^\circ$  for more than one year of model time. There are  $360 * 178 * 100 \approx 6.5 \cdot 10^6$  grid points and for each grid point we need  $O(20)$  multiplications and additions. The time step is  $\approx 0.6s$  and one year of model time has  $\approx 3.15 \cdot 10^7s$ . Altogether, we need  $N \approx 6.7 \cdot 10^{15}$  operations which on a machine with  $8GFLOPS$  can be done in  $8.5 \cdot 10^5s \approx 10d$ . For a smaller domain, which only extends to  $85^\circ$  in latitude, only about  $2d$  are necessary.

## 7 First test: Rectangular domain

Domain:

$$\begin{aligned}\lambda &= -20^\circ \dots 20^\circ \\ \vartheta &= -20^\circ \dots 20^\circ \\ \Delta\lambda &= \Delta\vartheta = 0.4^\circ \\ H &= 1000m = \text{const.} \\ \Rightarrow \Delta t &= 422s\end{aligned}\tag{40}$$

Initial conditions:

Adjustment should be done after  $O(20 \cdot 2 \cdot 111km / \sqrt{gH}) = O(4.5 \cdot 10^4s) = O(100)\Delta t$ .