

# Solving the FX cross-smiles problem - rate of convergence for Sinkhorn marginals, and the finite-option case

Martin Forde

19th Oct 2025

## Abstract

We adapt the arguments in [Guy20] to the problem of constructing a joint law for two FX rates  $X$  and  $Y$  consistent with observed European option prices on all three cross-rates. The usual Gibbs-type exponential ansatz for  $\mu$  leads to three coupled integral equations for the optimal European option portfolios  $u, v, w$ , and we prove  $O(\frac{1}{\sqrt{n}})$  convergence for the three marginals (in the total variation metric) for the average of the Sinkhorn iterates under natural assumptions on the target densities, assuming an admissible law exists (typically using  $\approx 50$  iterations in practice). The primary application here is risk-neutral pricing of e.g. Basket, Quanto, Best-of or other varieties of Rainbow options on  $X$  and  $Y$  so as to be consistent with all three smiles, which is useful if we can't or don't wish to estimate the true market model (i.e. the  $\mathbb{P}$  measure) so indifference pricing is infeasible or undesirable. We also consider the finite-option case via the usual duality relation between minimal entropy risk-neutral measures and utility maximization from chapter 3 in Föllmer&Schied[FS04]. The original constrained primal entropy minimization problem, and its dual (portfolio optimization) problem can also typically be solved directly (after appropriate discretization) using CVXPY with the MOSEK interior point solver even with a large number of options in the calibration set.<sup>1 2</sup>

## 1 Background

Regularized optimal transport methods have been a game changer for exotic option calibration problems in recent years, because it allows the inner inf in the minmax problem which arises from dualization to be computed explicitly (see e.g. pg 8 in [Guy20] or [GLOW22]). For regularized problems in the discrete-time setting, if we use an entropic penalty then can apply the Sinkhorn fixed point scheme (with provable convergence in certain standard cases, see e.g. [Nutz22]), for which the most notable practical application has been a discrete-time solution to the SPX-VIX calibration problem in [Guy20]. In particular, [Guy20] uses the Donsker-Varadhan/Gibbs variational formula  $\inf_{\mu \in \mathcal{P}} \{H(\mu|\bar{\mu}) - \mathbb{E}^\mu(X)\} = -\log \mathbb{E}^\mu(e^X)$  (see e.g. Eq 1.1 in Dupuis et al.[ACD15]) to simplify the supinf which arises from dualization.

In the continuous-time setting, we can also solve calibration problems using a Markov local and/or stochastic volatility models with finite tradeable European and/or VIX or barrier options using more general penalty terms, by numerically solving a HJB equation which emerges from dualization (see e.g. [GLOW22] and further discussion below), although rigorous duality results with viscosity solutions are more cumbersome to establish.

In principle, a bivariate [Carr09] Local Variance Gamma (LVG) model (see also [CN17]) can also give an exact fit to three cross-smiles, but in practice the implied local correlation function  $\rho_{loc}(x, y) := (\sigma_X(x)^2 + \sigma_Y(y)^2 - \sigma_Z(x/y)^2)/(2\sigma_X(x)\sigma_Y(y))$  that comes out from the three smiles typically falls outside  $[-1, 1]$  (similarly a two-maturity standard Carr LVG model can be fitted to a single SPX and VIX smile with the same maturity, but in practice the vol function for the 2nd epoch typically has a singularity, and a simpler (albeit somewhat unrealistic) model that achieves the same goal is described in Eq 6.3 in [Guy20]).

Guo et al. [GLW22],[GLOW22] (see also [HL19] and [Guy22]) show how to construct a generalized local/stochastic volatility model consistent with a finite number of European tradeable options at multiple maturities by minimizing a cost function over calibrated models which penalizes deviations from a standard reference model (e.g. Black-Scholes or Heston), and then re-casting the problem via dualization as an (unconstrained) minmax problem in terms of a non-linear HJB equation (so the cost function effectively regularizes the problem). If options at multiple maturities are used in the calibration set, the HJB equation unfortunately also includes Dirac source terms (but this can be avoided using a nested PDE), and this method is extended to include VIX options in section 3.3 in [GLOW22], by re-expressing  $V_t$  for the reference model in terms of  $\mathbb{E}(\int_t^T \sigma_s^2 ds | \mathcal{F}_t)$  (this analysis can be simplified using that  $VIX_t^2$  is just an affine function of  $V_t$  when the drift of  $V$  under the reference model has an affine drift). This approach is mathematically rich and exciting albeit numerically intensive since it requires numerically solving a non-linear HJB equation using e.g. fiddly implicit policy-iteration finite difference schemes, and then maximizing over the option weights vector. If path-dependent options are included in the calibration set we also have the issue that the prices may admit arbitrage.

<sup>1</sup>We thank Joseph Sullivan and Zhuoran Li for demonstrating this

<sup>2</sup>We thank David Hobson and Amir Dembo for sharing their insights. This document is an updated version of the older article <https://martinforde.github.io/FX-CrossSmiles.pdf>

[JLO23] construct the two-maturity Bass martingale  $M$ , by considering a standard 1d Brownian motion  $W$  with initial distribution  $\alpha_0$  with  $M_t = \mathbb{E}(F_1(W_T)|\mathcal{F}_t^W)$ . If  $\mu_t$  denotes the density of  $M_t$ , then from elementary arguments we can easily check that  $\mu_t = (R_{1-t} * F)_{\#}(R_t * \alpha_0)$  (where  $R_t$  is the Brownian density) and take  $\mu_0$  and  $\mu_1$  as given target laws, which leads to two coupled integral equations for  $\alpha_0$  and  $F_1$  when we set  $t = 0$  and  $t = 1$ . This naturally leads to an alternating Sinkhorn-type scheme for  $(\alpha_0, F_1)$  (see section 3.2 of [JLO23]) which they call the measure preserving martingale Sinkhorn (MPMS) algorithm. On page 6 with a few more lines they verify this formulation is equivalent to the fixed-point integral equation in [CL21] for the cdf  $G_{\alpha_0}$  of  $\alpha_0$  given by  $G_{\alpha_0} = G_{\mu_0} \circ (R_1 * F) = G_{\mu_0} \circ (R_1 * (G_{\mu_1}^{-1} \circ (R_1 * G_{\alpha_0})))$ , where  $G_{\nu}(\cdot)$  denotes the cdf of a probability measure  $\nu$ , and they give a short formal proof of the contraction property for this equation which was originally proved in Acciaio et al.[AMP25].

Recall that  $M$  in the preceding paragraph is also the solution to the martingale Benamou-Brenier problem (mBB):  $\inf_{M_0 \sim \mu, M_T \sim \nu, M_t = M_0 + \int_0^t \sigma_s dW_s} \mathbb{E}(\int_0^T (\sigma_t - 1)^2 dt)$  ( $T = 1$  above), hence the soubriquet “stretched Brownian motion”, or equivalently maximizes  $\mathbb{E}(\int_0^T |\sigma_t| dt)$  and  $\mathbb{E}(M_T W_T)$  over the same set. Using the HJB/dualization approach we can obtain the dual representation for this problem in Eq 2 in [JLO23] using the Fenchel-Moreau theorem (see [Loep25]) which can be re-written in a (possibly) more intuitive/familiar form as  $\sup_{\varphi} (-\int \varphi(y) \mu_T(dy) + \int u^{\varphi}(x, 0) \mu_0(dx))$  where  $u^{\varphi}$  satisfies the HJB equation  $u_t + \inf_{\sigma} (\frac{1}{2} \sigma^2 u_{xx} + \frac{1}{2} (\sigma^2 - 1)) = 0$  (if a classical solution exists) with  $u(x, T) = \varphi$ , which is the same as the usual form for [GLW22],[GLOW22] type problems with a single terminal target law  $\mu_T$  except now we integrate  $u^{\varphi}(x, 0)$  over  $\mu_0$  since this is a two-marginals problem. [BST23] give an alternate more explicit formulation for the dual here as  $\inf_{\psi \in L^1(\nu), \psi \text{ convex}} (\int \psi d\nu - \int (\psi^* * \gamma_1)^* d\mu)$  ( $\nu = \mu_T$  above) which we can prove just using basic properties of derivatives of Legendre transforms, but the [JLO23] formula is more easily adapted to the more general mBB(H) problem of minimizing  $\mathbb{E}(\int_0^T H(\sigma_t^2) dt)$ .

Note if we solve an entropically regularized M.O.T. problem of the form  $\inf_{\mu \in \mathcal{M}(\mu_1, \mu_2)} (H(\mu|\bar{\mu}) + \mathbb{E}^{\mu}(c(X, Y)))$  and similarly for  $(\mu_2, \mu_3)$ , with  $\mu_1, \mu_2, \mu_3$  in convex order, then concatenating the two conditional laws for  $Y|X$  yields a (Markov) martingale measure consistent with all three target laws (see [Bour25], which can then be used to price path-dependent options (of course one can also use the Carr Local Variance Gamma model to do this, cf. [Carr09],[CN17])). [Bour25],[BG24] also discusses use of Newton’s method to speed up Sinkhorn algorithms.

## 1.1 Outline of article

In this article, we adapt the Schrödinger bridge approach in [Guy20] to the problem of constructing a joint density for two FX rates  $X$  and  $Y$  consistent with observed European option prices on all three cross-rates. The cross-smile problem we consider in this article (as outlined in the abstract) falls under the general class of problems of the form  $\inf_{\mu \in C} H(\mu|\bar{\mu})$  for a convex set  $C$  which is closed under the total variation metric (see Lemma 2.2), for which there is a general existence/uniqueness result in Theorem 1.10 in [Nutz22].

Two of the Sinkhorn equations are of the usual fixed-point type but the final equation requires numerical root finding as for the fourth and fifth equations in the first order optimality conditions for the SPX-VIX problem in [Guy20]). This can also be solved as a pure optimal transport (OT) (linear programming) problem where we compute the max or min price of a non-tradeable option given tradeable European options on all three cross-rates (as indeed can Guyon’s SPX-VIX calibration problem<sup>3</sup>, see also [GMN17]. However, the Schrödinger bridge solution is more realistic since the optimal coupling  $\mu^*$  has a joint density unlike OT problems where the support of  $\mu^*$  is typically just the graph of  $n$ -functions. To cite a famous example, the left curtain coupling in [BJ16] is the solution  $\mathbb{P}^*$  to  $\sup_{\mathbb{P} \in \mathcal{M}(\mu, \nu)} \mathbb{E}^{\mathbb{P}}(c(X, Y))$  if  $c$  satisfies the *Spence-Mirless condition*  $c_{xyy} > 0$  and  $\mu \ll \text{Leb}$ , and  $Y = T_u(X)$  or  $T_d(X)$  if  $X$  lies in a certain range (i.e. a binomial model) and  $Y = X$  otherwise so  $\text{card}(\text{supp}(\mathbb{P}^*)) \leq 2$  (see also slide 14 in [dm18], and the heat maps on slides 16 and 38), with  $T_u(x) \geq x$  non-decreasing and  $T_d(x) \leq x$  non-increasing. Explicit derivations of the non-linear coupled ODEs for  $T_u(\cdot)$  and  $T_d(\cdot)$  are given on page 12 in [HLT16], and the associated hedge quantities on page 18 in [HLT16]). Variants of the left-curtain coupling are derived in the context of robust pricing/hedging for at-the-money forward straddle options using a Lagrangian argument in [HK15] ( $n = 3$ ), and [HN12] ( $n = 2$ ), and (as above) such problems can be efficiently approximated in MOSEK (one can also compute e.g. minimal price of an American puts with finite exercise dates given two marginals, see [Norg25],[HN19],[HN25] for theoretical results in this direction).

Returning to the current article, the usual Gibbs-type exponential ansatz for  $\mu(x, y)$  leads to three non-linear coupled integral equations for  $(u, v, w)$ , and we characterize the rate convergence for the three marginals (in the total variation metric) for the associated (quasi) fixed-point Sinkhorn scheme under certain assumptions on the target densities (which are computationally reasonable), assuming an admissible law exists with finite entropy. We also discuss more realistic choices for the reference measure e.g. using the Gaussian copula (which we numerically implement), and one can also adapt the Sinkhorn scheme for the forward-starter calibration problem in the usual setup with one asset and two maturities which leads to a fourth integral equation to enforce the martingale condition  $\mathbb{E}(\Delta(X)(Y - X))$  for all  $\Delta \in C_b$ ,<sup>4</sup> which also requires numerical root-finding. We later consider the finite-option case where the standard Sinkhorn algorithm is not directly applicable, using the duality between minimal entropy

<sup>3</sup>We thank Rowan Austin for demonstrating this in MOSEK using sparse flattened matrices to deal with the constraints.

<sup>4</sup>We thank Youpeng Chen and Max Chellev for demonstrating this

equivalent risk-neutral measures and the concave exponential utility maximization problem in chapter 3 in [FS04], and how to compute sensitivities of the value function in terms of the optimal portfolio weights which are the Lagrange multipliers. As mentioned in the abstract, we can typically use Python (or better Python with MOSEK or Clarabel) to solve the primal or dual problems directly (hence no need for Sinkhorn), and numerical results for the latter are given in Figure 3.

Finally, in section 3 (under mild assumptions) we show that the upper (resp. lower) bound for options on  $X/Y$  which pay  $(X - KY)^+$  in the original currency are attained by the lower (resp. upper) Fréchet-Hoeffding bound (see also Hobson et al. [HLW05], [HLW05b] for a more involved analysis with basket options where the weights are assumed to be non-negative or 1 unlike our example here).

## 2 Outline of problem

For major currencies like the Euro, Dollar and Pound, European options are very actively traded on all three cross-rates. A natural question to ask is: how to we construct a model so as to be consistent with observed European option prices at multiple strikes (at a single maturity) on all three cross-rates. This problem is attempted in [Aus11] but the article does not check that the resulting joint density is non-negative.

Assuming our home currency is dollars, we let  $X$  denote the price of 1 Euro in dollars at time  $T$  in the future (known as the EUR/USD rate), and  $Y$  denote the price of 1 pound in dollars at time  $T$  (the GBP/USD rate). A European option on EUR/GBP (i.e. on the cross-rate  $Z = X/Y$ ) pays  $(Z - K)^+$  pounds at time  $T$ , or equivalently  $Y(Z - K)^+$  dollars. If we assume interest rates are zero for simplicity (otherwise we work with forward rates instead), then given a risk-neutral measure  $\mathbb{Q}$  associated to the US economy, the initial price of such an option (in pounds) is

$$\frac{1}{Y_0} \mathbb{E}^{\mathbb{Q}}(Y(\frac{X}{Y} - K)^+) = \mathbb{E}^{\tilde{\mathbb{Q}}}((Z - K)^+)$$

where  $Y_0 = \mathbb{E}^{\mathbb{Q}}(Y)$ , and we have re-written the expectation on the left side here using the GBP numéraire measure:

$$\tilde{\mathbb{Q}}(A) := \mathbb{E}^{\mathbb{Q}}(\frac{Y}{\mathbb{E}^{\mathbb{Q}}(Y)} A)$$

associated with using GBP as the home currency. From here, we will just use  $\mathbb{E}(\cdot)$  to denote expectations under  $\mathbb{Q}$ .

From the Breeden-Litzenberger formula, we can extract the law of  $X$  and  $Y$  under  $\mathbb{Q}$  from European call prices at all strikes on  $X$ , and the law of  $X/Y$  under  $\tilde{\mathbb{Q}}$ , and in practice this would typically be done using an SVI-type parametrization to interpolate between tradeable strikes with parameters chosen so as to preclude butterfly arbitrage (see the many articles by Gatheral, Jacquier, Martini, Mingone et al. on this theme).

### 2.1 Formulation of the problem

We assume  $\mathbb{R}_+ = (0, \infty)$  throughout. Now suppose we are given three target laws  $\mu_X$ ,  $\mu_Y$  and  $\mu_Z$  on  $\mathbb{R}_+$  and (without loss of generality) we assume that

$$\int_{\mathbb{R}_+} x \mu_X(dx) = \int_{\mathbb{R}_+} y \mu_Y(dy) = \int_{\mathbb{R}_+} z \mu_Z(dz) = 1 \quad (1)$$

i.e. the initial rates  $X_0 = Y_0 = 1$  and hence  $Z_0 = X_0/Y_0 = 1$  as well. Let  $\Pi(\mu, \nu)$  denote the space of joint probability measures on  $\mathbb{R}_+^2$  with marginals  $\mu$  and  $\nu$  respectively. We wish to find a  $\mu^* \in \Pi(\mu_X, \mu_Y)$  such that if  $X, Y$  have joint distribution  $\mu^*$ , then  $X/Y \sim \mu_Z$  under the probability measure  $\tilde{\mathbb{Q}}$  defined above.

**Remark 2.1** If such a  $\mu^*$  exists, it can then be used to price quanto options like e.g. Best-of options which pay  $(X - K_1)^+/K_1 \wedge (Y - K_2)^+/K_2$ , or a quanto option which pays  $(\frac{X}{Y} - K)^+$  dollars (as opposed to EUR).

### 2.2 Technical results

Consider a measurable space  $(\Omega, \mathcal{F})$ , and let  $\mathcal{P}(\Omega)$  the collection of probability measures on  $(\Omega, \mathcal{F})$ . For  $\mathbb{P} \in \mathcal{P}(\Omega)$ ,  $H(\mathbb{Q}|\mathbb{P}) := \mathbb{E}^{\mathbb{Q}}(\log \frac{d\mathbb{Q}}{d\mathbb{P}}) = \mathbb{E}^{\mathbb{P}}(\frac{d\mathbb{Q}}{d\mathbb{P}} \log \frac{d\mathbb{Q}}{d\mathbb{P}})$  denotes the *entropy* of  $\mathbb{Q}$  with respect to  $\mathbb{P}$ . Moreover, since  $h(z) = z \log z$  is convex,  $H(\mathbb{Q}|\mathbb{P}) = \mathbb{E}(h(\frac{d\mathbb{Q}}{d\mathbb{P}})) \geq h(\mathbb{E}(\frac{d\mathbb{Q}}{d\mathbb{P}})) = 0$ .  $H(\mathbb{Q}|\mathbb{P})$  is jointly l.s.c. and convex, see e.g. Lemma 1.3 in [Nutz22]).

To specialize to our problem above, we now let  $\mathcal{E}$  denote the set of probability measures  $\mu$  on  $(\mathbb{R}_+^2, \mathcal{B}(\mathbb{R}_+^2))$  such that  $\int u(x) \mu(dx, dy) = \int u d\mu_X$ ,  $\int v(y) \mu(dx, dy) = \int v d\mu_Y$  and  $\int y w(\frac{x}{y}) \mu(dx, dy) = \int w d\mu_Z$  for all  $u, v, w \in C_b(\mathbb{R}_+)$ , and we make the following standing assumption:

**Assumption 2.1**  $\mathcal{E}$  is non-empty, and contains an element  $\mu$  with  $H(\mu|\bar{\mu}) < \infty$ .

Then  $\mathcal{E}$  is convex, and the following property is readily verified:

**Lemma 2.2**  $\mathcal{E}$  is closed in the weak topology (and hence in the total variation metric).

**Proof.** See Appendix A. ■

Then from Lemma 1.10 in [Nutz22] (or Theorem 2.1 in [Csi75]),  $H(\mu|\bar{\mu})$  attains a unique minimum  $\mu^*$  on  $\mathcal{E}$ , and (from Theorem 2.1 in [Csi75]),

$$\mu^* = e^g \bar{\mu} \quad (2)$$

where  $g$  lies in the closed subspace of  $L^1(\mu^*)$  spanned by the functions of the form  $u(x), v(y)$  and  $yw(\frac{x}{y})$ .

Unfortunately, this does not automatically imply that  $g$  is of the form  $u(x) + v(y) + yw(\frac{x}{y})$ . For this reason, we will not solve the problem  $\inf_{\mu \in \mathcal{E}} H(\mu|\bar{\mu})$  directly or use the result in (2) (which we don't require anyway)<sup>5</sup>, but rather we use

$$\mu^*(x, y) = e^{u^*(x) + v^*(y) + yw^*(\frac{x}{y})} \bar{\mu}(x, y) \quad (3)$$

as an ansatz for an admissible  $\mu$  (i.e. in  $\mathcal{E}$ ), and (assuming  $\mathcal{E}$  is non-empty) we show that the marginals for the associated Sinkhorn scheme (defined below) converge under appropriate conditions on the target laws, which is all we need for our calibration problem. This circumvents the need for dualization arguments as in [Guy20], but we do discuss duality for the finite-option case below where it becomes more practically relevant.

### 2.3 Integral equations for $u, v, w$

From here on we assume the following:

**Assumption 2.3**  $\mu_X, \mu_Y$  and  $\mu_Z$  have densities.

Let  $\bar{\mu}$  denote a reference probability measure, typically chosen to be the product measure of  $\mu_X$  and  $\mu_Y$ , so  $\bar{\mu}$  already satisfies 2 of the 3 marginal constraints. Then using the ansatz in (3) the marginal constraints can be written

$$\begin{aligned} \mu_X(x) &= e^{u(x)} \int_0^\infty e^{v(y) + yw(\frac{x}{y})} \bar{\mu}(x, y) dy \\ \mu_Y(y) &= e^{v(y)} \int_0^\infty e^{u(x) + yw(\frac{x}{y})} \bar{\mu}(x, y) dx \\ \mu_Z(K) &= \int_0^\infty \int_0^\infty y \delta(\frac{x}{y} - K) e^{u(x) + v(y) + yw(\frac{x}{y})} \bar{\mu}(x, y) dy dx \\ &= \int_0^\infty \int_0^\infty \frac{x}{z} \delta_K(dz) e^{u(x) + v(\frac{x}{z}) + \frac{x}{z} w(z)} \frac{x}{z^2} \bar{\mu}(x, \frac{x}{z}) dz dx \\ &= \int_0^\infty \frac{x}{K} e^{u(x) + v(\frac{x}{K}) + \frac{x}{K} w(K)} \frac{x}{K^2} \bar{\mu}(x, \frac{x}{K}) dx \end{aligned} \quad (4)$$

where we have made the transformation  $z = x/y$  (i.e.  $y = x/z$ ), so  $dy = -\frac{x}{z^2} dz$  in the inner integral. We can formally obtain the same three equations using a variational argument as on pg 8 in [Guy20] (see also Remark 3.4 in [Nutz22]).

If the integrals in the expressions for  $\mu_X(x)$  and  $\mu_Y(y)$  are finite and non-zero, the first two equations can be re-arranged as

$$\begin{aligned} u(x) &= \log \mu_X(x) - \log \int_0^\infty e^{v(y) + yw(\frac{x}{y})} \bar{\mu}(x, y) dy \\ v(y) &= \log \mu_Y(y) - \log \int_0^\infty e^{u(x) + yw(\frac{x}{y})} \bar{\mu}(x, y) dx \end{aligned}$$

and from here on we make the usual choice that  $\bar{\mu}(x, y) = \mu_X(x)\mu_Y(y)$ , so these simplify further to

$$u(x) = -\log \int_0^\infty e^{v(y) + yw(\frac{x}{y})} \mu_Y(y) dy, \quad v(y) = -\log \int_0^\infty e^{u(x) + yw(\frac{x}{y})} \mu_X(x) dx$$

and setting  $K = z$ , we see that

$$H(z; u, v, w) = \int_0^\infty e^{u(x) + v(\frac{x}{z}) + \frac{x}{z} w} \frac{x^2}{z^3} \mu_X(x) \mu_Y(\frac{x}{z}) dx - \mu_Z(z) = 0 \quad (5)$$

where (with mild abuse of notation)  $w$  is a scalar here unlike above.

<sup>5</sup>Solving  $\inf_{\mu \in \mathcal{E}} H(\mu|\bar{\mu})$  also lead to issues with having to prove attainment for the dual problem to justify the exponential ansatz which (even for the much simpler problem considered in Theorem 2.1 in [Nutz22]) requires pages of intermediate lemmas to prove, see section 2.3 in [Nutz22] and in particular the discussion below Eq 2.8.

## 2.4 The Sinkhorn scheme

The Sinkhorn-type fixed point iterative scheme to solve these coupled equations is then given by

$$u_{n+1}(x) := -\log \int_0^\infty e^{v_n(y)+yw_n(\frac{x}{y})} \mu_Y(y) dy \quad (6)$$

$$v_{n+1}(y) := -\log \int_0^\infty e^{u_{n+1}(x)+yw_n(\frac{x}{y})} \mu_X(x) dx \quad (7)$$

$$0 = H(z; u^{n+1}, v^{n+1}, w^{n+1}) \quad (8)$$

and we typically set  $u^0 \equiv v^0 \equiv w^0 \equiv 0$ . Note we can re-write Eq (5) as

$$\int_0^\infty e^{u(z y)+v(y)+y w} y^2 \mu_X(y z) \mu_Y(y) dy - \mu_Z(z) = 0$$

by letting  $x = yz$ , so  $dx = z dy$ . Then if  $w^*(z) = \inf\{w : H(z; u, v, w) + \mu_Z(z) = +\infty\}$ , then (from the monotone convergence theorem) we know that

$$\lim_{w \nearrow w^*(z)} \int_0^\infty e^{u(z y)+v(y)+y w} y^2 \mu_X(y z) \mu_Y(y) dy = \int_0^\infty e^{u(z y)+v(y)+y w^*(z)} y^2 \mu_X(y z) \mu_Y(y) dy.$$

The right hand side may or may not be infinite, but we clearly need it to be  $\geq \mu_Z(z)$  to be able to find a root of  $H(z; u, v, w)$ . We now impose more stringent conditions on the target densities to avoid problems with this root finding.

**Assumption 2.4**  $\text{supp}(\mu_X) = [0, \bar{X}]$ ,  $\text{supp}(\mu_Y) = [\underline{Y}, \bar{Y}]$ ,  $\text{supp}(\mu_Z) = [\underline{Z}, \bar{Z}]$  and  $\mu_Y(y) \in [\underline{\mu}_Y, \bar{\mu}_Y]$  and  $\mu_Z(y) \in [\underline{\mu}_Z, \bar{\mu}_Z]$  where all constants here are positive and finite.

Then if  $v_n$  and  $w_n$  lie in finite intervals  $[\underline{v}_n, \bar{v}_n]$  and  $[\underline{w}_n, \bar{w}_n]$ , we see that

$$\begin{aligned} u_{n+1}(x) &\in [-\bar{v}_n - \bar{Y} \bar{w}_n, -\underline{v}_n - \underline{Y} \underline{w}_n] \\ v_{n+1}(y) &\in [-\bar{u}_{n+1} - \bar{Y} \bar{w}_n, -\underline{u}_{n+1} - \underline{Y} \underline{w}_n] \end{aligned}$$

and hence  $u_{n+1}$  and  $v_{n+1}$  are also bounded from below.

Let  $h(x; w, z) = e^{u(x)+v(\frac{x}{z})+\frac{x}{z} w} \mu_X(x) \mu_Y(\frac{x}{z})$  denote the integrand in the expression for  $I_3(w, z) := H(z; u, v, w)$  where we have dropped the dependence on  $u$  and  $v$  to ease notation.  $I_3(w, z)$  is clearly strictly monotonically increasing in  $w$ . Moreover (under Assumption 2.4) if  $u, v$  are bounded then  $H(z; u, v, w)$  is finite for all  $w \in \mathbb{R}$ , and  $h(x; w, z)$  tends monotonically to  $+\infty$  as  $w \rightarrow \infty$  and 0 as  $w \rightarrow -\infty$ . Hence (from the monotone convergence theorem)  $I_3(w, z)$  tends to  $+\infty$  as  $w \rightarrow \infty$  and tends to  $-\mu_Z(z) < 0$  as  $w \rightarrow -\infty$ . Then considering an arbitrary sequence  $w^{(k)} \rightarrow w$ , we see that  $I_3(\cdot, z)$  is continuous (from the bounded convergence theorem), so a unique finite root  $w(z)$  exists (where we have emphasized the dependence on  $z$  here).

If  $u_{n+1}, v_{n+1}$  are bounded from above, then from the third integral equation

$$\int_0^\infty e^{u(x)+v(\frac{x}{z})+\frac{x}{z} w} \mu_Y(\frac{x}{z}) \mu_X(x) dx - \mu_Z(z) = 0$$

we see that  $w_{n+1} \leq \bar{w}$ , where  $e^{\bar{X} \bar{w} / \bar{Z}} e^{\bar{u}_{n+1} + \bar{v}_{n+1}} \frac{\bar{X}^2}{\bar{Z}^3} \bar{\mu}_Y = \bar{\mu}_Z$ , and  $w_{n+1} \geq \underline{w}$  where  $e^{\bar{X} \underline{w} / \underline{Z}} e^{\bar{u}_{n+1} + \bar{v}_{n+1}} \frac{\bar{X}^2}{\underline{Z}^3} \bar{\mu}_Y = \underline{\mu}_Z$ . Hence by induction,  $(u_{n+1}, v_{n+1}, w_{n+1})$  are bounded for each  $n$  so the Sinkhorn scheme doesn't blow up at any finite  $n$ . This does not show that  $(u_n, v_n, w_n)$  are uniformly bounded in  $n$ , but in the next section we prove convergence of the marginals for the scheme in the total variation metric.

The Sinkhorn scheme typically converges very quickly in practice, and note the third equation here requires numerical root-finding as for the SPX-VIX calibration problem discussed in [Guy20], but we have one less dimension here since we only have to compute  $w(z)$  for a range of values for the one argument  $z$  not a two-variable function of the form  $\Delta(s_1, v)$  as in [Guy20].

## 2.5 Order of convergence for the Sinkhorn scheme

Set  $\mu_{i,j,k}(x, y) = e^{u_i(x)+v_j(y)+yw_k(x/y)} \bar{\mu}(x, y)$ , and let  $\mu_{i,j,k}^X(x)$  (resp.  $\mu_{i,j,k}^Y(y)$ ) denote the first (resp. second) marginal of  $\mu_{i,j,k}$ , and  $\mu_{i,j,k}^Z(z)$  denote the  $Z$ -marginal of  $\mu_{i,j,k}$  under  $\tilde{\mathbb{Q}}$ .

**Proposition 2.5** *Under Assumptions 2.1, 2.3 and 2.4, the marginals for the Sinkhorn scheme converge; specifically we have that  $\|\mu_{n,n,n}^X - \mu_X\|_{TV} \rightarrow 0$  and  $\|\mu_{n,n,n}^Y - \mu_Y\|_{TV} \rightarrow 0$  as  $n \rightarrow \infty$  (where  $\|\cdot\|_{TV}$  denotes the total variation distance, see [Nutz22]) and  $\mu_{n,n,n}^Z = \mu_Z$  for all  $n \geq 1$ . Moreover there exists a constant  $A$  such that  $\|(\frac{1}{n} \sum_{i=0}^{n-1} \mu_{i,i,i})^X - \mu_X\|_{TV} \leq \frac{A}{\sqrt{n}}$  and  $\|(\frac{1}{n} \sum_{i=0}^{n-1} \mu_{i,i,i})^Y - \mu_Y\|_{TV} \leq \frac{A}{\sqrt{n}}$  i.e. the marginals of the average of the Sinkhorn iterates have  $O(\frac{1}{\sqrt{n}})$  convergence.*

**Proof.** (of Proposition 2.5). Then (following Lemma 6.4 in [Nutz22]), we see that

$$\begin{aligned} H(\mu_{n+1,n,n}|\mu_{n,n,n}) &= \int \log \frac{d\mu_{n+1,n,n}}{d\mu_{n,n,n}} \mu_{n+1,n,n}(x,y) dx dy = \int \int (u_{n+1}(x) - u_n(x)) \mu_{n+1,n,n}(x,y) dx dy \\ &= \int (u_{n+1} - u_n) d\mu_X \end{aligned} \quad (9)$$

since  $\mu_{n+1,n+1,n}$  has the correct marginal for  $X$  by construction (see also Eq 6.5 in [Nutz22]), and similarly

$$\begin{aligned} H(\mu_{n+1,n+1,n}|\mu_{n+1,n,n}) &= \int (v_{n+1} - v_n) d\mu_{n+1,n+1,n} = \int (v_{n+1} - v_n) d\mu_Y \\ H(\mu_{n+1,n+1,n+1}|\mu_{n+1,n+1,n}) &= \int y(w_{n+1} - w_n) d\mu_{n+1,n+1,n+1} = \int (w_{n+1} - w_n) d\mu_Z \end{aligned} \quad (10)$$

for all  $n \geq 0$ . From Assumption 2.1, there exist a  $\mu_* \in \mathcal{E}$  with  $H(\mu_*|\bar{\mu}) < \infty$ , so (from Lemma 1.4b) in [Nutz22]) we see that

$$H(\mu_*|\bar{\mu}) - H(\mu_*|\mu_{n,n,n}) = \mathbb{E}^{\mu_*}(\log \frac{d\mu_{n,n,n}}{d\bar{\mu}}) \quad (11)$$

which (from the definition of the scheme) we can re-write as

$$\mathbb{E}^{\mu_*}(\log \frac{d\mu_{n,n,n}}{d\bar{\mu}}) = \mathbb{E}^{\mu_*}(u_n(X) + v_n(Y) + Yw_n(X/Y)) = \mathbb{E}^{\mu_X}(u_n(X)) + \mathbb{E}^{\mu_Y}(v_n(Y)) + \mathbb{E}^{\mu_Z}(w_n(Z)). \quad (12)$$

Re-writing the right hand side here as a telescoping sum using the three equations at the beginning of the proof, we see that

$$H(\mu_*|\mu_{n,n,n}) = H(\mu_*|\bar{\mu}) - \sum_{i=0}^{n-1} (H(\mu_{i+1,i,i}|\mu_{i,i,i}) + H(\mu_{i+1,i+1,i}|\mu_{i+1,i,i}) + H(\mu_{i+1,i+1,i+1}|\mu_{i+1,i+1,i})) \quad (13)$$

so  $0 \leq H(\mu_*|\mu_{n,n,n}) \leq H(\mu_*|\bar{\mu})$ , and we see that  $H(\mu_*|\mu_{n,n,n})$  is non-increasing<sup>6</sup> (recall  $H(\mu_*|\bar{\mu})$  is finite by assumption).

The sum on the right in (13) is bounded by  $H(\mu_*|\bar{\mu})$ , so the summand tends to zero. Then from the Data Processing Inequality (Example 1.7 in [Nutz22]), for  $i \geq 1$ , the third term in the summand in (13) dominates  $H(\mu_{i+1,i+1,i+1}^Y|\mu_Y)$  and the first term dominates  $H(\mu_X|\mu_{i,i,i}^X)$  (note the ordering is now the other way round, but this won't matter when we apply Pinsker's inequality below) so these two quantities also tend to zero as  $i \rightarrow \infty$  (so clearly  $H(\mu_{i,i,i}^Y|\mu_Y) \rightarrow 0$  as well), and  $\mu_{i,i,i}^Z$  satisfies the third marginal condition exactly by construction for all  $i \geq 1$ . Then from Pinsker's inequality (see Lemma 1.2 in [Nutz22] with the 2 typo corrected),

$$\begin{aligned} \|\mu_{i,i,i}^X - \mu_X\|_{TV}^2 &\leq \frac{1}{2} H(\mu_X|\mu_{i,i,i}^X) \rightarrow 0 \\ \|\mu_{i,i,i}^Y - \mu_Y\|_{TV}^2 &\leq \frac{1}{2} H(\mu_{i,i,i}^Y|\mu_Y) \rightarrow 0 \end{aligned}$$

so we have convergence for the three marginals as claimed. Finally, to get the order of convergence, from the joint convexity of  $H(.,.)$  and the Data processing inequality,

$$\begin{aligned} H(\mu_X | (\frac{1}{n} \sum_{i=0}^{n-1} \mu_{i,i,i})^X) &= H(\mu_X | \frac{1}{n} \sum_{i=0}^{n-1} \mu_{i,i,i}^X) \leq \frac{1}{n} \sum_{i=0}^{n-1} H(\mu_X | \mu_{i,i,i}^X) \leq \frac{1}{n} \sum_{i=0}^{n-1} H(\mu_{i+1,i,i}|\mu_{i,i,i}) \leq \frac{H(\mu_*|\bar{\mu})}{n} \\ H((\frac{1}{n} \sum_{i=1}^n \mu_{i,i,i})^Y | \mu_Y) &= H(\frac{1}{n} \sum_{i=1}^n \mu_{i,i,i}^Y | \mu_Y) \leq \frac{1}{n} \sum_{i=1}^n H(\mu_{i,i,i}^Y | \mu_Y) \leq \frac{1}{n} \sum_{i=1}^n H(\mu_{i,i,i}|\mu_{i,i,i-1}) \leq \frac{H(\mu_*|\bar{\mu})}{n} \end{aligned}$$

Hence (from Pinsker's inequality)  $\|(\frac{1}{n} \sum_{i=0}^{n-1} \mu_{i,i,i})^X - \mu_X\|_{TV}$  and  $\|(\frac{1}{n} \sum_{i=1}^n \mu_{i,i,i})^Y - \mu_Y\|_{TV}$  are  $O(\frac{1}{\sqrt{n}})$  and we know  $\mu_{i+1,i+1,i+1}^Z = \mu_Z$  by construction, and (from the definition of the total variation metric the contribution from  $\frac{1}{n} \mu_{0,0,0}$  is irrelevant in the limit). ■

**Remark 2.2** See also Section 6.2 in [Nutz22] for discussion on the rate of convergence for Sinkhorn schemes.

<sup>6</sup>From (11) and (12) we also see that  $\mathbb{E}^{\mu_*}(\log \frac{d\mu_{n,n,n}}{d\bar{\mu}}) = \mathbb{E}^{\mu_X}(u_n(X)) + \mathbb{E}^{\mu_Y}(v_n(Y)) + \mathbb{E}^{\mu_Z}(w_n(Z)) \leq H(\mu_*|\bar{\mu}) \leq \infty$  i.e. we have a uniform bound

## 2.6 The finite-option case

In this section, we mimic the setup in sections 1.1 (and section 3.1) for a one-period market model in [FS04] (see also Example 1.18 in [Nutz22]). Assume we now only have  $d + 1$  tradeable assets with (dollar) prices  $S^i$  at time 1 of the form  $g_i(X, Y)$  for some measurable non-negative functions  $g_i$  with  $S^0 = 1$  (i.e. cash), and  $S^1 = X$  and  $S^2 = Y$  (i.e. forwards on  $X$  and  $Y$  respectively), where  $X$  and  $Y$  are defined on the probability space  $(\Omega, \mathcal{F}, \mathbb{P}) = (\mathbb{R}_+^2, \mathcal{B}(\mathbb{R}_+^2), \bar{\mu})$ . As before,  $X$  and  $Y$  represent the EUR/USD and GBP/USD exchange rates at time 1, and we assume these  $d + 1$  assets have (dollar) market prices  $\pi_i$  at time 0. In practice we are typically interested in the case when the assets for  $i > 2$  comprise European calls (resp. puts) on  $X$  and  $Y$  and cross-rate options which pay  $(X - KY)^+$  (resp.  $(KY - X)^+$ ) in the home currency.

We set  $Y^i = S^i - \pi^i$ , and assume interest rates are zero W.L.O.G., and assume the prices  $\pi_i$  are non-arbitrageable (see Eq 3.5 in [FS04] for definition), and the payoffs  $S^i$  are non-redundant (see Definition 3.3 in [FS04]), so in particular do not a forward on the cross-rate  $Z = X/Y$  since this is worth  $X \cdot \frac{Y}{X} = S^1$  dollars at time 1.

Then under the (somewhat stringent) moment condition that  $\mathbb{E}^{\bar{\mu}}(e^{\lambda \cdot Y}) < \infty$  for all  $\lambda \in \mathbb{R}^{d+1}$ , from Corollary 3.27 and Theorem 3.30 in [FS04] (with  $m = 0$ ), there exists a unique equivalent probability measure  $\mu^* \sim \bar{\mu}$  and a  $\lambda^*$  such that

$$H(\mu^* | \bar{\mu}) = \inf_{\mu \sim \bar{\mu} : \mathbb{E}^\mu(Y^i) = 0, \forall i} H(\mu | \bar{\mu}) = \sup_{\lambda \in \mathbb{R}^{d+1}} -\log \mathbb{E}^{\bar{\mu}}(e^{\lambda \cdot Y}) = -\log \mathbb{E}^{\bar{\mu}}(e^{\lambda^* \cdot Y}) \quad (14)$$

with  $\frac{d\mu^*}{d\bar{\mu}} = \frac{e^{\lambda^* \cdot Y}}{\mathbb{E}^{\bar{\mu}}(e^{\lambda^* \cdot Y})}$  (i.e. we minimize the mgf of  $Y$  under  $\bar{\mu}$  which is strictly concave and smooth, see e.g. Lemma 3.20 and Theorem 3.5 in [FS04]). The constraints  $\mathbb{E}^\mu(Y) = 0$  correspond to the first-order optimality conditions (FOC):  $\frac{\partial}{\partial \lambda} \mathbb{E}^{\bar{\mu}}(Y e^{\lambda \cdot Y})|_{\lambda=\lambda^*} = \mathbb{E}^{\bar{\mu}}(Y e^{\lambda^* \cdot Y}) = 0$  (see Proposition 3.9 in [FS04]).

This is essentially the FTAP (in one direction) for a one-period market (see Remark 3.13 in [FS04]), and (14) is the finite-dimensional analog of the concave maximization problem in Eq 5.5 in [Guy20] (appropriately adapted for our problem). Figure 3 displays numerical results for this maximization scheme (using CVXPY with MOSEK) applied to the FX option (and forward) prices in Table 1.

**Remark 2.3** From the envelope theorem, the Lagrange multipliers  $\lambda_i^*$  yield the sensitivities of the certainty equivalent:

$$-\frac{\partial}{\partial \pi_i} \log \mathbb{E}^{\bar{\mu}}(e^{\lambda^* \cdot (S - \pi)}) = \lambda_i^*$$

(recall  $\lambda^*$  depends on  $\pi$ , but the contribution from  $\frac{\partial \lambda^*}{\partial \pi_i}$  doesn't show up in the final answer because due to the FOC).

**Remark 2.4** Note we can either choose  $\bar{\mu}$  to be the real world market model (i.e. the  $\mathbb{P}$ ) measure, in which case the dual (maximization) problem here is the classical problem of maximizing exp utility from the available assets, or (if e.g. we cannot easily estimate  $\mathbb{P}$  from historical data), we can just choose  $\bar{\mu}$  as a fictitious reference measure which we can specify to be consistent with the tradeable European options prices on  $X$  and  $Y$  to give a warm starting point for the concave maximization scheme (see also Remark 13 in [Guy24]).

## 2.7 Using a copula for $\bar{\mu}$

A more flexible choice for  $\bar{\mu}$  is to use e.g. use the Gaussian copula, and fit the correlation parameter  $\rho$  to e.g. historical data or use the *Margrabe formula*:  $\sigma_Z^2 = \sigma_X^2 + \sigma_Y^2 - 2\rho\sigma_X\sigma_Y$  (used for pricing EUR/GBP options under a bivariate Black-Scholes model) to back out  $\rho$  using implied volatilities.

To compute  $\bar{\mu}$  for the Gaussian copula, we set  $X = F_X^{-1}(\Phi(Z_1))$  and  $Y = F_Y^{-1}(\Phi(Z_2))$  where  $Z_1, Z_2 \sim N(0, 1)$  with  $\mathbb{E}(Z_1 Z_2) = \rho$ , then  $X \sim \mu_X$  and  $Y \sim \mu_Y$  as required, and

$$\mathbb{P}(X \leq x, Y \leq y) = \mathbb{P}(F_X^{-1}(\Phi(Z_1)) \leq x, F_Y^{-1}(\Phi(Z_2)) \leq y) = \mathbb{P}(Z_1 \leq \Phi^{-1}(F_X(x)), Z_2 \leq \Phi^{-1}(F_Y(y))).$$

$\bar{\mu}$  is then given by

$$\bar{\mu}(x, y) = \frac{\partial^2}{\partial x \partial y} \mathbb{P}(X \leq x, Y \leq y) = F(x)G(x)f(\Phi^{-1}(F_X(x)), \Phi^{-1}(F_Y(y)))$$

where  $f$  is the joint density of  $Z_1$  and  $Z_2$ , and  $F(x) = \frac{d}{dx}(\Phi^{-1}(F_X(x)))$  and  $G(x) = \frac{d}{dy}(\Phi^{-1}(F_Y(y)))$  (see numerical results in Figure 3).

**Remark 2.5** Using the Margrabe formula with all combinations of the  $5 \times 5 \times 5$  (mid) implied vols in the table below for the EUR-USD-GBP triangle leads to a range for the implied correlation for EUR/USD and GBP/USD of [0.7445, 0.8156].

EUR/USD	1.0567	1.0680	1.0798	1.0950	1.1025
Implied vol	5.69/6.315%	5.621/5.966%	5.54/5.815%	5.509/5.854%	5.45/6.075%
$a, b, \sigma, \rho, m$	-0.0005100	0.009510	0.08579	0.30719	0.03433
GBP/USD	1.2331	1.2480	1.2632	1.2718	1.2919
Implied vol	6.148/7.239%	6.125/6.727%	5.985/6.465%	5.873/6.475%	5.681/6.772%
$a, b, \sigma, \rho, m$	0.0002773	0.002254	0.01867	-0.3272	0.003880
EUR/GBP	0.84261	0.84852	0.85478	0.86142	0.8681
Implied vol	3.414/4.584%	3.59/4.232%	3.66/4.17%	3.735/4.373%	3.681/4.841%
$a, b, \sigma, \rho, m$	0.0001047	0.001959	0.01139	0.2032	-0.001484

Table 1: Table of (bid-ask) implied volatilities with 1-month maturity (with strikes going horizontally in the 1st, 3rd and 5th rows) on 11th Feb 2024, and SVI parameters fitted to the mid-implied vols. The forward prices here are 1.0796 for EUR/USD, 1.2630 for GBP/USD and .85483 for EUR/GBP (data obtained from Bloomberg), and these SVI parameters are used for the Sinkhorn scheme.

### 3 Extremal prices for the cross-rate options given full marginals $\mu_X$ and $\mu_Y$ and Kantorovich duality

In this section we compute robust price bounds (and the corresponding sub/super hedge) for cross-rate options which pay  $(X - KY)^+$  dollars, i.e. we solve the (non-entropic) optimal transport problem.

To this end, we first note the triangle-type inequality:

$$(x - y)^+ \leq (x - x_1)^+ + (x_1 - y)^+ \quad (15)$$

for all  $x, y, x_1 \in \mathbb{R}$  (we just have to check all cases to verify this identity). Now let  $y = p(x) := F_{\mu_Y}^{-1}(1 - F_{\mu_X}(x))$  where  $F_{\mu}(x) := \mu([0, x])$  so  $Y = p(X) \sim \mu_Y$  if  $X \sim \mu_X$  i.e. the Fréchet-Hoeffding lower bound coupling, and note that  $p$  is strictly decreasing if we assume  $\mu_X$  and  $\mu_Y$  have strictly positive densities. We further assume there is a unique root  $x_*$  of  $p(x) = x$ , with  $p(x) > x_*$  for  $x < x_*$  and vice versa (see first plot in Figure 4 below). Then setting  $x_1 = x^*$  and  $y = p(x)$  and assuming  $x > x_*$ , (15) becomes

$$x - p(x) \leq x - x_* + x_* - p(x)$$

(since  $p(x) < x$ ) i.e. an equality. Conversely if  $y = p(x)$  and  $x < x_*$ ,  $p(x) > x_* > x$  so both sides of (15) are zero. Hence  $(x - x_*)^+ + (x_* - y)^+$  (i.e. a call on  $X$  plus a put on  $Y$ , both with strike  $x^*$ ) is a superhedge for  $c(x, y) = (x - y)^+$ , and equality is attained by the coupling  $Y = p(X)$ , so this coupling is optimal for the max problem, i.e.

$$\int_0^1 (F_{\mu_X}^{-1}(u) - F_{\mu_Y}^{-1}(1 - u))^+ du = \sup_{\mu \in \Pi(\mu_X, \mu_Y)} \mathbb{E}^\mu((X - Y)^+)$$

(see also [HLW05],[HLW05b] who look at this problem in the context of basket options). For the general case when  $c(x, y) = (x - Ky)^+$ , we just regard  $KY$  as the new  $Y$  variable, then  $x_*$  is now the root of  $p(x) = Ky$  in the proof above, which is otherwise unchanged except now  $x_*$  depends on  $K$ .

For the lower bound, we first note that

$$(x - y)^+ \geq -(x_1 - x)^+ + (x_1 - y)^+ \quad (16)$$

for all  $x, y, x_1 \in \mathbb{R}$ . Now let  $y = p(x) = F_{\mu_Y}^{-1}(F_{\mu_X}(x))$  so  $Y = p(X) \sim \mu_Y$  if  $X \sim \mu_X$  i.e. the Fréchet-Hoeffding upper bound coupling, and note that  $p$  is strictly increasing if we assume  $\mu_X$  and  $\mu_Y$  have strictly positive densities. We again assume there is a unique root  $x^*$  of  $p(x) = x$ , but now with  $p(x) > x^*$  for  $x > x^*$  and vice versa (see 3rd plot in Figure 4). Then again setting  $x_1 = x^*$  and  $y = p(x)$  and assuming  $x \leq x^*$ , (16) becomes

$$x - p(x) \geq -(x^* - x) + (x^* - p(x)) = x - p(x)$$

so we have equality. Conversely, if  $x > x^*$ , both sides of (16) vanish. Hence (repeating the same arguments as above), the coupling with  $Y = p(X)$  is optimal for the min problem, i.e.

$$\int_0^1 (F_{\mu_X}^{-1}(u) - F_{\mu_Y}^{-1}(u))^+ du = \inf_{\mu \in \Pi(\mu_X, \mu_Y)} \mathbb{E}^\mu((X - Y)^+)$$

and again we can extend this argument to the general case when  $c(x, y) = (x - Ky)^+$ . Note our assumption that  $p(x) = x$  has a unique root fails for our EUR-USD-GBP triangle example (see second plot in Figure 4) where  $p(x) = x$  at three distinct  $x$ -values (see also Corollary 1.2 in [BJ16], although their result requires strict convexity condition on  $c(x, y) = h(y - x)$ ).






Figure 1: Here we have plotted the maximizing  $u$ ,  $v$  and  $w$  respectively after 40 iterations of the Sinkhorn algorithm, using  $[.9, 1.1]^2$  as the domain for numerical integration for  $X, Y$ .




Figure 2: Here we have plotted the Sinkhorn smiles (blue) as a function of moneyiness  $K/F_0$  associated with the  $u$ ,  $v$  and  $w$  in the plot above verses the original (mid)-market smiles (grey crosses) from Table 1 and the SVI interpolated smiles (grey line, which can barely be seen as it's very close to the blue curve) for EUR/USD, GBP/USD and EUR/GBP on 11th Feb 2024. Five options were used for each cross-rate (At-the-money, and .10, .25, .75, and .90 Delta calls, as is customary in FX options markets), using the standard SVI parametrization to interpolate between them and 400 point Gaussian quadrature for the single and double integrals which appear in the Sinkhorn equations, and the bisection method for the root-finding.




Figure 3: Here we have plotted the EUR/USD, GBP/USD and EUR/GBP smiles below) by solving the finite-dimensional concave maximization problem on the right hand side of (14) for the same option data in Table 1, using a Gaussian copula for  $\bar{\mu}$  with correlation  $\rho = .6$ , with weights:

$u = [68.5272, 2.37849786, -5.13262340, 3.39533437, -6.45805503, 8.71979949]$ ,

$v = [-8.66738944e+01, 8.45538689, 6.15522364, 5.80792640, 6.20318995, 9.92863218]$ ,

$w = [0, -6.23988170, -3.90680833e+01, -5.94003754e+01, -1.73172616e+01, -2.17798761e-02]$ . We impose that  $w[0] = 0$ .




Figure 4: On the left we see the optimal transport map  $y = p(x)$  (in blue) from Section 4 using the lower Frechet-Hoeffding bound versus the  $y = x$  line (dashed). In the 2nd plot we see the  $p(x)$  map for the minimal price for EUR/GBP options which comes from the upper Frechet-Hoeffding bound (and we see there are three points of equality which is a problem case), and the final plot shows the same  $p(x)$  function for EUR/USD options (given EUR/JPY and USD/JPY smiles on another date) which only has 1 point of equality (non-problem case) as assumed in the proof in subsection .

**Remark 3.1** For a single cross-rate option paying  $(x - Ky)^+$  with market price  $P$ , we can construct a “model” consistent with  $P$  and the given marginals for  $X$  and  $Y$  if and only if  $P$  lies within the upper and lower price bounds computed above, in which case the model can just be chosen as a weighted coin toss between the two extremal models with the weight chosen to match  $P$ . It is an interesting open problem as to whether we can construct a consistent model if we have prices for two (or  $n$ ) such cross-rate options when the market prices for each these options lie between their respective upper and lower price bounds <sup>7</sup>

---

<sup>7</sup>We thank David Hobson for pointing this out.

## References

- [ACD15] Atar, R., K. Chowdhary, and P. Dupuis, “Robust bounds on risk-sensitive functionals via Rényi divergence”, *SIAM Journal on Uncertainty Quantification*, 3(1):1–30, 2015.
- [AMP25] Acciaio, B., A.Marini, G.Pammer, “Calibration of the Bass Local Volatility model”, *SIAM Journal on Financial Mathematics*, 16/3, 803-833, 2025.
- [Aus11] Austing, P., “Repricing the Cross Smile”, **risk.net**, 2011
- [BBHK20] Backhoff-Veraguas, J., M.Beiglbock, M.Huesmann and S.Källblad, “Martingale Benamou–Brenier: A Probabilistic Perspective”, *The Annals of Probability*, Vol. 48, No. 5, 2258–2289, 2020
- [BST23] Backhoff-Veraguas, J., W.Schachermayer, and B.Tschiderer, “The Bass functional of martingale transport”, preprint, 2023
- [BJ16] Beiglbock, M. and N.Juillet, “On a problem of optimal transport under marginal martingale constraints”, *The Annals of Probability*, 2016, Vol. 44, No. 1, 42–106
- [BCH18] Beiglbock, M., A.M.G.Cox, and M.Huesmann, “Optimal Transport and Skorokhod Embedding”, *Invent. Math.*, 2018
- [Bert15] Bertsekas, D., “Convex Optimization Algorithms”, Athena Scientific, 2015.
- [BG24] Bourgey, F. and J.Guyon, “Fast Exact Joint S&P 500/VIX Smile Calibration in Discrete and Continuous Time”. **risk.net**, Feb 2024
- [Bour25] Bourgey, F. “Fast Smile Calibration in Discrete and Continuous Time Using Sinkhorn Algorithms”, SIAM conference talk, Miami, July 2025.
- [Burke] Burke, J., Lecture Notes, <https://sites.math.washington.edu/~burke/crs/407/notes/section4.pdf>
- [Carr09] Carr, P., “Local Variance Gamma Option Pricing Model”, presentation Bloomberg/Courant Institute, April 28, 2009.
- [CN17] Carr, P. and S.Nadtochiy, “Local Variance Gamma and Explicit Calibration to Option Prices”, *Mathematical Finance*, 27(1):151–193, 2017
- [CL21] Conze, A. and P.Henry-Labordère, “Bass Construction with Multi-Marginals: Lightspeed Computation in a New Local Volatility Model”, preprint, 2021.
- [Csi75] Csiszár, I. “ $I$ -divergence geometry of probability distributions and minimization problems”, *The Annals of Probability*, 3(1):146–158, 1975.
- [dMHL15] de Marco, S. and P.Henry-Labordère, “Linking Vanillas and VIX options: A constrained martingale optimal transport problem”, *SIAM J. Finan. Math*, 4(54), pp. 1171–1194, 2015
- [DZ10] Dembo, A. and O.Zeitouni, “Large deviations techniques and applications”, Jones and Bartlet publishers, Boston, second edition, 2010.
- [F24] Forde, M., “Rough Bergomi revisited - exact and minimal-variance hedging for VIX and European options, and exact calibration to multiple smiles”, preprint, 2024.
- [FS23] Forde, M. and B.Smith, “Markovian stochastic volatility with stochastic correlation - joint calibration and consistency of SPX/VIX short-maturity smiles”, *Int. J. Theor. Appl. Finance*, 26, 2-3, 2023.
- [FS04] Föllmer, H. and A. Schied, “Stochastic Finance: An Introduction in Discrete Time”, W. de Gruyter, Berlin, 3rd edition, 2004.
- [GLOW22] Guo, I., G.Loeper, J.Oblój, S.Wang, “Optimal transport for model calibration”, **risk.net**, Jan 2022
- [GLW22] Guo, I., G.Loeper, S.Wang, “Calibration of Local-Stochastic Volatility Models by Optimal Transport”, *Mathematical Finance*, 32(1), 46-77, 2022.
- [Guy13] Guyon, J., “A New Class of Local Correlation Models”, Quantitative Research, Bloomberg LP, 2013
- [Guy20] Guyon, J., “The Joint S&P 500/VIX Smile Calibration Puzzle Solved”, **risk.net**, April 2020
- [Guy22] Guyon, J., “Dispersion-Constrained Martingale Schrödinger Bridges: Joint Entropic Calibration of Stochastic Volatility Models to S&P 500 and VIX Smiles”, preprint, 2022

- [Guy24] Guyon, J., “Dispersion-Constrained Martingale Schrödinger Problems and the Exact Joint S&P 500/VIX Smile Calibration Puzzle”, *Finance and Stochastics*, 28(1):27-79, 2024
- [Guy24b] Guyon, J., “Fast exact joint S&P 500/VIX smile calibration in discrete and continuous time”, London Mathematical Finance Seminar, Feb 2024
- [GMN17] Guyon, J., R.Menegaux and M.Nutz, “Bounds for VIX Futures given S&P 500 Smiles”, *Finance & Stochastics*, Vol. 21, No. 3, pp. 593-630, 2017
- [dM18] De March, H., “Multi-dimensional Martingale Optimal Transport: Local structure and Numerics”, presentation at Oxford University, 2018
- [HN12] Hobson, D. and A.Neuberger, “Robust bounds for forward start options”, *Mathematical Finance*, 22, 31-56, 2012
- [HK15] Hobson, D. and M.Klimmek, “Robust price bounds for the forward starting straddle”, *Finance and Stochastics*, Vol 19, Issue 1, 189-214, 2015
- [HL19] Henry-Labordère, P., “From (Martingale) Schrodinger bridges to a new class of Stochastic Volatility Models”, preprint, 2019.
- [HLW05] Hobson, D., P.Laurence, T-H.Wang, “Static-arbitrage upper bounds for the prices of basket options”, *Quantitative Finance*, 5 (4) 329-342, 2005
- [HLW05b] Hobson, D., P.Laurence, T-H.Wang, “Static-arbitrage optimal subreplicating strategies for basket options.” *Insurance: Mathematics and Economics*, 37(3):553–572, 2005.
- [HLT16] Henry-Labordere, P. and N.Touzi, “An explicit martingale version of the one-dimensional Brenier theorem”, *Finance and Stochastics*, 20(3):635–668, 2016 (version from 10th April 2015 cited here)
- [HLT16] Henry-Labordere, P. and N.Touzi, Hadrien De March, “Building arbitrage-free implied volatility: Sinkhorn’s algorithm and variants”, preprint.
- [HN19] Norgilas, N., D.Hobson, “Robust bounds for the American put Volume 23, pages 359–395, (2019)
- [HN25] Norgilas, N., D.Hobson, “Model-independent upper bounds for the prices of Bermudan options with convex payoffs”, preprint.
- [Loep25] Loeper, G., “Sinkhorn algorithms for Schrödinger and Bass problems: a PDE point of view”, VCMF conference, Vienna, July 2025
- [JLO23] Joseph, B., G.Loeper and Jan Oblój, “The Measure Preserving Martingale Sinkhorn Algorithm”, preprint
- [Norg25] Norgilas, N., “Model-free price bounds of Bermudan contracts”, SIAM conference talk, Miami, July 2025.
- [Nutz22] Nutz, M., “Introduction to Entropic Optimal Transport”, Lecture Notes, 2022
- [Nutz24] Nutz, M., “Quadratically Regularized Optimal Transport: Existence and Multiplicity of Potentials”, preprint
- [Sion58] Sion, M., “On general minimax theorems”, *Pacific Journal of Mathematics*, 8(1), 171–176, 1958.

## A Proof of Lemma 2.2

Let  $\mu_n \in \mathcal{E}$  with  $\mu_n \xrightarrow{w} \mu$ . Then for any  $u, v \in C_b(\mathbb{R}_+)$ ,  $\int u(x)\mu^X(dx) = \int u d\mu_n \rightarrow \int u d\mu$  and similarly for  $v$ , so  $\mu$  has the correct  $X$  and  $Y$  marginals. Let  $f_K(x, y) := (x - Ky)^+$  and  $f_{K,m} := (x - Ky)^+ \wedge m$ . Then

$$0 \leq f_K - f_{K,m} = ((x - Ky)^+ - m)^+ \leq (x - m)^+ \leq x 1_{x>m}.$$

Fix  $\varepsilon > 0$ . Since  $\int x d\mu_X < \infty$ , we can choose  $m$  sufficiently large so that  $\int x 1_{\{x>m\}} d\mu_X \leq \varepsilon$ , and hence

$$\sup_n \int (f_K - f_{K,m}) d\mu_n \leq \varepsilon, \quad \int (f_K - f_{K,m}) d\mu = \int x 1_{\{x>m\}} d\mu_X \leq \varepsilon$$

since  $\mu^X = \mu_n^X = \mu_X$  (from the first part of the proof). Then from the triangle inequality

$$\left| \int f_K d\mu_n - \int f_K d\mu \right| \leq \int (f_K - f_{K,m}) d\mu_n + \left| \int f_{K,m} d\mu_n - \int f_{K,m} d\mu \right| + \int (f_K - f_{K,m}) d\mu.$$

Using weak convergence for the middle term, we see that

$$\limsup_{n \rightarrow \infty} \left| \int f_K d\mu_n - \int f_K d\mu \right| \leq 2\varepsilon.$$

Then since  $\varepsilon > 0$  is arbitrary, the claim follows.