

Sinkhorn revisited - joint calibration to European and One-Touch prices

Martin Forde

Draft version, 30th Sept 2026

Abstract

We apply a Sinkhorn scheme to the problem of constructing an admissible joint law for X_T and the supremum S_T of a continuous martingale X , given target marginals μ and ν . This corresponds to fitting European and One-Touch prices at all strikes and barrier levels at maturity T . ν necessarily lies between the classical Azéma-Yor and Perkins bounds, but we show this condition is not sufficient with a simple counterexample. Our approach avoids constrained entropy minimization and duality arguments, and after each Sinkhorn cycle, the law of S_T and the Rogers[Rog93] condition are satisfied exactly, and the X_T -marginal converges to μ (under a finite-entropy feasibility assumption) at rate $1/\sqrt{n}$ in the total variation metric. As [Rog12] remarks, this problem can also be solved as an optimal transport linear programming problem where we compute the max or min price of a non-tradable option given tradable options, but the latter typically leads to a sparse joint law which may be less realistic from a financial standpoint.

1 Background

We first recall the following fundamental result (cf. Theorem 3.1 and Lemma 2.3 in [Rog93]):

Theorem 1.1 *Fix a $T > 0$ and $x_0 \in \mathbb{R}$. A probability measure π on $\{(s, x) : -\infty < x \leq s, s \geq x_0\}$ is the joint law of (S_T, X_T) (where $S_t = \sup_{0 \leq u \leq t} X_u$) for some continuous martingale X with $X_0 = x_0$ if and only if $\int |x| \pi(ds, dx) < \infty$, $\int x \pi(ds, dx) = x_0$, $\lim_{a \rightarrow \infty} a \bar{F}_\nu(a) = 0$ and*

$$\eta(ds) = s\nu(ds) - \bar{F}_\nu(s)ds \quad (\text{as measures on } (x_0, \infty)), \quad (1)$$

where $\nu(ds) = \int_{x \leq s} \pi(ds, dx)$, $\eta(ds) = \int_{x \leq s} x \pi(ds, dx)$, and $\bar{F}_\nu(s) = \nu((s, \infty))$.¹

Remark 1.1 We give a sketch proof of the sufficiency part of the [Rog93] proof in Appendix D. Eq (1) has the physical interpretation

$$\mathbb{P}(S_T > b)db = \mathbb{E}((S_T - X_T)1_{S_T \in db}) = \mathbb{E}((b - X_T)1_{S_T \in db}). \quad (2)$$

Remark 1.2 (The integrated form of the Rogers condition) In [Rog12], the characterisation in Theorem 1.1 is stated in the integrated form: for π supported on $\{x \leq s, s \geq x_0\}$ with $\int |x| \pi(ds, dx) < \infty$,

$$\int_{\{s \geq a\}} (x - a) \pi(ds, dx) = 0 \quad \text{for all } a \geq x_0, \quad (3)$$

i.e. $\mathbb{E}(X_T - a; S_T \geq a) = 0$ for all $a \geq x_0$, which is optional stopping at the first hitting time to a . The integrability condition $\int |x| \pi(ds, dx) < \infty$ is a standing hypothesis in both formulations: it is the first condition of Theorem 1.1, and it is what makes η and the integral in (3) well defined. Under it, (3), which in terms of η and ν reads $\eta([a, \infty)) = a\nu([a, \infty))$ for all $a \geq x_0$ is equivalent to the remaining three conditions of Theorem 1.1, namely $\int x \pi(ds, dx) = x_0$, $\lim_{a \rightarrow \infty} a \bar{F}_\nu(a) = 0$ and (1):

- (i) At $a = x_0$, (3) is the mean condition, since π is supported on $s \geq x_0$.
- (ii) For $a > 0$, $0 \leq a \bar{F}_\nu(a) \leq a\nu([a, \infty)) = \eta([a, \infty)) = \int x 1_{\{s \geq a\}} \pi(ds, dx)$. The integrand is dominated by $|x|$, which is π -integrable by the standing hypothesis, and it tends to zero pointwise as $a \rightarrow \infty$ since $S_T < \infty$; hence $\lim_{a \rightarrow \infty} a \bar{F}_\nu(a) = 0$ by dominated convergence.
- (iii) For (1), we differentiate $\eta([a, \infty)) = a\nu([a, \infty))$ in a in the sense of measures to get:

$$-\eta(da) = d_a[a\nu([a, \infty))] = \nu([a, \infty))da - a\nu(da) = \bar{F}_\nu(a)da - a\nu(da) \quad \text{on } (x_0, \infty),$$

which is (1). Conversely, integrating (1) over $[a, \infty)$ recovers (3) for $a > x_0$, and the tail condition makes the boundary term at infinity vanish, and at $a = x_0$ the mean condition supplies the atom $\eta(\{x_0\}) = x_0\nu(\{x_0\})$.

Thus, given integrability, (3) contains the mean and tail conditions automatically, whereas the local form (1) on its own does not, as Example 2.1 shows; it is nevertheless the local form that the Sinkhorn scheme below is built on.

¹See also Remark (i) after Theorem 3.1 in [Rog93]. The results in [Rog93] are formulated for uniformly integrable (UI) martingales on $[0, \infty)$ and their terminal value M_∞ . They apply here because a martingale X on $[0, T]$, extended by $X_t = X_T$ for $t \geq T$, is a UI martingale on $[0, \infty)$ with $X_\infty = X_T$ and $\sup_{t \geq 0} X_t = S_T$, and conversely a UI continuous martingale can be time-changed onto $[0, T]$ without altering its terminal value or its supremum. Theorem 1.1 is thus a restatement of Theorem 3.1 and Lemma 2.3 in [Rog93], together with Remark (i) there for $x_0 \neq 0$.

2 Outline of problem

Our aim is to construct a continuous martingale X with prescribed marginals μ and ν for X_T and S_T . We refer to (1) as the local Rogers condition, and this condition will form the basis of the Sinkhorn scheme below.

Based on Theorem 1.1, we impose the following:

Assumption 2.1 $\int |x|\mu(dx) < \infty$, $\int x\mu(dx) = x_0$ and $\lim_{a \rightarrow \infty} a\bar{F}_\nu(a) = 0$, and ν lies between the Perkins and Azéma-Yor model-independent bounds, i.e.

$$\sup_{K < B} \left\{ \mu([B, \infty)) + \frac{C_\mu(B) - P_\mu(K)}{B - K} \right\} \leq \nu([B, \infty)) \leq \inf_{K < B} \frac{C_\mu(K)}{B - K}$$

² for all $B \geq x_0$, where $C_\mu(K) = \int (x - K)^+ \mu(dx)$ and $P_\mu(K) = \int (K - x)^+ \mu(dx)$ denote the call and put pricing functions.

Remark 2.1 If we relax the condition that X is continuous, the upper bound remains unchanged but the Perkins bound is replaced by the trivial lower bound $\mathbb{P}^{X \sim \mu}(\max(X, X_0) \geq B)$, and there is an explicit solution to the problem addressed in this paper described at the end of page 459 and page 460 in [Rog93] using a “lost time” excursion theory construction, whereby X_T and S_T are linked via the simple co-monotonic coupling, so there is no need for a Sinkhorn scheme in this case.

Example 2.1 (Brownian motion stopped at a lower barrier) (see Remark (ii) after Theorem 3.1 of [Rog93]). Let B be a standard Brownian motion with $B_0 = 0$, let $\tau = \inf\{t \geq 0 : B_t = -1\}$, and let $X_t = B_{t \wedge \tau}$. Then X is a continuous local martingale with $X_\infty = -1$ almost surely, and its supremum $S_\infty = \sup_{t \geq 0} X_t$ has law

$$\nu(ds) = \frac{ds}{(1+s)^2}, \quad s \geq 0, \quad \text{so that} \quad \bar{F}_\nu(s) = \nu((s, \infty)) = \frac{1}{1+s}.$$

The joint law of (S_∞, X_∞) satisfies the local Rogers condition (1) on $(0, \infty)$, but X is not uniformly integrable: $\mathbb{E}(X_\infty) = -1 \neq 0$ and $s\bar{F}_\nu(s) \rightarrow 1$ as $s \rightarrow \infty$.

Proof. For $s \geq 0$, the event $\{S_\infty > s\}$ is the event that B reaches s before -1 , which has probability $1/(1+s)$ (the gambler’s ruin probability, obtained by applying the optional stopping theorem to B stopped at the first exit from $(-1, s)$); this gives $\bar{F}_\nu$ and, on differentiating, the density $\nu(s) = (1+s)^{-2}$. Since $X_\infty = -1$ on every path, the measure $\eta(ds) = \int x\pi(ds, dx)$ is $\eta(ds) = -\nu(ds)$. Hence, for $s > 0$,

$$s\nu(s) - \bar{F}_\nu(s) = \frac{s}{(1+s)^2} - \frac{1}{1+s} = \frac{s - (1+s)}{(1+s)^2} = -\frac{1}{(1+s)^2} = -\nu(s)$$

so $\eta(ds) = s\nu(ds) - \bar{F}_\nu(s)ds$ on $(0, \infty)$, which is Eq (1). On the other hand $\mathbb{E}X_\infty = -1 \neq X_0$, so X is not a UI martingale, and correspondingly $s\bar{F}_\nu(s) = s/(1+s) \rightarrow 1 \neq 0$, so the tail condition of Theorem 1.1 fails. ■

As alluded to in the abstract, the following counterexample shows that ν being a probability measure and lying between the Perkins and Azéma-Yor bounds is not a sufficient condition for an admissible joint law to exist.

Counterexample 2.1 Let $\mu \sim U(-1, 1)$, so $x_0 = 0$, and let ν be the law on $(0, 1)$ with tail cdf

$$\bar{F}_\nu(s) = 1 - s - \frac{7}{2}s(1-s)^5, \quad 0 \leq s \leq 1.$$

The Azéma-Yor and Perkins bounds for μ are $1-s$ and $1-\sqrt{s}$ respectively, and $1-\sqrt{s} \leq \bar{F}_\nu(s) \leq 1-s$ on $(0, 1)$, so ν satisfies Assumption 2.1, but there is no continuous martingale X with $X_T \sim \mu$ and $S_T \sim \nu$.

Proof. Suppose such a martingale exists, and let π be the joint law of (S_T, X_T) . By Theorem 1.1, π satisfies (1): $\eta(ds) = s\nu(ds) - \bar{F}_\nu(s)ds$ on $(0, \infty)$, where $\eta(ds) = \int_{x \leq s} x\pi(ds, dx)$. Since $X_T \geq -1$ almost surely, $\eta(ds) \geq -\int_{-1 \leq x \leq s} \pi(ds, dx) = -\nu(ds)$. Substituting the expression for η from Eq (1) into the left-hand side, this reads $s\nu(ds) - \bar{F}_\nu(s)ds \geq -\nu(ds)$, i.e.

$$(1+s)\nu(ds) \geq \bar{F}_\nu(s)ds \quad \text{on } (0, 1),$$

so $(1+s)\nu(s) \geq \bar{F}_\nu(s)$ for Lebesgue almost every $s \in (0, 1)$. But this inequality is violated for $\nu(s) = -\bar{F}'_\nu(s) = 1 + \frac{7}{2}(1-s)^4(1-6s)$ on a neighbourhood of $s = \frac{1}{3}$. ■

Motivated by financial applications, we henceforth assume μ only has support on the positive real line. If X is an asset price and interest rates are zero, μ can be inferred from European option prices via the Breeden-Litzenberger formula, and we can interpret $\nu([B, \infty))$ to be the price of a One-Touch option on X with barrier level $B > X_0$.

²the bounds are sharp here, and attained by the Perkins and Azéma-Yor Skorokhod embeddings respectively, see the original papers [AY79],[Per86] and [Obl04],[Cox04],[Hob11] for details.

2.1 Sinkhorn Integral equations

We henceforth further assume that μ, ν have densities with respect to the Lebesgue measure, and set

$$r(s) = s\nu(s) - \bar{F}_\nu(s)$$

i.e. the density on the right-hand side of (1).

Let $p(s, x)$ be a candidate joint density for (S, X) . Based on the $\mathbb{E}(X_T 1_{S_T \in db})$ term in (2) and the entropic optimal transport literature, we make the exponential ansatz:

$$p(s, x) = R(s, x)e^{u(x)+v(s)+x\lambda(s)}$$

where R is a reference density (see Assumption 2.2 for conditions on R). Then the X -marginal condition is

$$\mu(x) = e^{u(x)} \int_x^\infty R(s, x)e^{v(s)+\lambda(s)x} ds$$

which we can rearrange as

$$u(x) = \log \mu(x) - \log \left(\int_x^\infty R(s, x)e^{v(s)+\lambda(s)x} ds \right) \quad (4)$$

if the integral term lies in $(0, \infty)$ (this will be addressed in Lemma C.1). Similarly, the S -marginal condition is

$$\nu(s) = e^{v(s)} \int_0^s R(s, x)e^{u(x)+\lambda(s)x} dx$$

and the local Rogers condition in (1) is

$$r(s) = e^{v(s)} \int_0^s x R(s, x)e^{u(x)+\lambda(s)x} dx.$$

Dividing by the S -marginal equation then gives

$$\frac{\int_0^s x R(s, x)e^{u(x)+\lambda(s)x} dx}{\int_0^s R(s, x)e^{u(x)+\lambda(s)x} dx} = b(s) := \frac{r(s)}{\nu(s)} \quad (5)$$

where $b(s)$ is defined for ν -almost every s . Under suitable technical conditions (see Lemma 2.3 for details), $\lambda(s)$ can be computed from (5) as a one-dimensional root-finding exercise. The S -marginal equation gives

$$v(s) = \log \nu(s) - \log \left(\int_0^s R(s, x)e^{u(x)+\lambda(s)x} dx \right). \quad (6)$$

Hence we see that (4), (5) and (6) are the three coupled equations for u, λ and v .

2.2 The Sinkhorn scheme

The coupled equations above suggest the following iterative scheme. Set $u_0 = v_0 = \lambda_0 = 0$. Given (u_n, v_n, λ_n) , first set

$$u_{n+1}(x) = \log \mu(x) - \log \left(\int_x^\infty R(s, x)e^{v_n(s)+\lambda_n(s)x} ds \right). \quad (7)$$

This enforces the X -marginal exactly. Next, choose $\lambda_{n+1}(s)$ as the solution of

$$\frac{\int_0^s x R(s, x)e^{u_{n+1}(x)+\lambda_{n+1}(s)x} dx}{\int_0^s R(s, x)e^{u_{n+1}(x)+\lambda_{n+1}(s)x} dx} = b(s). \quad (8)$$

This is a scalar root-finding problem in λ for each s ; existence and uniqueness of the root for ν -almost every s are established in Lemma 2.3. Finally set

$$v_{n+1}(s) = \log \nu(s) - \log \left(\int_0^s R(s, x)e^{u_{n+1}(x)+\lambda_{n+1}(s)x} dx \right). \quad (9)$$

The next u -update restores the X -marginal, which is the only constraint disturbed by the completed (λ, v) update. In practice, one iteration therefore consists of an explicit u -update, one scalar root-find for $\lambda_{n+1}(s)$ for every s , and an explicit v -update.

Definition 2.1 *A probability measure on $\{(s, x) : -\infty < x \leq s < \infty, s \geq x_0\}$ is called **admissible** if it has X -marginal μ , S -marginal ν , and satisfies the conditions in Theorem 1.1.*

For probability measures P and Q we write $H(P|Q) = \int \log \frac{dP}{dQ} dP$ if $P \ll Q$ and $H(P|Q) = +\infty$ otherwise. The integral is well defined in $[0, \infty]$: writing $\rho = dP/dQ$, $\int (\log \rho)^- dP = \int \rho (\log \rho)^- dQ \leq 1/e$, since $\rho \log \rho \geq -1/e$ for $\rho \geq 0$, and $H(P|Q) \geq 0$ by Jensen's inequality.³

From here on we make the following further assumption:

Assumption 2.2 $\mu = 0$ outside $0 \leq x \leq K$, $\nu = 0$ outside $X_0 \leq s \leq K$ for some $K < \infty$ ⁴, and $\mu(x) > 0$ for Lebesgue almost every $x \in (0, K)$ and $\nu(s) > 0$ for Lebesgue almost every $s \in (X_0, K)$. The reference density R is a probability density supported on

$$s \geq X_0, \quad 0 < x \leq s$$

with $R(s, x) > 0$ for every $s \geq X_0$ and Lebesgue almost every $x \in (0, s)$. Finally, there exists an admissible probability density p^* such that $H(p^*|R) < \infty$.

Since $H(p^*|R) < \infty$, we have $p^* \ll R$, so $0 < X \leq S$ p^* -almost surely. Since p^* is a density, $p^*(X = S) = 0$. Hence, using $b(s) = \mathbb{E}_{p^*}(X|S = s)$, (by the local Rogers condition), we obtain

$$0 < b(s) < s \tag{10}$$

for ν -almost every s .

Remark 2.2 (Reference density for Geometric Brownian motion) A natural choice for the reference density R is the joint law of (S_T, X_T) for driftless geometric Brownian motion $X_t = X_0 e^{\sigma W_t - \sigma^2 t/2}$. Writing $Y_t = \log X_t$, a Brownian motion with drift $-\sigma^2/2$ and volatility σ started at $\log X_0$, the reflection principle and Girsanov's theorem give the joint density of $M_T = \max_{t \leq T} Y_t$ and Y_T , and the change of variables $s = e^b$, $x = e^y$ then yields

$$R(s, x) = \frac{2 \ell(s, x)}{\sigma^3 \sqrt{2\pi T^3} s x} \sqrt{\frac{X_0}{x}} \exp\left(-\frac{\ell(s, x)^2}{2\sigma^2 T} - \frac{\sigma^2 T}{8}\right), \quad \ell(s, x) = \log \frac{s^2}{x X_0},$$

for $x > 0$ and $s \geq \max(x, X_0)$, and $R(s, x) = 0$ otherwise. Its S -marginal is

$$\bar{F}_R(B) = \int_B^\infty \int_0^s R(s, x) dx ds = \Phi\left(\frac{-\log(B/X_0) - \sigma^2 T/2}{\sigma \sqrt{T}}\right) + \frac{X_0}{B} \Phi\left(\frac{-\log(B/X_0) + \sigma^2 T/2}{\sigma \sqrt{T}}\right), \quad B \geq X_0,$$

where Φ is the standard normal distribution function. Since geometric Brownian motion is a continuous martingale, R satisfies the local Rogers condition (1) exactly, and it is positive on the interior of its support, as required in Assumption 2.2. Note however that R is not supported on $s \leq K$, so $q_0 = R$ charges the region $\{x > K\}$; this is why several statements in Section 2 and Appendix C are restricted to $n \geq 1$.

Lemma 2.3 Suppose Assumption 2.2 holds. Let u be measurable and finite Lebesgue almost everywhere on $(0, K)$ and suppose that, for ν -almost every s , there exists at least one finite value of λ for which $0 < \int_0^s R(s, x) e^{u(x) + \lambda x} dx < \infty$. Then for ν -almost every s there exists a unique finite $\lambda(s)$ such that

$$\frac{\int_0^s x R(s, x) e^{u(x) + \lambda(s)x} dx}{\int_0^s R(s, x) e^{u(x) + \lambda(s)x} dx} = b(s). \tag{11}$$

Moreover, λ admits a measurable version, and for ν -almost every s , the integral $\int_0^s R(s, x) e^{u(x) + \lambda x} dx$ is positive and finite for every finite λ , and in particular for $\lambda = \lambda(s)$.

Proof. See Appendix A. ■

3 Convergence for the Sinkhorn scheme

For the convergence argument, define $p_{i,j,k}(s, x) = R(s, x) e^{u_i(x) + v_j(s) + \lambda_k(s)x}$. Thus one complete cycle is $p_{n,n,n} \rightarrow p_{n+1,n,n} \rightarrow p_{n+1,n+1,n} \rightarrow p_{n+1,n+1,n+1}$, corresponding to the u -update, the λ -update and the v -update respectively. By Eq (7), $(p_{n+1,n,n})^X = \mu$, so $p_{n+1,n,n}$ is a probability density. By Eq (9), $(p_{n+1,n+1,n+1})^S = \nu$, so every completed iterate is also a probability density. Moreover, Eq (8) gives

$$\frac{\int_0^s x p_{n+1,n+1,n+1}(s, x) dx}{\int_0^s p_{n+1,n+1,n+1}(s, x) dx} = b(s).$$

Since the denominator is $\nu(s)$ by construction, we see that $\int_0^s x p_{n+1,n+1,n+1}(s, x) dx = b(s)\nu(s) = r(s)$. Hence every completed iterate $p_{n,n,n}$ with $n \geq 1$ not only has the correct S -marginal but satisfies the local Rogers condition. The intermediate density $p_{n+1,n,n+1}$ need not have mass one, so in the entropy calculation below the λ - and v -updates are treated together.

³The same bound justifies computing $H(P|Q)$ as a double integral computed one variable at a time whenever P and Q are given by densities.

⁴note there is mild overloading of notation here; this K is not the strike price variable K in Assumption 2.1

Lemma 3.1 For every $n \geq 0$ for which the u -update is well defined,

$$\frac{p_{n+1,n,n}(s, x)}{p_{n,n,n}(s, x)} = \frac{\mu(x)}{(p_{n,n,n})^X(x)} \quad (12)$$

and

$$H(p_{n+1,n,n}|p_{n,n,n}) = H(\mu|(p_{n,n,n})^X). \quad (13)$$

Proof. Since $p_{n+1,n,n}$ and $p_{n,n,n}$ differ only in the u -potential, $p_{n+1,n,n}(s, x)/p_{n,n,n}(s, x) = e^{u_{n+1}(x)-u_n(x)}$. Integrating $p_{n,n,n}(s, x) = R(s, x)e^{u_n(x)+v_n(s)+\lambda_n(s)x}$ over s gives $(p_{n,n,n})^X(x) = e^{u_n(x)} \int_x^\infty R(s, x)e^{v_n(s)+\lambda_n(s)x} ds$, so Eq (7) can be written as $e^{u_{n+1}(x)} = \mu(x)e^{u_n(x)}/(p_{n,n,n})^X(x)$, which gives Eq (12). Hence, since $(p_{n+1,n,n})^X = \mu$,

$$H(p_{n+1,n,n}|p_{n,n,n}) = \int \int p_{n+1,n,n}(s, x) \log \frac{\mu(x)}{(p_{n,n,n})^X(x)} dx ds = \int \mu(x) \log \frac{\mu(x)}{(p_{n,n,n})^X(x)} dx = H(\mu|(p_{n,n,n})^X),$$

which is Eq (13). ■

Proposition 3.2 Suppose Assumption 2.2 holds. Then $\|(p_{n,n,n})^X - \mu\|_{\text{TV}} \rightarrow 0$ as $n \rightarrow \infty$.⁵

Proof. By Lemma C.1, every step of the scheme is well defined and $H(p^*|p_{n,n,n})$ and $H(p^*|p_{n+1,n,n})$ are finite for every $n \geq 0$. Adding Eqs (C-1) and (C-2), the term $H(p^*|p_{n+1,n,n})$ cancels and we obtain, for every $n \geq 0$,

$$H(p_{n+1,n,n}|p_{n,n,n}) + H(p_{n+1,n+1,n+1}|p_{n+1,n,n}) = H(p^*|p_{n,n,n}) - H(p^*|p_{n+1,n+1,n+1}). \quad (14)$$

Now sum Eq (14) over $n = 0, 1, \dots, N-1$. On the right-hand side the sum telescopes, since the second term for index n cancels the first term for index $n+1$, leaving only the first term for $n = 0$ and the second term for $n = N-1$:

$$\begin{aligned} \sum_{n=0}^{N-1} (H(p_{n+1,n,n}|p_{n,n,n}) + H(p_{n+1,n+1,n+1}|p_{n+1,n,n})) &= \sum_{n=0}^{N-1} (H(p^*|p_{n,n,n}) - H(p^*|p_{n+1,n+1,n+1})) \\ &= H(p^*|p_{0,0,0}) - H(p^*|p_{N,N,N}) \\ &= H(p^*|R) - H(p^*|p_{N,N,N}) \leq H(p^*|R), \end{aligned}$$

since $p_{0,0,0} = R$ and entropy is non-negative. Letting $N \rightarrow \infty$ gives

$$\sum_{n=0}^{\infty} (H(p_{n+1,n,n}|p_{n,n,n}) + H(p_{n+1,n+1,n+1}|p_{n+1,n,n})) \leq H(p^*|R) \quad (15)$$

so in particular, $H(p_{n+1,n,n}|p_{n,n,n}) \rightarrow 0$. By Lemma 3.1, $H(p_{n+1,n,n}|p_{n,n,n}) = H(\mu|(p_{n,n,n})^X)$. Therefore $H(\mu|(p_{n,n,n})^X) \rightarrow 0$ and Pinsker's inequality gives $\|(p_{n,n,n})^X - \mu\|_{\text{TV}} \rightarrow 0$. ■

3.1 Rate of convergence

Proposition 3.3 Under the assumptions of Proposition 3.2, for every $n \geq 1$, $H((p_{n,n,n})^X|\mu) \leq \frac{H(p^*|R)}{n}$. Hence $\|(p_{n,n,n})^X - \mu\|_{L^1} \leq \sqrt{\frac{2H(p^*|R)}{n}}$, and $(p_{n,n,n})^X \rightarrow \mu$ in total variation at rate $O(n^{-1/2})$.

Proof. Write $q_n = p_{n,n,n}$. Since $(p_{n+1,n,n})^X = \mu$, the Data Processing inequality (DPI, cf. Lemma 1.6 in [Nutz22]) gives

$$H(q_{n+1}^X|\mu) \leq H(q_{n+1}|p_{n+1,n,n}). \quad (16)$$

From Eq (12) $p_{n+1,n,n}(s, x)/q_n(s, x) = \mu(x)/q_n^X(x)$ for $n \geq 1$, so $q_n(s, x)/p_{n+1,n,n}(s, x) = q_n^X(x)/\mu(x)$ q_n -almost everywhere, since for $n \geq 1$ the density q_n^X is supported in $(0, K)$, where $\mu > 0$ almost everywhere. Hence

$$H(q_n|p_{n+1,n,n}) = \int \log \frac{q_n(s, x)}{p_{n+1,n,n}(s, x)} q_n(s, x) ds dx = \int \log \frac{q_n^X(x)}{\mu(x)} q_n(s, x) ds dx = \int \log \frac{q_n^X(x)}{\mu(x)} q_n^X(x) dx = H(q_n^X|\mu). \quad (17)$$

In particular, by Eq (16) with $n-1$ in place of n and Eq (15), $H(q_n|p_{n+1,n,n}) = H(q_n^X|\mu) \leq H(q_n|p_{n,n-1,n-1}) \leq H(p^*|R) < \infty$ for $n \geq 1$.

For $n \geq 1$, both q_n and q_{n+1} have S -marginal ν and conditional mean $b(s)$ by Lemma C.1. Moreover, as shown in the proof of Lemma C.1, $q = p_{n+1,n,n}$ satisfies the hypotheses of Lemma B.1, with $\lambda(s) = \lambda_{n+1}(s) - \lambda_n(s)$ and with $q' = p_{n+1,n+1,n+1} = q_{n+1}$. Hence Lemma B.1, with $p = q_n$, gives

$$H(q_n|p_{n+1,n,n}) = H(q_n|q_{n+1}) + H(q_{n+1}|p_{n+1,n,n}). \quad (18)$$

⁵see section 1.1 in [Nutz22] for the definition of the total variation distance $\|\cdot\|_{\text{TV}}$ here

By Eq (18), $H(q_{n+1}|p_{n+1,n,n}) \leq H(q_n|p_{n+1,n,n})$. Combining this with Eqs (16) and (17) gives

$$H(q_{n+1}^X|\mu) \leq H(q_{n+1}|p_{n+1,n,n}) \leq H(q_n|p_{n+1,n,n}) = H(q_n^X|\mu).$$

On the other hand, Eqs (16) and (15) give

$$\sum_{n=1}^{\infty} H(q_n^X|\mu) \leq \sum_{n=0}^{\infty} H(q_{n+1}|p_{n+1,n,n}) \leq H(p^*|R).$$

Since $H(q_n^X|\mu)$ is non-increasing and non-negative, $nH(q_n^X|\mu) \leq \sum_{k=1}^n H(q_k^X|\mu) \leq H(p^*|R)$, which proves the first claim. Finally, Pinsker's inequality gives $\|q_n^X - \mu\|_{L^1} \leq \sqrt{2H(q_n^X|\mu)} \leq \sqrt{\frac{2H(p^*|R)}{n}}$. ■




Figure 1: Here we have plotted u, v, λ for a target law ν which is a 60/40 mixture of the Azéma-Yor and Perkins solutions respectively (and hence admissible) with $\mu \sim U[-1, 1]$, using 400 (equidistant) steps for X direction and 200 for S .




Figure 2: Here we have plotted the error for the Rogers condition and the μ -marginal condition for the plots above.




Figure 3: Here we have plotted u, v, λ for a target law ν with cdf $\nu([0, s]) = s^p$ for $s \in [0, 1]$ with $p = 3/4$, again with $\mu \sim U[-1, 1]$.

References

- [AY79] Azéma, J. and M.Yor, “Une solution simple au problème de Skorokhod”, *Séminaire de Probabilités XIII*, Lecture Notes in Mathematics 721, Springer, 90–115, 1979.
- [BHR01] Brown, H., Hobson, D., Rogers, L.C.G., “Robust hedging of barrier options”, *Mathematical Finance*, 11(3), 285–314, 2001.

- [BBBL26] Backhoff, J., M. Beiglböck, G. Bifronte, A. Ley., “Bridging Classical and Martingale Schrödinger Bridges”, preprint
- [Cox04] Cox, A.M.G., “Skorokhod embeddings: non-centred target distributions, diffusions and minimality”, PhD thesis, University of Bath, 2004.
- [DR15] Duembgen, M., Rogers, L.C.G., “The joint law of the extrema, final value and signature of a stopped random walk”, in *In Memoriam Marc Yor – Séminaire de Probabilités XLVII*, Lecture Notes in Mathematics 2137, Springer, 321–338, 2015.
- [For26] Forde, M., “Solving the FX cross-smiles problem - rate of convergence for Sinkhorn marginals, and the finite-option case”, *Statistics and Probability Letters*, 237, 110805, 2026.
- [Guy20] Guyon, J., “The Joint S&P 500/VIX Smile Calibration Puzzle Solved”, *Risk*, April 2020.
- [HOST16] Henry-Labordère, P., Oblój, J., Spoida, P., Touzi, N., “The maximum maximum of a martingale with given n marginals”, *Annals of Applied Probability*, 26(1), 1–44, 2016.
- [Hob11] Hobson, D., “The Skorokhod embedding problem and model-independent bounds for option prices”, in *Paris–Princeton Lectures on Mathematical Finance 2010*, Lecture Notes in Mathematics 2003, Springer, 267–318, 2011.
- [Nutz22] Nutz, M., “Introduction to Entropic Optimal Transport”, Lecture Notes, 2022.
- [Obl04] Oblój, J., “The Skorokhod embedding problem and its offspring”, *Probability Surveys*, 1, 321–392, 2004.
- [Per86] Perkins, E., “The Cereteli–Davis solution to the H^1 -embedding problem and an optimal embedding in Brownian motion”, in *Seminar on Stochastic Processes, 1985*, Progress in Probability and Statistics 12, Birkhäuser, Boston, 172–223, 1986.
- [Rog93] Rogers, L.C.G., “The joint law of the maximum and terminal value of a martingale”, *Probability Theory and Related Fields*, 95(4), 451–466, 1993.
- [Rog12] Rogers, L.C.G., “Extremal martingales”, Talk, Statistical Laboratory, University of Cambridge, 2012.

A Proof of Lemma 2.3

Proof. Fix an s for which the hypothesis of the lemma holds, i.e. for which $Z_s(\lambda) \in (0, \infty)$ for some finite λ , where $Z_s(\lambda) = \int_0^s R(s, x)e^{u(x)+\lambda x}dx$; by assumption this is the case for ν -almost every s . Suppose that, for some finite λ_2 ,

$$0 < Z_s(\lambda_2) = \int_0^s R(s, x)e^{u(x)+\lambda_2 x}dx < \infty.$$

Then, for any other finite λ_1 ,

$$Z_s(\lambda_1) = \int_0^s e^{(\lambda_1-\lambda_2)x} R(s, x)e^{u(x)+\lambda_2 x}dx.$$

Since $x \in (0, s)$, the factor $e^{(\lambda_1-\lambda_2)x}$ is bounded above and below by strictly positive constants. Hence (by comparison with the previous expression) $0 < Z_s(\lambda_1) < \infty$. Therefore $Z_s(\lambda)$ is positive and finite for every finite λ . Since $x \in (0, s)$, differentiation under the integral sign is justified locally in λ by dominated convergence. Moreover

$$\frac{d}{d\lambda} \log Z_s(\lambda) = \frac{\int_0^s x R(s, x)e^{u(x)+\lambda x}dx}{\int_0^s R(s, x)e^{u(x)+\lambda x}dx} =: m_s(\lambda),$$

which is the mean of x under the probability density $R(s, x)e^{u(x)+\lambda x}/Z_s(\lambda)$ on $(0, s)$, and (using the quotient rule) we can easily check that $m'_s(\lambda)$ is the corresponding variance,

$$m'_s(\lambda) = \int_0^s (x - m_s(\lambda))^2 \frac{R(s, x)e^{u(x)+\lambda x}}{Z_s(\lambda)} dx.$$

By Assumption 2.2, $R(s, x) > 0$ for Lebesgue almost every $x \in (0, s)$. Therefore, since $u(x)$ is finite Lebesgue almost everywhere, we have $\frac{R(s, x)e^{u(x)+\lambda x}}{Z_s(\lambda)} > 0$ for Lebesgue almost every $x \in (0, s)$. The integrand is strictly positive for Lebesgue almost every $x \in (0, s)$ except the single point $x = m_s(\lambda)$. Therefore

$$m'_s(\lambda) > 0$$

for every finite λ . Thus m_s is continuous and strictly increasing. Moreover, since $R(s, x)e^{u(x)} > 0$ for Lebesgue almost every $x \in (0, s)$, for every $\varepsilon \in (0, s)$ we have $\int_0^\varepsilon R(s, x)e^{u(x)}dx > 0$ and $\int_{s-\varepsilon}^s R(s, x)e^{u(x)}dx > 0$. Now fix $\lambda > 0$. Since

$$Z_s(\lambda) = \int_0^s R(s, x)e^{u(x)}e^{\lambda x}dx,$$

and the integrand is non-negative, $Z_s(\lambda) \geq \int_{s-\varepsilon}^s R(s, x) e^{u(x)} e^{\lambda x} dx$. For $x \in (s - \varepsilon, s)$, $e^{\lambda x} \geq e^{\lambda(s-\varepsilon)}$, so

$$Z_s(\lambda) \geq e^{\lambda(s-\varepsilon)} \int_{s-\varepsilon}^s R(s, x) e^{u(x)} dx.$$

Also, since $x < s$, $e^{\lambda x} \leq e^{\lambda s}$, and hence $Z_s(\lambda) \leq e^{\lambda s} \int_0^s R(s, x) e^{u(x)} dx = e^{\lambda s} Z_s(0)$. Therefore

$$e^{\lambda(s-\varepsilon)} \int_{s-\varepsilon}^s R(s, x) e^{u(x)} dx \leq Z_s(\lambda) \leq e^{\lambda s} Z_s(0).$$

Taking logarithms and dividing by λ gives

$$s - \varepsilon + o(1) \leq \frac{\log Z_s(\lambda)}{\lambda} \leq s + o(1)$$

as $\lambda \rightarrow \infty$. Since $\varepsilon > 0$ is arbitrary, $\lim_{\lambda \rightarrow \infty} \log Z_s(\lambda)/\lambda = s$. Similarly, using $\int_0^\varepsilon R(s, x) e^{u(x)} dx > 0$ and $e^{\lambda x} \geq e^{\lambda \varepsilon}$ for $x \in (0, \varepsilon)$ and $\lambda < 0$, $\lim_{\lambda \rightarrow -\infty} \frac{\log Z_s(\lambda)}{\lambda} = 0$. We now use the following elementary fact: for a differentiable convex function f on $\mathbb{R}$, the limits $\lim_{\lambda \rightarrow \pm\infty} f'(\lambda)$ exist in $[-\infty, \infty]$ and $\lim_{\lambda \rightarrow \pm\infty} f(\lambda)/\lambda = \lim_{\lambda \rightarrow \pm\infty} f'(\lambda)$. The function $f = \log Z_s$ is differentiable with $f' = m_s$, and convex because $f'' = m'_s \geq 0$ by the variance formula above. Applying this to $f = \log Z_s$ and combining it with the two limits just computed, we obtain

$$\lim_{\lambda \rightarrow -\infty} m_s(\lambda) = 0, \quad \lim_{\lambda \rightarrow \infty} m_s(\lambda) = s.$$

Recall from Eq (10) that $0 < b(s) < s$, so (by the intermediate value theorem) there exists a unique finite $\lambda(s)$ satisfying

$$m_s(\lambda(s)) = b(s).$$

Since $m_s(\lambda)$ is measurable in s for every fixed λ and continuous and strictly increasing in λ for ν -almost every s , the unique root $\lambda(s)$ admits a measurable version.

The final assertion of the lemma was established at the start of the proof, where $Z_s(\lambda)$ was shown to be positive and finite for every finite λ . ■

B The entropy projection onto the S -marginal and Rogers constraints.

The following lemma describes what happens when the (λ, ν) -update is applied to a probability density q : the resulting density q' satisfies two of the three constraints, namely the S -marginal and the local Rogers conditions, and for any other density p satisfying these two constraints the relative entropies of p and q' with respect to q are related by the Pythagorean identity $H(p|q) = H(p|q') + H(q'|q)$. In particular q' is the density closest to q in relative entropy among those satisfying the two constraints.

Lemma B.1 *Let $D = \{(s, x) : s \geq X_0, 0 < x \leq s\}$, and let q be a probability density on D whose S -marginal q^S is positive ν -almost everywhere. For a given measurable function λ which is finite ν -almost everywhere, define*

$$q_\lambda(x|s) = \frac{e^{\lambda(s)x} q(s, x)}{\int_0^s e^{\lambda(s)y} q(s, y) dy}, \quad x \in (0, s], \text{ for every } s \text{ with } q^S(s) > 0,$$

and suppose that λ is such that

$$\int_0^s x q_\lambda(x|s) dx = b(s) \quad \text{for } \nu\text{-almost every } s. \quad (\text{B-1})$$

Set $q'(s, x) = \nu(s) q_\lambda(x|s)$ for s with $q^S(s) > 0$ and $q'(s, x) = 0$ otherwise, i.e. q' has S -marginal ν and conditional density $q_\lambda(\cdot|s)$ given $S = s$.

Then q' is a probability density on D , and for any probability density p on D with S -marginal ν , conditional mean $\int_0^s x p(x|s) dx = b(s)$ for ν -almost every s , and $H(p|q) < \infty$, we have $H(p|q) = H(p|q') + H(q'|q)$.

Proof. Step 1: properties of q' . For s with $\nu(s) > 0$, integrating the definition of q' over x gives $\int_0^s q'(s, x) dx = \nu(s)$, and for s with $\nu(s) = 0$ both sides vanish; hence q' has S -marginal ν and is a probability density on D . Its conditional density given $S = s$ is $q'(x|s) = e^{\lambda(s)x} q(s, x) / \int_0^s e^{\lambda(s)y} q(s, y) dy$, so the defining equation for λ says precisely that $\int_0^s x q'(x|s) dx = b(s)$ for ν -almost every s : q' has the same conditional mean as p . Moreover, dividing by $q(x|s) = q(s, x)/q^S(s)$,

$$\frac{q'(x|s)}{q(x|s)} = \frac{q^S(s) e^{\lambda(s)x}}{\int_0^s e^{\lambda(s)y} q(s, y) dy} \quad \text{for } \nu\text{-almost every } s, \quad (\text{B-2})$$

whose right-hand side is, for ν -almost every s , positive and finite for every $x \in (0, s)$, since $q^S(s) > 0$, $\lambda(s)$ is finite and $q(s, \cdot)$ is integrable. Hence $q'(\cdot|s)$ and $q(\cdot|s)$ vanish on the same set, i.e. they are mutually absolutely continuous, which we write $q(\cdot|s) \sim q'(\cdot|s)$. Taking logarithms, and noting that neither $q^S(s)$ nor the integral in the denominator depends on x ,

$$\log \frac{q'(x|s)}{q(x|s)} = \lambda(s)x + c(s), \quad c(s) := \log q^S(s) - \log \int_0^s e^{\lambda(s)y} q(s, y) dy, \quad (\text{B-3})$$

so the log-density ratio is an affine function of x .

Step 2: decomposing relative entropy along S . Let P and Q be probability densities on D with $P \ll Q$, and write $P(s, x) = P^S(s)P(x|s)$ and $Q(s, x) = Q^S(s)Q(x|s)$. Then $\log \frac{P(s, x)}{Q(s, x)} = \log \frac{P^S(s)}{Q^S(s)} + \log \frac{P(x|s)}{Q(x|s)}$ for P -almost every (s, x) , and integrating against P gives

$$H(P|Q) = H(P^S|Q^S) + \int P^S(s) H(P(\cdot|s)|Q(\cdot|s)) ds. \quad (\text{B-4})$$

Step 3: the identity fibrewise. Applying Eq (B-4) with $P = p$, $Q = q$ and using $H(p|q) < \infty$, both terms on the right-hand side are finite; in particular $H(p(\cdot|s)|q(\cdot|s)) < \infty$ for ν -almost every s . Fix such an s . Since $q'(\cdot|s)$ is equivalent to $q(\cdot|s)$ by Step 1, Lemma 1.4(b) in [Nutz22] gives

$$H(p(\cdot|s)|q(\cdot|s)) = H(p(\cdot|s)|q'(\cdot|s)) + \int_0^s \log \frac{q'(x|s)}{q(x|s)} p(x|s) dx.$$

By Eq (B-3), for any probability density ρ on $(0, s)$ the integral $\int_0^s \log \frac{q'(x|s)}{q(x|s)} \rho(x) dx = \lambda(s) \int_0^s x \rho(x) dx + c(s)$ depends on ρ only through its mean. Since $p(\cdot|s)$ and $q'(\cdot|s)$ both have mean $b(s)$ (by hypothesis and by Step 1 respectively), the last integral above is unchanged if $p(x|s)$ is replaced by $q'(x|s)$, and equals $H(q'(\cdot|s)|q(\cdot|s))$. Therefore

$$H(p(\cdot|s)|q(\cdot|s)) = H(p(\cdot|s)|q'(\cdot|s)) + H(q'(\cdot|s)|q(\cdot|s)) \quad \text{for } \nu\text{-almost every } s. \quad (\text{B-5})$$

Step 4: integrating over s . Apply Eq (B-4) to the three pairs (p, q) , (p, q') and (q', q) ; the absolute continuity required holds since $p \ll q$ by hypothesis, $q' \ll q$ by Step 1, and $p \ll q'$ because $p(\cdot|s) \ll q(\cdot|s)$ (from $p \ll q$) and $q(\cdot|s) \sim q'(\cdot|s)$ (Step 1, after Eq (B-2)) for ν -almost every s , and both p and q' have S -marginal ν .

Multiplying Eq (B-5) by $\nu(s)$ and integrating over s gives

$$\int \nu(s) H(p(\cdot|s)|q(\cdot|s)) ds = \int \nu(s) H(p(\cdot|s)|q'(\cdot|s)) ds + \int \nu(s) H(q'(\cdot|s)|q(\cdot|s)) ds. \quad (\text{B-6})$$

We now express each of the three integrals in Eq (B-6) via Eq (B-4), whose first term on the right is the relative entropy of the S -marginals. Here $p^S = \nu$ by hypothesis and $q'^S = \nu$ by Step 1, while q^S is the S -marginal of q . Recall (B-4) states that $H(P|Q) = H(P^S|Q^S) + \int P^S(s) H(P(\cdot|s)|Q(\cdot|s)) ds$. Then taking $(P, Q) = (p, q)$, (p, q') and (q', q) respectively gives

$$H(p|q) = H(\nu|q^S) + \int \nu(s) H(p(\cdot|s)|q(\cdot|s)) ds, \quad (\text{B-7})$$

$$H(p|q') = \int \nu(s) H(p(\cdot|s)|q'(\cdot|s)) ds, \quad (\text{B-8})$$

$$H(q'|q) = H(\nu|q^S) + \int \nu(s) H(q'(\cdot|s)|q(\cdot|s)) ds. \quad (\text{B-9})$$

Substituting Eqs (B-7), (B-8) and (B-9) into Eq (B-6), i.e. replacing the three integrals there by $H(p|q) - H(\nu|q^S)$, $H(p|q')$ and $H(q'|q) - H(\nu|q^S)$ respectively, the two terms $H(\nu|q^S)$ cancel and we obtain $H(p|q) = H(p|q') + H(q'|q)$. $\blacksquare$

C Well-definedness of the Sinkhorn scheme

For $n \geq 1$, we regard u_n and v_n as extended-real-valued functions, with $u_n(x) = -\infty$ for $x \notin (0, K)$ and $v_n(s) = -\infty$ for $s \notin (X_0, K)$. The multiplier λ_n is only relevant ν -almost everywhere, and we set $\lambda_n(s) = 0$ whenever $\nu(s) = 0$. Thus $p_{n,n,n}(s, x) = R(s, x) e^{u_n(x) + v_n(s) + \lambda_n(s)x}$ vanishes whenever $x \notin (0, K)$ or $s \notin (X_0, K)$.

Lemma C.1 *Suppose Assumption 2.2 holds and set $u_0 = v_0 = \lambda_0 = 0$. Then every step of the Sinkhorn scheme is well defined. For every $n \geq 0$, $H(p^*|p_{n,n,n}) < \infty$, $H(p^*|p_{n+1,n,n}) < \infty$ and*

$$H(p^*|p_{n,n,n}) - H(p^*|p_{n+1,n,n}) = H(p_{n+1,n,n}|p_{n,n,n}), \quad (\text{C-1})$$

$$H(p^*|p_{n+1,n,n}) - H(p^*|p_{n+1,n+1,n+1}) = H(p_{n+1,n+1,n+1}|p_{n+1,n,n}). \quad (\text{C-2})$$

The functions u_n are finite Lebesgue almost everywhere on $(0, K)$, and λ_n and v_n are finite ν -almost everywhere. Every completed iterate $p_{n,n,n}$ with $n \geq 1$ has S -marginal ν and satisfies the local Rogers condition.

Proof. We proceed by induction. Since $p_{0,0,0} = R$, Assumption 2.2 gives $H(p^*|p_{0,0,0}) < \infty$. Also $u_0 = 0$, $\lambda_0 = 0$ and $v_0 = 0$ are finite, so the finiteness assertions hold at $n = 0$. As the inductive hypothesis, we assume that $H(p^*|p_{n,n,n}) < \infty$, u_n is finite Lebesgue almost everywhere on $(0, K)$, and that λ_n and v_n are finite ν -almost everywhere.

Write $q_n = p_{n,n,n}$ and let q_n^X denote its X -marginal. Since the X -marginals of p^* and q_n are μ and q_n^X , respectively, the DPI gives

$$H(\mu|q_n^X) \leq H(p^*|q_n) < \infty. \quad (\text{C-3})$$

Since $H(\mu|q_n^X) < \infty$, we have $\mu \ll q_n^X$, so the set $\{x \in (0, K) : q_n^X(x) = 0\}$ is μ -null; as $\mu > 0$ Lebesgue almost everywhere on $(0, K)$ by Assumption 2.2, this set is Lebesgue-null, i.e. $q_n^X(x) > 0$ for Lebesgue almost every $x \in (0, K)$. Moreover, since q_n^X is a probability density, $q_n^X(x) < \infty$ for Lebesgue almost every $x \in (0, K)$.

Recall the u -update, Eq (7): $u_{n+1}(x) = \log \mu(x) - \log(\int_x^\infty R(s, x) e^{v_n(s) + \lambda_n(s)x} ds)$. Integrating $q_n(s, x) = R(s, x) e^{u_n(x) + v_n(s) + \lambda_n(s)x}$ over s gives $q_n^X(x) = e^{u_n(x)} \int_x^\infty R(s, x) e^{v_n(s) + \lambda_n(s)x} ds$, so the integral inside the logarithm above equals $e^{-u_n(x)} q_n^X(x)$, and therefore

$$u_{n+1}(x) = \log \mu(x) - \log q_n^X(x) + u_n(x)$$

for Lebesgue almost every $x \in (0, K)$. Hence $p_{n+1,n,n}(s, x)/q_n(s, x) = e^{u_{n+1}(x) - u_n(x)} = \mu(x)/q_n^X(x)$, so $p_{n+1,n,n} \ll q_n$, and $p^* \ll q_n$ since $H(p^*|q_n) < \infty$ by the inductive hypothesis.

Lemma 1.4(b) in [Nutz22] gives

$$H(p^*|q_n) - H(p^*|p_{n+1,n,n}) = \mathbb{E}^{p^*}(\log \frac{p_{n+1,n,n}(S, X)}{q_n(S, X)}) = \mathbb{E}^{p^*}(\log \frac{\mu(X)}{q_n^X(X)}).$$

Since the X -marginal of p^* is μ , $\mathbb{E}^{p^*}(\log \frac{\mu(X)}{q_n^X(X)}) = H(\mu|q_n^X)$, and therefore $H(p^*|q_n) - H(p^*|p_{n+1,n,n}) = H(\mu|q_n^X)$. Since $H(p^*|q_n) < \infty$ and $H(\mu|q_n^X) < \infty$, it follows that $H(p^*|p_{n+1,n,n}) < \infty$.

Moreover, by Eq (13), $H(p_{n+1,n,n}|q_n) = H(\mu|q_n^X)$. Hence

$$H(p^*|q_n) - H(p^*|p_{n+1,n,n}) = H(p_{n+1,n,n}|q_n)$$

which is Eq (C-1). By Eq (13), $H(p_{n+1,n,n}|q_n) = H(\mu|q_n^X)$ and by (C-3) this quantity is finite.

Since $\mu(x) > 0$ and $0 < q_n^X(x) < \infty$ for Lebesgue almost every $x \in (0, K)$ (shown above), and u_n is finite Lebesgue almost everywhere on $(0, K)$ by the inductive hypothesis, the identity $u_{n+1}(x) = \log \mu(x) - \log q_n^X(x) + u_n(x)$ shows that u_{n+1} is finite Lebesgue almost everywhere on $(0, K)$.

We now apply Lemma 2.3 directly to u_{n+1} . Since $p_{n+1,n,n}(s, x) = R(s, x) e^{u_{n+1}(x) + v_n(s) + \lambda_n(s)x}$, its S -marginal is

$$(p_{n+1,n,n})^S(s) = e^{v_n(s)} \int_0^s R(s, x) e^{u_{n+1}(x) + \lambda_n(s)x} dx.$$

Since $p_{n+1,n,n}$ is a probability density, $(p_{n+1,n,n})^S$ is integrable, hence finite for Lebesgue almost every s . As shown above, $H(p^*|p_{n+1,n,n}) < \infty$. Since the S -marginal of p^* is ν , the DPI gives $H(\nu|(p_{n+1,n,n})^S) \leq H(p^*|p_{n+1,n,n}) < \infty$, hence $\nu \ll (p_{n+1,n,n})^S$, i.e. $(p_{n+1,n,n})^S(s) > 0$ for ν -almost every s .

We have already shown that $0 < (p_{n+1,n,n})^S(s) < \infty$ for ν -almost every s . Moreover, v_n is finite ν -almost everywhere by the inductive hypothesis, and hence $0 < e^{-v_n(s)} < \infty$ for ν -almost every s . Therefore

$$0 < \int_0^s R(s, x) e^{u_{n+1}(x) + \lambda_n(s)x} dx < \infty \quad (\text{C-4})$$

for ν -almost every s . Hence the hypotheses of Lemma 2.3 hold, so the λ -update is well defined and there exists a unique finite measurable function $\lambda_{n+1}(s)$ satisfying

$$\frac{\int_0^s x R(s, x) e^{u_{n+1}(x) + \lambda_{n+1}(s)x} dx}{\int_0^s R(s, x) e^{u_{n+1}(x) + \lambda_{n+1}(s)x} dx} = b(s).$$

By Assumption 2.2 and the fact that ν is a probability density, $0 < \nu(s) < \infty$ for ν -almost every s . Now recall Eq (9):

$$v_{n+1}(s) = \log \nu(s) - \log(\int_0^s R(s, x) e^{u_{n+1}(x) + \lambda_{n+1}(s)x} dx).$$

Then $\log \nu(s)$ is finite for ν -almost every s , and the second logarithm in Eq (9) is finite for ν -almost every s by Lemma 2.3 applied with $u = u_{n+1}$ and $\lambda = \lambda_{n+1}(s)$, so the v -update is well defined and v_{n+1} is finite ν -almost everywhere.

We now apply Lemma B.1 with $q = p_{n+1,n,n}$ and $\lambda(s) = \lambda_{n+1}(s) - \lambda_n(s)$. Its hypotheses hold. First, the S -marginal of q is positive ν -almost everywhere (shown above). Second, by definition $q_\lambda(x|s) \propto e^{\lambda(s)x} q(s, x)$; with $\lambda = \lambda_{n+1} - \lambda_n$ and $q = p_{n+1,n,n} = R e^{u_{n+1}(x) + v_n(s) + \lambda_n(s)x}$ this is

$$e^{(\lambda_{n+1}(s) - \lambda_n(s))x} R(s, x) e^{u_{n+1}(x) + v_n(s) + \lambda_n(s)x} = e^{v_n(s)} R(s, x) e^{u_{n+1}(x) + \lambda_{n+1}(s)x},$$

Hence $q_\lambda(x|s) = R(s, x)e^{u_{n+1}(x)+\lambda_{n+1}(s)x} / \int_0^s R(s, y)e^{u_{n+1}(y)+\lambda_{n+1}(s)y}dy$, whose mean $\int_0^s xq_\lambda(x|s)dx$ is the left-hand side of Eq (8), hence equal to $b(s)$. Finally, by the definition of q' in Lemma B.1,

$$q'(s, x) = \nu(s) q_\lambda(x|s) = \frac{\nu(s)}{\int_0^s R(s, y)e^{u_{n+1}(y)+\lambda_{n+1}(s)y}dy} R(s, x)e^{u_{n+1}(x)+\lambda_{n+1}(s)x} = e^{v_{n+1}(s)} R(s, x)e^{u_{n+1}(x)+\lambda_{n+1}(s)x} = p_{n+1, n+1, n+1}$$

where the third equality is Eq (9).

p^* has S -marginal ν , conditional mean b and $H(p^*|p_{n+1, n, n}) < \infty$ (shown above), so Lemma B.1 gives

$$H(p^*|p_{n+1, n, n}) = H(p^*|p_{n+1, n+1, n+1}) + H(p_{n+1, n+1, n+1}|p_{n+1, n, n}) \quad (\text{C-5})$$

which is Eq (C-2). Since the left-hand side is finite and both terms on the right are nonnegative,

$$H(p^*|p_{n+1, n+1, n+1}) < \infty.$$

Finally, $(p_{n+1, n+1, n+1})^S = \nu$ by Lemma B.1, since $p_{n+1, n+1, n+1}$ is the density q' there, and by Eq (8) $\int_0^s xp_{n+1, n+1, n+1}(s, x)dx = b(s)\nu(s) = r(s)$, so the completed iterate satisfies the local Rogers condition. This completes the induction. ■

D Sketch proof of the Rogers' construction

For completeness we sketch the construction in Section 3 of [Rog93] of a continuous martingale with a prescribed joint law of (S_T, X_T) , in the case relevant here where the joint law has a density. It is convenient to work with $Y = S - X$ and to write $\mu(b, y)$ for the joint density of (S_T, Y_T) on $[x_0, \infty) \times (0, \infty)$, so that $\mu(b, y) = \pi(b, b - y)$ in the notation of Theorem 1.1.

Let W be a Brownian motion started at x_0 with running maximum $\bar{W}_t = \max_{u \leq t} W_u$, and let $\{Z_b\}_{b \geq x_0}$ be a family of random variables, independent of W and of each other, with Z_b having density $f(\cdot; b)$ on $(0, \infty)$. Define $\tau = \inf\{t \geq 0 : \bar{W}_t - W_t \geq Z_{\bar{W}_t}\}$. The rate of exclusions below the level y is $\frac{1}{y}$, so the excursions that reach Z_b arrive at rate $\int_0^\infty y^{-1} f(y; b) dy$, and let τ is the time of the first such excursion. Then

$$\begin{aligned} \mathbb{P}(\bar{W}_\tau > b) &= \exp\left[-\int_{x_0}^b \int_0^\infty \frac{1}{y} f(y; s) dy ds\right], \\ \mathbb{P}(\bar{W}_\tau \in db, \bar{W}_\tau - W_\tau \in dy) &= \mathbb{P}(\bar{W}_\tau \geq b) \frac{1}{y} f(y; b) dy db. \end{aligned} \quad (\text{D-1})$$

Now let $\mu(b, y)$ be the target density and $\rho(b) = \int_b^\infty \int_0^\infty \mu(s, u) du ds = \mathbb{P}(S_T > b)$. Matching (D-1) with μ requires $\mathbb{P}(\bar{W}_\tau \geq b) = \rho(b)$ and

$$f(y; b) = \frac{y\mu(b, y)}{\rho(b)} \quad (\text{D-2})$$

This is a legitimate choice only if $f(\cdot; b)$ is a probability density in y for each b , i.e.

$$\int_0^\infty y\mu(b, y) dy = \rho(b) \quad \text{for almost every } b \geq x_0. \quad (\text{D-3})$$

Substituting $\mu(b, y) = \pi(b, b - y)$, the left-hand side of (D-3) is $\int_{-\infty}^b (b - x)\pi(b, x) dx = b\nu(b) - \eta(b)$ and $\rho(b) = \bar{F}_\nu(b)$, so (D-3) is exactly the local Rogers condition (1).