A re-scaled rough Heston model with jumps - weak convergence for the log stock price to the marginal of a Lévy process

Alessandro Bondi* Martin Forde†

26th March 2026

Abstract

We show weak convergence of the marginals for a re-scaled rough Heston model to the marginal of a Normal Inverse Gaussian (NIG) Lévy process. This shows we can obtain such a limit without having to impose that the true Hurst exponent H for the model is $\frac{1}{2}$ as in [AC24],[1] or that $H \searrow -\frac{1}{2}$ as in [AAR25], so the result potentially has increased financial relevance. We also extend to the case when V has jumps, where we show weak convergence of the log stock price marginals to the marginal of an arithmetic Brownian motion evaluated at the inverse hitting time process of a linear combination of the same Brownian motion plus another independent Brownian motion with drift plus an independent Lévy process.

1 Introduction

qRHeston

The Rough Heston stochastic volatility model was introduced in [JR16], and (using $C-$tightness arguments) they show that the model arises naturally as a weak limit of a high-frequency market microstructure model driven by two nearly unstable Hawkes processes with a heavy-tailed kernel. [ALP19],[ER19] and [GK19] show that the characteristic function of the log stock price for Rough Heston-type models can be expressed in terms of the solution to a non-linear Volterra integral equation (VIE) (see also [ER18] and [BLP24],[BPS24], for extensions to jumps in V), which allows for accurate option pricing for $H \ll 1$ (and even $H = 0$) using an Adams scheme to solve the VIE numerically. This avoids Monte Carlo (MC) techniques for which are notoriously inaccurate for $H \ll 1$ (both in terms of bias and sample variance). From the stochastic Fubini theorem, it is well known that $A_t = \int_0^t V_s ds$ satisfies an equation of the form: $A_t = G_0(t) + \int_0^t \kappa(t-s) W_{A_s} ds$ for some Brownian motion W (see e.g. Theorem 3.1 in [JR20]), which is a non-linear VIE for A in terms of the a.s. $(\frac{1}{2} - \varepsilon)-$Hölder continuous function $W_{(\cdot)}$, and is still well defined even if κ is only in $L^1(0,T)$; this also allows us to consider the hyper-rough regime $H \in (-\frac{1}{2}, 0]$. If we discretize this SVE and re-write in terms of the final (discrete-time) increments of A and W, then we can use an independent sequence of Inverse Gaussian random variables to perform an approximate Monte Carlo simulation of A (see Algorithm 1 in [AA25] for details) which is naturally suited to the regimes $H \ll 1$ and $H \in (-\frac{1}{2}, 0]$. The variance process V is $(H - \varepsilon)-$Hölder continuous like fBM (see e.g. Theorem 3.2 in [JR16]) and the spot and VIX smiles exhibit power-law skew for implied volatility in the small-time limit (see [FGS21], [FGS22] for details). See also [ALP19], [Cuch22], [FGS21], [FS21] for further related results on rough Heston-type models.

[BL24] improve accuracy in estimating the tail part of the integral for Fourier inversion (for e.g. call options under the rough Heston model) using a sinh contour which goes outside the usual strip of analyticity for the mgf (but is admissible as long as it avoids poles of the characteristic function), and also discuss refinements to the usual Adams schemes for solving the rough Heston VIE.

[AC24] show functional weak convergence on the Skorokhod space of càdlàg paths on $[0,T]$ endowed with the M_1 topology for a re-scaled standard Markov Heston model with fast mean-reversion and large vol-of-vol (via an H parameter which is not the Hurst exponent) to one of three different models (either Black-Scholes, a Normal Inverse Gaussian or a Normal Lévy model), depending on whether their H parameter is $>$, $=$, or $< -\frac{1}{2}$. [AAR25] obtain a similar result without any ε parameter but instead letting $H \searrow -\frac{1}{2}$ for the so-called *hyper-rough Heston model* (see also Section 5 in [FGS21] and Section 7 in [AJ21] for more on this model), and exploiting Dirac-type behaviour in their Lemma 2.4.

In this note, we fill in the gap between the two extreme cases considered in [AC24] and [AAR25], by showing that a conceptually similar result is obtained for any $H \in (0, \frac{1}{2})$. We later extend the model to allow positive jumps in V^ε, and in this case (using the Laplace transform of the hitting time to a lower barrier for a spectrally positive Lévy process), we show weak convergence of the log stock price marginals to an infinitely divisible law which is described more explicitly in the abstract above.

*Department of AI, Data and Decision Sciences, Luiss University, Rome abondi@luiss.it

†https://martinforde.github.io

[1] The H parameter in [AC24] is not fixed to be $\frac{1}{2}$, but the Hurst exponent for their family of V^ε processes is of course $\frac{1}{2}$ since the models are all standard Heston for $\varepsilon > 0$.

2 Asymptotics for the terminal log stock price for a re-scaled rough Heston model

We work on a probability space $(\Omega, \mathcal{F}, \mathbb{Q})$ throughout with filtration $(\mathcal{F}_t)_{t\geq 0}$ satisfying the usual conditions.

Proposition 2.1 *Consider a re-scaled rough Heston model for a log stock price process X_t^ε:*

$$\begin{aligned} dX_t^\varepsilon &= -\frac{1}{2}V_t^\varepsilon dt + \sqrt{V_t^\varepsilon}dB_t, \\ V_t^\varepsilon &= V_0 + \frac{1}{\varepsilon\Gamma(\alpha)}\int_0^t (t-s)^{\alpha-1}(\lambda(\theta - V_s^\varepsilon)ds + \sigma\sqrt{V_s^\varepsilon}dW_s), \end{aligned} \tag{1}$$

where B and W are two Brownian motions with $dB_t dW_t = \rho dt$ with $\rho \in [-1,0]$, $\alpha \in (\frac{1}{2}, 1)$ and $\lambda, \sigma > 0$. Then (for $t > 0$ fixed) X_t^ε tends weakly to X_t as $\varepsilon \to 0$, where X_t has the marginal law of a Normal Inverse Gaussian Lévy process which does not depend on $H = \alpha - \frac{1}{2}$ or V_0.

Proof. Let $I^\alpha(f)(t) = \frac{1}{\Gamma(\alpha)}\int_0^t (t-s)^{\alpha-1}f(s)ds$ denote the αth-order fractional integral of a function f for $\alpha \in (0,1]$, and (without loss of generality), we assume $X_0^\varepsilon = 0$. Then for $p \in (0,1)$ (which will be sufficient for our purposes when we invoke the [Bill86] weak convergence result below)

$$\mathbb{E}[e^{pX_t^\varepsilon}] = e^{V_0 I^{1-\alpha}\phi_\varepsilon(t) + \frac{1}{\varepsilon}\lambda\theta I^1\phi_\varepsilon(t)} \tag{2}$$

where ϕ_ε is the unique solution of the non-linear Volterra integral equation (VIE):

$$\phi_\varepsilon(t) = \frac{1}{\Gamma(\alpha)}\int_0^t (t-s)^{\alpha-1}(\frac{1}{2}(p^2-p) + \frac{1}{\varepsilon}(\rho p\sigma - \lambda)\phi_\varepsilon(s) + \frac{1}{\varepsilon^2}\frac{1}{2}\sigma^2\phi_\varepsilon(s)^2)ds$$

(see e.g. Section 4 in [ER19] or Section 7 in [ALP19]). Now let $\phi_\varepsilon(t) = \varepsilon\psi(\varepsilon^q t)$. Then

$$\begin{aligned} \varepsilon\psi(\varepsilon^q t) &= \frac{1}{\Gamma(\alpha)}\int_0^t (t-s)^{\alpha-1}(\frac{1}{2}(p^2-p) + (\rho p\sigma - \lambda)\psi(\varepsilon^q s) + \frac{1}{2}\sigma^2\psi(\varepsilon^q s)^2)ds \\ &= \frac{1}{\Gamma(\alpha)}\int_0^{\varepsilon^q t} (t-u\varepsilon^{-q})^{\alpha-1}(\frac{1}{2}(p^2-p) + (\rho p\sigma - \lambda)\psi(u) + \frac{1}{2}\sigma^2\psi(u)^2)du\,\varepsilon^{-q} \\ &= \frac{\varepsilon^{-q(\alpha-1)}}{\Gamma(\alpha)}\int_0^{\varepsilon^q t} (\varepsilon^q t-u)^{\alpha-1}(\frac{1}{2}(p^2-p) + (\rho p\sigma - \lambda)\psi(u) + \frac{1}{2}\sigma^2\psi(u)^2)du\,\varepsilon^{-q} \end{aligned} \tag{3}$$

where we set $\varepsilon^q s = u$ in the second line, so $\varepsilon^q ds = du$. Then setting $\varepsilon^q t \mapsto t$, we see that

$$\varepsilon\psi(t) = \frac{\varepsilon^{-q\alpha}}{\Gamma(\alpha)}\int_0^t (t-u)^{\alpha-1}F(\psi(u))du \tag{4}$$

where $F(w) = \frac{1}{2}(p^2-p) + (\rho p\sigma - \lambda)w + \frac{1}{2}\sigma^2 w^2$. If now we let $q = -\frac{1}{\alpha}$, the VIE (4) is independent of ε, so

$$\phi_\varepsilon(t) = \varepsilon\psi(\frac{t}{\varepsilon^{\frac{1}{\alpha}}}).$$

Hence for every $t > 0$ (from Lemma 4.4 in [FGS21])

$$\lim_{\varepsilon\to 0}\frac{1}{\varepsilon}\phi_\varepsilon(t) = \lim_{t\to\infty}\psi(t) = \psi(\infty) = U_1(p) := \frac{1}{\sigma^2}[\lambda - p\sigma\rho - \sqrt{\lambda^2 - 2\lambda\rho\sigma p + \sigma^2 p(1-p\bar{\rho}^2)}]$$

for $p \in (0,1)$, where $\psi(\infty)$ is the smallest root of F and $\bar{\rho}^2 = 1-\rho^2$, and the convergence to $\psi(\infty)$ is rapid for $\varepsilon \ll 1$ since $\frac{1}{\varepsilon}\phi_\varepsilon(t) = \psi(\frac{t}{\varepsilon^{\frac{1}{\alpha}}})$. More precisely, Lemma 4.5 in [FGS21] can be applied to $-\psi$, which solves the following VIE:

$$-\psi(t) = \frac{1}{\Gamma(\alpha)}\int_0^t (t-u)^{\alpha-1}F_1(-\psi(u))du, \quad \text{with } F_1(w) = \frac{1}{2}(p-p^2) + (\rho p\sigma - \lambda)w - \frac{1}{2}\sigma^2 w^2$$

on $\mathbb{R}_+$, since $F_1(0) > 0$ and F_1 is analytic and decreasing on $\mathbb{R}_+$, the aforementioned result in [FGS21] implies that $-\psi(t)$ converges as $t \to \infty$ to the positive root of F_1, which coincides with $-U_1(p)$. Then for the exponent in (2), we know that $\frac{1}{\varepsilon}\phi_\varepsilon(.) = \psi(\frac{.}{\varepsilon^{\frac{1}{\alpha}}})$ is continuous on $\mathbb{R}_+$ and admits finite limit at ∞, and thus is bounded, so (by the dominated convergence theorem) we see that

$$V_0 I^{1-\alpha}\phi_\varepsilon(t) + \frac{\lambda\theta}{\varepsilon}I^1\phi_\varepsilon(t) = \frac{V_0}{\Gamma(1-\alpha)}\int_0^t (t-s)^{-\alpha}\phi_\varepsilon(s)ds + \lambda\theta\int_0^t \frac{1}{\varepsilon}\phi_\varepsilon(s)ds \to 0 + \lambda\theta U_1(p)t$$

as $\varepsilon \to 0$, and $\lambda\theta U_1(p)t$ is the log mgf of a Normal Inverse Gaussian Lévy process on its effective domain (see e.g. Remark 2.3 in [FJ11]). Then from e.g. the solution to Problem 30.4 on Page 573 in [Bill86] (which is also used in [GK19]), X_t^ε tends weakly to the marginal law of an NIG process. ■

2.1 Adding jumps into the V^ε process

We now generalize the V^ε process to the following:

$$V_t^\varepsilon \;=\; V_0 + \frac{1}{\Gamma(\alpha)}\int_0^t (t-s)^{\alpha-1}(\frac{1}{\varepsilon}\lambda(\theta - V_s^\varepsilon)ds + \frac{1}{\varepsilon}\sigma\sqrt{V_s^\varepsilon}dW_s + \frac{1}{\varepsilon}d\tilde{J}_s^\varepsilon) \tag{5}$$

where $\tilde{J}_t^\varepsilon = \int_0^t\int_{\mathbb{R}_+} x(N^\varepsilon(dx,ds) - V_s^\varepsilon\nu(x)dxds)$ and $N^\varepsilon(dx,dt)$ is a (time-inhomogeneous) Poisson random measure with (random) intensity $V_t^\varepsilon\nu(x)dxdt$; We assume ν only has positive support with $\nu(\{0\}) = 0$ and $\int_{\mathbb{R}_+} e^{px}\nu(x)dx < \infty$ for p in some open interval including zero, so $\tilde{J}^\varepsilon$ has positive-only jumps. We refer to [BPS24] for details, applications and generalizations of (5).

Setting $z = x/\varepsilon$, $dx = \varepsilon dz$, we see that

$$\frac{1}{\varepsilon}\tilde{J}_t^\varepsilon \;=\; \frac{1}{\varepsilon}\int_0^t\int_0^\infty x(N^\varepsilon(dx,ds) - V_s^\varepsilon\nu(x)dxds) \;=\; \int_0^t\int_0^\infty z(N^\varepsilon(\varepsilon dz,ds) - V_s^\varepsilon\nu(\varepsilon z)\varepsilon dzds)$$

and set $V_\varepsilon(p) = \int_0^\infty(e^{pz} - 1 - pz)\varepsilon\nu(\varepsilon z)dz$ so

$$V_\varepsilon(\varepsilon p) \;=\; \int_0^\infty (e^{p\varepsilon z} - 1 - \varepsilon pz)\varepsilon\nu(\varepsilon z)dz \;=\; \int_0^\infty (e^{px} - 1 - px)\nu(x)dx \;=\; V_1(p) \tag{6}$$

Then adapting the chain of equalities in (3) to account for the jumps, we find that

$$\begin{aligned}
\varepsilon\psi(\varepsilon^q t) &= \frac{1}{\Gamma(\alpha)}\int_0^t (t-s)^{\alpha-1}(\frac{1}{2}(p^2-p) + (\rho p\sigma - \lambda)\psi(\varepsilon^q s) + \frac{1}{2}\sigma^2\psi(\varepsilon^q s)^2 + V_\varepsilon(\varepsilon\psi(\varepsilon^q s)))ds \\
&= \frac{1}{\Gamma(\alpha)}\int_0^{\varepsilon^q t} (t - u\varepsilon^{-q})^{\alpha-1}(\frac{1}{2}(p^2-p) + (\rho p\sigma - \lambda)\psi(u) + \frac{1}{2}\sigma^2\psi(u)^2 + V_\varepsilon(\varepsilon\psi(u)))du\,\varepsilon^{-q} \\
&= \frac{\varepsilon^{-q(\alpha-1)}}{\Gamma(\alpha)}\int_0^{\varepsilon^q t} (\varepsilon^q t - u)^{\alpha-1}(\frac{1}{2}(p^2-p) + (\rho p\sigma - \lambda)\psi(u) + \frac{1}{2}\sigma^2\psi(u)^2 + V_1(\psi(u)))du\,\varepsilon^{-q}
\end{aligned}$$

where we set $\varepsilon^q s = u$ in the second line, so $\varepsilon^q ds = du$. Then setting $\varepsilon^q t \mapsto t$, we see that

$$\varepsilon\psi(t) \;=\; \frac{\varepsilon^{-q\alpha}}{\Gamma(\alpha)}\int_0^t (t-u)^{\alpha-1}F(\psi(u))du \tag{7}$$

where $F(w) = \frac{1}{2}(p^2-p) + (\rho p\sigma - \lambda)w + \frac{1}{2}\sigma^2 w^2 + V_1(w)$. If now we let $q = -\frac{1}{\alpha}$, the VIE (7) is independent of ε as before.
$V_1^{(n)}(0) = \int_0^\infty x^n\nu(x)dx < \infty$ for $n \geq 2$, so V_1 is still analytic at zero, so (again from Lemma 4.4 in [FGS21]), $\frac{1}{\varepsilon}\phi_\varepsilon(t)$ tends to $U_J(p)$ which is the smallest root of $F_J(u) = \frac{1}{2}(p^2-p) + (\rho p\sigma - \lambda)u + \frac{1}{2}\sigma^2u^2 + V_1(u)$ (note V_1 and F_J are convex, see e.g. Lemma 2.2.5 in [DZ98]).

We now construct a time-changed arithmetic Brownian motion whose marginal at time t has log mgf $t\lambda\theta U_J(p)$.

2.2 Characterizing the limiting law for X^ε in terms of a Lévy process

Proposition 2.2 *For $t > 0$ fixed, X_t^ε tends weakly to X_t as $\varepsilon \to 0$, where X_t has the marginal law of a time-changed arithmetic Brownian motion with log mgf $\lambda\theta U_J(p)t$.*

Proof. Let Z be a spectrally positive Lévy process with Lévy triple (b, σ^2, ν) with $b = \rho p\sigma - \lambda$ and log mgf $\Lambda(u) = (\rho p\sigma - \lambda)u + \frac{1}{2}\sigma^2u^2 + V_1(u)$ at $t = 1$. Then from the optional stopping theorem

$$\mathbb{E}(e^{-qH_a}) \;=\; e^{-\Phi(q)a} \;=\; e^{\Phi(q)|a|} \tag{8}$$

for $q \geq 0$ and $a < 0$ where $H_a = \inf\{t \geq 0 : Z_t < a\}$ and $\Phi(q) \leq 0$ is smallest root of $\Lambda(u) = q$ (since we have a *lower* barrier here), see e.g. Eq 2.5 in [KKR13], Theorem 46.3 in [Sato99] or Appendix C in [BF25] [2], and (8) also holds for $q < 0$ if q is greater than the minimum value attained by $\Lambda(.)$.

If we now let $W_t^{(-\frac{1}{2})} = W_t^\perp - \frac{1}{2}t$ where $W^\perp$ is another Brownian motion independent of Z and everything else, then for p in some non-zero interval around zero, we see that

$$\mathbb{E}(e^{pW_{H_{-t}}^{(-\frac{1}{2})}}) \;=\; \mathbb{E}(e^{\frac{1}{2}(p^2-p)H_{-t}}) \;=\; e^{t\Phi_p(-\frac{1}{2}(p^2-p))} \tag{9}$$

[2] in [KKR13] they take the largest root since they are considering an upper barrier

and $\Phi_p(-\frac{1}{2}(p^2-p))$ is $U_J(p)$ at the end of the previous subsection (note that H_{-t} also depends on p so we cannot say this is the mgf of $W_{H_{-t}}^{(-\frac{1}{2})}$).

It turns out that (9) is the mgf of $\eta_{\tau_t^-}$, where $\eta_t = W_t - \frac{1}{2}t$ and $\tau_t^- = \inf\{s : \xi_s < -t\}$, where $\xi_t = -\lambda t + \sigma(\rho W_t + \bar{\rho}\tilde{W}_t) + J_t$, and $\tilde{W}$ is another Brownian motion independent of W and J is a spectrally positive Lévy process (independent of $(W, \tilde{W})$), with $\log \mathbb{E}(e^{uJ_1}) = V_1(u)$. To see this, we first note that

$$f_p(u) \quad := \quad \log \mathbb{E}(e^{u\xi_1 + p\eta_1}) \quad = \quad \Lambda_p(u) + \frac{1}{2}(p^2 - p)$$

with $\Lambda_p(u) = -\lambda u + \frac{1}{2}\sigma^2 u^2 + V_1(u) + \rho\sigma p u$, since $\mathrm{Cov}(\xi_1, \eta_1) = \rho\sigma$. Now define $\kappa(p)$ to be the root u of $\Lambda_p(u) + \frac{1}{2}(p^2 - p) = 0$ for which $\kappa(0) = 0$. Note $\Lambda_p(.)$ is finite and convex and $\Lambda_p(0) = 0$ and $-\frac{1}{2}(p^2 - p) > 0$ for $p \in (0, 1)$, so two roots exist for $p \in (0, 1)$, and since $\mathbb{E}(\xi_1) = \Lambda_0'(0) = -\lambda < 0$ where $\Lambda_0(u) = -\lambda u + \frac{1}{2}\sigma^2 u^2 + V_1(u)$ and $V_1'(0) = 0$, $\tau_t^- < \infty$ a.s. and the left root is the correct one when $p \in (0, 1)$. Then $\kappa(p) = \Phi_p(-\frac{1}{2}(p^2 - p))$ (as defined above), and from the optional stopping theorem

$$\mathbb{E}(e^{\kappa(p)\xi_{\tau_t^-} + p\eta_{\tau_t^-}} 1_{\tau_t^- \le n}) + \mathbb{E}(e^{\kappa(p)\xi_n + p\eta_n} 1_{\tau_t^- > n}) \quad = \quad 1\,.$$

By the monotone convergence theorem, the first term tends to $\mathbb{E}(e^{\kappa(p)\xi_{\tau_t^-} + p\eta_{\tau_t^-}}) = \mathbb{E}(e^{-t\kappa(p) + p\eta_{\tau_t^-}})$ as $n \to \infty$ because ξ has no negative jumps, and $\lim_{n\to\infty} 1_{\tau_t^- \le n} = 1_{\tau_t^- < \infty} = 1$.

To deal with the second term, we introduce the Esscher tilt $\frac{d\mathbb{Q}^{(p)}}{d\mathbb{P}}|_{\mathcal{F}_t} = e^{\kappa(p)\xi_t + p\eta_t}$ and note that $\mathbb{E}(e^{\kappa(p)\xi_t + p\eta_t}) = 1$. Then

$$\mathbb{E}(e^{\kappa(p)\xi_n + p\eta_n} 1_{\tau_t^- > n}) \quad = \quad \mathbb{Q}^{(p)}\{\tau_t^- > n\}$$

and

$$\mathbb{E}^{\mathbb{Q}^{(p)}}(e^{u\xi_t}) \quad = \quad e^{(\Lambda_p(u+\kappa(p)) + \frac{1}{2}(p^2-p))t} \quad = \quad e^{t(\Lambda_p(u+\kappa(p)) - \Lambda_p(\kappa(p)))}$$

and $\frac{\partial}{\partial u}(\Lambda_p(u + \kappa(p)) - \Lambda_p(\kappa(p)))|_{u=0} = \Lambda_p'(\kappa(p))$. We need to show that $\Lambda_p'(\kappa(p)) = f_p'(\kappa(p)) < 0$.

For $p \in (0, 1)$, $f_p(0) = \frac{1}{2}(p^2 - p) < 0$, and by definition of $\kappa(p)$, $f_p(\kappa(p)) = 0$. Since $\kappa(p)$ is the smaller root, necessarily $\kappa(p) < 0$. Because f_p is convex, f_p' is increasing. Suppose for contradiction that $f_p'(\kappa(p)) \ge 0$. Then for every $u \in (\kappa(p), 0)$,

$$f_p'(u) \quad \ge \quad f_p'(\kappa(p)) \quad \ge \quad 0.$$

Hence f_p is non-decreasing on $(\kappa(p), 0)$, so $f_p(0) \ge f_p(\kappa(p)) = 0$, which contradicts $f_p(0) = \frac{1}{2}(p^2 - p) < 0$. Therefore

$$f_p'(\kappa(p)) < 0.$$

Since $f_p'(u) = \Lambda_p'(u)$, this yields that $\Lambda_p'(\kappa(p)) < 0$. Hence $\xi_t \to -\infty$ as $t \to \infty$ $\mathbb{Q}^{(p)}$-a.s., so $\mathbb{Q}^{(p)}\{\tau_t^- > n\} \to \mathbb{Q}^{(p)}\{\tau_t^- = \infty\} = 0$. Thus we have shown that

$$\mathbb{E}(e^{p\eta_{\tau_t^-}}) \quad = \quad e^{\kappa(p)t}$$

for $p \in (0, 1)$, and (as above) the result follows from the solution to Problem 30.4 on Page 573 in [Bill86]. Thus X_t^ε tends weakly to the law of $\eta_{\tau_{\lambda\theta t}^-}$. ■

References

[AJ21] Abi Jaber, E., "Weak existence and uniqueness for affine stochastic Volterra equations with L^1-kernels", *Bernoulli, 27*(3), 1583-1615, 2021.

[AA25] Abi Jaber, E. and E.Attal, "Simulating integrated Volterra square-root processes and Volterra Heston models via Inverse Gaussian", preprint, 2025.

[AAR25] Abi Jaber, E., E.Attal and M.Rosenbaum, "From Hyper Roughness to Jumps as $H \to -\frac{1}{2}$", Preprint, 2025.

[ACLP21] Abi Jaber, E., C.Cuchiero, M.Larsson and S.Pulido, "A weak solution theory for stochastic Volterra equations of convolution type", *Annals of Applied Probability, 31*(6), 2924-2952, 2021.

[AC24] Abi Jaber, E. and N.De Carvalho, "Reconciling rough volatility with jumps", *SIAM Journal on Financial Mathematics, 15*(3), 785-823, 2024.



Figure 1: Here we see convergence of the VIE $\psi(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} F(\psi(s))ds$ to the asymptotic solution $U_J(p)$ for $H = .2$, $\sigma = 1$; $\lambda = 1$, $\rho = -.65$, $p = \frac{1}{2}$ and $\nu(x) = \frac{e^{-Mx}}{x^{1+Y}} 1_{x>0}$ with $M = 2$ and $Y = 1.5$ using an Adams scheme with 2000 time steps.



Figure 2: The log mgf $U_J(p)$ of $\eta_{\tau_t^-}$ at $t = 1$ for the same parameters in the plot above. $U_J(p) = \infty$ for p outside the domain shown in the plot.



Figure 3: Here we have plotted the implied volatility smile as a function of log-moneyness $x = \log \frac{S_0}{K}$ for the asymptotic model $S_t = e^{\eta_{\tau_{\lambda\theta t}^-}}$ in Proposition 2.2 for the same parameters as above but with $C_1 = 1$ (blue), $C_1 = .25$ (yellow) and $C_1 = .01$ (green) and $\rho = -.5$, $\lambda = 1$, $\theta = .04$ and maturity $T = 1$, using the usual Lewis Fourier inversion formula in Eq 1.6 in [GR24]. Note the asymptotic smile does not depend on V_0 or H.

[ALP19] Abi Jaber, E., M.Larsson and S.Pulido, "Affine Volterra processes", *Annals of Applied Probability, 29*(5), 3155-3200, 2019.

[Bill86] Billingsley, P., "*Probability and measure*", John Wiley&Sons, 2nd edition, 1986.

[BLP24] Bondi, A., G.Livieri and S.Pulido, "Affine Volterra processes with jumps", *Stochastic Processes and their Applications, 168*, 1042-64, 2024.

[BF25] Bondi, A., and M.Forde, "Lévy processes as weak limits of rough Heston models", preprint 2025

[BPS24] Bondi, A., S. Pulido and S.Scotti, "The rough Hawkes Heston stochastic volatility model", *Mathematical Finance, 34*(4), 1197-1241, 2024.

[BL24] Boyarchenko, S. and S.Levendorskii, "Correct implied volatility shapes and reliable pricing in the rough Heston model", Preprint, 2024.

[Cuch22] Cuchiero, C., "Modelling rough covariance processes", Talk, February 2022.

[DZ98] Dembo, A. and O.Zeitouni, "*Large deviations techniques and applications*", Jones and Bartlett publishers, Boston, 1998.

[ER18] El Euch, O. and M.Rosenbaum, "Perfect hedging in rough Heston models", *Annals of Applied Probability, 28*(6), 3813-3856, 2018.

[ER19] El Euch, O. and M.Rosenbaum, "The characteristic function of Rough Heston models", *Mathematical Finance, 29*(1), 3-38, 2019.

[FGS21] Forde, M., S. Gerhold and B.Smith, "Small-time, large-time and $H \to 0$ asymptotics for the rough Heston model", *Mathematical Finance, 31*(1), 203-241, 2021.

[FGS22] Forde, M., S. Gerhold and B.Smith, "Small-time VIX smile and the stationary distribution for the Rough Heston model", Working paper, 2022.

[FJ11] Forde, M. and A.Jacquier, "The large-maturity smile for the Heston model", *Finance and Stochastics, 15*(4), 755-780, 2011.

[FS21] Forde, M and B.Smith, "Rough Heston with jumps - joint calibration to SPX/VIX level and skew as $T \to 0$, and issues with the quadratic rough Heston model", preprint, 2021.

[GK19] Gatheral, J. and M.Keller-Ressel, "Affine forward variance models", *Finance and Stochastics, 23*, 501-533, 2019.

[GR24] Gatheral, J. and Radoicic, "A generalization of the rational rough Heston approximation", preprint, 2024

[GGP19] Gerhold, S., C.Gerstenecker and A.Pinter, "Moment Explosions In The Rough Heston Model", *Decisions in Economics and Finance, 42*(2), 575-608, 2019.

[Jai15] Jaisson, T., "Market impact as anticipation of the order flow imbalance", *Quantitative Finance, 15*(7), 1123-1135, 2015.

[JR16] Jaisson, T. and M.Rosenbaum, "Rough fractional diffusions as scaling limits of nearly unstable heavy tailed Hawkes processes", *Annals of Applied Probability, 26*(5), 2860-2882, 2016.

[JR20] Jusselin, P. and M.Rosenbaum, "No-arbitrage implies power-law market impact and rough volatility", *Mathematical Finance, 30*(4), 1309-1336, 2020.

[KKR13] Kuznetsov, A., A.Kyprianou and V.Rivero, "The theory of scale functions for spectrally negative Lévy processes", Review article, 2013.

[Lew00] Lewis, A.L., "Option valuation under stochastic volatility", Finance Press, 2000.

[MORS26] Muhle-Karbe, J., Y.Chahdi, M.Rosenbaum and G.Szymanski, "A unified theory of order flow, market impact, and volatility", preprint.

[Sato99] Sato, K., "Lévy Processes and Infinitely Divisible Distributions", Cambridge University Press, 1999.