


Patients in staircase periods (n=1800)

Not sampled for nested staircase outcomes (n=1000):
. Outside randomised shifts (n=920)
. Other (n=80)

Sampled for nested staircase outcomes (n=800)

Sequence 1



Lost/excluded (n=14; reasons):
. Lost to follow-up (n=8)
. Withdrew consent (n=3)
. Other (n=3)

Analysed for nested staircase outcomes (n=131)

Sequence 2



Lost/excluded (n=12; reasons):
. Lost to follow-up (n=8)
. Withdrew consent (n=2)
. Other (n=2)

Analysed for nested staircase outcomes (n=140)


Sequence 3



Lost/excluded (n=13; reasons):
. Lost to follow-up (n=8)
. Withdrew consent (n=3)
. Other (n=2)

Analysed for nested staircase outcomes (n=147)


Sequence 4



Lost/excluded (n=14; reasons):
. Lost to follow-up (n=9)
. Withdrew consent (n=2)
. Other (n=3)

Analysed for nested staircase outcomes (n=154)

Sequence 5



Lost/excluded (n=15; reasons):
. Lost to follow-up (n=10)
. Withdrew consent (n=3)
. Other (n=2)

Analysed for nested staircase outcomes (n=160)

Before ATLS training
In staircase periods (n=925)
Sampled (n=411)
Analysed (n=376)
Excluded (n=35; reasons):
. Lost to follow-up (n=22)
. Withdrew consent (n=7)
. Died before follow-up (n=2)
. Other (n=4)

After ATLS training
In staircase periods (n=875)
Sampled (n=389)
Analysed (n=356)
Excluded (n=33; reasons):
. Lost to follow-up (n=21)
. Withdrew consent (n=6)
. Died before follow-up (n=3)
. Other (n=3)

Analysed for nested staircase outcomes (n=732)

Excluded from nested staircase analyses (n=68):
. Lost to follow-up (n=43)
. Withdrew consent (n=13)
. Died before follow-up (n=5)
. Other (n=7)

Staircase: standard care
Staircase: intervention

Transition
Outside staircase