12. Comorbidities (Charlson Comorbidity Index) Myocardial infarction - History of definite or probable MI (EKG changes and/or enzyme changes) Congestive heart failure - Exertional or paroxysmal nocturnal dyspnea and has responded to digitalis, diuretics, or afterload reducing agents Peripheral vascular disease - Intermittent claudication or past bypass for chronic arterial insufficiency, history of gangrene or acute arterial insufficiency, or untreated thoracic or abdominal aneurysm (≥6 cm) Cerebrovascular disease - History of a cerebrovascular accident with minor or no residua and transient ischemic attacks Dementia - Chronic cognitive deficit Chronic pulmonary disease - Asthma, chronic bronchitis, emphysema, and other lung disease who have ongoing symptoms such as dyspnea or cough, with mild or moderate activity. Peptic ulcer disease - Any history of treatment for ulcer disease or history of ulcer bleeding Liver disease - Severe = cirrhosis and portal hypertension with variceal bleeding history, moderate = cirrhosis and portal hypertension but no variceal bleeding history, mild = chronic hepatitis (or cirrhosis without portal hypertension) Renal disease - Severe = on dialysis, status post kidney transplant, uremia, moderate = creatinine >3 mg/dL (0.27 mmol/L)	(Source: Medical record, treating physician or interview)
12.1 Please specify	_____
13. Severity of liver disease	(Source: Medical record, treating physician or interview)
14. Severity of diabetes	(Source: Medical record, treating physician or interview)
15. Severity of malignancy	(Source: Medical record, treating physician or interview)
Any other comments?	_____