

BIOAVAILABILITY AMPLIFIERS: DIETARY FATS AND Neu5Gc DELIVERY

The Role of Animal-Derived Fats as Structural Carriers of Xeno-Antigenic Load in the Hamer Loop

Martin Hamer, RN (Ret.)

Hamer Glycan Systems | Project: Acnerase | 2026

Confidentiality Notice: This document forms part of the Hamer Loop Manuscript Series (Documents 1–7), authored by Martin Hamer, RN (Ret.), Hamer Glycan Systems, 2026. All content is the original intellectual property of the author and is protected under Canadian copyright law. Unauthorised reproduction or distribution without express written consent is strictly prohibited.

Abstract

This document, the third in the Hamer Loop Manuscript Series, addresses the role of animal-derived dietary fats as structural amplifiers of Neu5Gc-mediated inflammation. Within the Hamer Loop framework, not all sources of N-glycolylneuraminic acid (Neu5Gc) produce equivalent inflammatory responses. The severity of an immune reaction depends substantially on how efficiently the antigen is delivered into the systemic circulation and, critically, into the lymphatic system — the compartment in which the Hamer Loop generates its proposed inflammatory reservoir.

Three intersecting mechanisms are proposed to explain why concentrated animal fats produce more severe, persistent, and anatomically deeper lesions than lower-fat dairy sources: elevated antigen concentration within fat-associated glycan structures, particularly the Milk Fat Globule Membrane (MFGM); chylomicron-mediated lymphatic delivery that bypasses hepatic first-pass metabolism and deposits the antigen directly into the compromised clearance system; and enhanced interstitial sequestration resulting from increased lymphatic viscosity under high lipid load.

Two negative controls of particular evidential value are examined: porcine fat (lard), which does not trigger the Hamer Loop despite being a structurally similar rendered animal fat, confirming that the operative trigger is bovine-specific Neu5Gc rather than animal fat as a class; and farmed Atlantic salmon, which despite being fed bovine-derived rations does not trigger the Hamer Loop, consistent with secondary metabolic processing neutralising the bovine antigenic signature.

Keywords: *Neu5Gc; dietary fat; chylomicron; lymphatic delivery; Milk Fat Globule Membrane; tallow; lard; bovine specificity; xenosialitis; interstitial sequestration; Hamer Loop*

1. Overview and Position Within the Series

Document 2 of this series established that lesion severity within the Hamer Loop framework is determined by two principal variables: the Neu5Gc antigenic load delivered to the immune system, and the presence or absence of ammonium-based catalysts that potentiate the immune response beyond what the antigen load alone would generate. Fat was identified as the primary structural determinant of antigen load: volume of antigen delivery is a fat function.

This document examines precisely how dietary fat delivers that antigenic load. The answer lies not merely in the quantity of Neu5Gc present in fat-rich bovine foods, but in the specific absorption and transport pathway that concentrated fats follow — a pathway that bypasses the liver entirely and deposits antigen directly into the lymphatic system, the very compartment in which the Hamer Loop produces its inflammatory reservoir.

Understanding the fat delivery mechanism is essential for two practical purposes: it explains why the severity hierarchy in the Neu5Gc Severity Index (Appendix II) correlates with fat concentration rather than Neu5Gc content alone, and it informs the clinical advice that complete avoidance of concentrated dairy fats and rendered bovine fats is more impactful than avoidance of lean red meat in reducing Hamer Loop activity.

2. Antigen Concentration: Glycan Density in Animal Fats

Neu5Gc is not uniformly distributed across the constituent fractions of animal-derived foods. In bovine sources, the antigen is disproportionately concentrated in fat-associated glycan structures — most notably the Milk Fat Globule Membrane (MFGM), the phospholipid bilayer surrounding each fat globule in bovine milk.

2.1 The Milk Fat Globule Membrane as an Antigen Concentrator

The MFGM is a glycoprotein-rich trilayer membrane that encases each fat droplet in bovine milk. It is disproportionately rich in sialic acid-bearing glycoproteins, including MUC1, butyrophilin, and lactadherin — all of which carry Neu5Gc in bovine species. When milk is processed into concentrated dairy products, the progressive reduction of the aqueous fraction concentrates the MFGM-associated glycan structures.

Butter, for instance, retains the MFGM at high density relative to liquid milk, while ghee — produced by cooking butter to remove the aqueous phase — concentrates the fat-soluble Neu5Gc fractions further still. Anhydrous Milk Fat (AMF), with greater than 99.8% fat content, represents the most extreme concentration of this process. The result is a product in which Neu5Gc-bearing glycan structures are packed at densities many times greater than in the liquid milk from which they originate.

2.2 Rendered Bovine Fat: Tallow

Tallow, derived from bovine adipose tissue, delivers a higher antigenic load per gram than lean bovine muscle meat, in which Neu5Gc is primarily associated with the cellular protein fraction

rather than the lipid fraction. This distinction has practical implications for risk stratification: rendered bovine fats are not interchangeable with lean meat cuts in terms of their potential to trigger Hamer Loop activity.

Tallow appears as a processing ingredient in a range of food products — commercial frying oils, confectionery glazes, processed meat binders, and cosmetic formulations — where its bovine origin is not immediately apparent from the label. This makes it a meaningful and frequently overlooked target for the Acnerase detection framework.

2.3 Porcine Fat (Lard): A Critical Negative Control

Lard, derived from porcine adipose tissue, is structurally comparable to bovine tallow as a rendered animal fat. Despite this structural similarity, lard has been confirmed as a consistent negative control within this framework across more than four decades of longitudinal self-provocation trials: no inflammatory cutaneous response has been observed following lard consumption.

This finding is significant and must be stated clearly. It demonstrates that the fat delivery mechanism described in this document is insufficient on its own to trigger the Hamer Loop. The bovine-specific antigenic component remains the necessary condition. Porcine fat, despite following the same chylomicron-mediated absorption route as bovine tallow, does not carry the relevant xeno-antigenic load in this subject. This antigen specificity within the rendered fat category reinforces the central claim of the series: the immune response is directed at bovine-derived Neu5Gc specifically — in its unmodified glycan configuration — not at animal fat as a class, not at rendered fat as a category, and not at Neu5Gc broadly across all mammalian species.

3. Chylomicron-Mediated Lymphatic Delivery: The Route That Matters

The delivery pathway of the antigen is proposed to be as consequential as the antigenic load itself. This is the mechanistic core of Document 3 and the primary explanation for why fat-associated Neu5Gc sources produce disproportionately severe Hamer Loop responses.

3.1 The Two Absorption Routes

Water-soluble compounds absorbed from the intestinal lumen enter the portal venous system and pass through the liver before reaching systemic circulation. The liver performs first-pass metabolism on a wide range of absorbed compounds, potentially modifying or attenuating antigenic molecules before they reach the lymphatic and vascular compartments where Hamer Loop activity occurs.

Dietary fats follow an entirely different route. In intestinal enterocytes, absorbed fatty acids and fat-soluble compounds are packaged into chylomicrons — large lipoprotein particles — which are absorbed not into the portal blood supply but into the lacteals, the lymphatic capillaries of the small intestinal villi. Chylomicrons travel through the intestinal lymphatics and enter the

systemic lymphatic circulation via the thoracic duct, bypassing hepatic first-pass metabolism entirely.

3.2 Why This Route Is Mechanistically Critical

The lymphatic system is precisely the compartment in which Hamer Loop activity produces its proposed inflammatory reservoir. Regional lymphatic flow becomes functionally limited during active inflammatory cycles, inflammatory compounds are not efficiently cleared, and interstitial sequestration occurs. Direct lymphatic entry of Neu5Gc-bearing chylomicrons therefore deposits the antigen into an already compromised clearance system — at exactly the site where sequestration is most likely to occur and persist.

This mechanism provides a coherent explanation for one of the most reproducible clinical observations in this framework: high-fat triggers such as aged cheese, butter, or tallow consistently produce more persistent, deeper, and slower-resolving lesions than liquid milk containing equivalent Neu5Gc by mass. The antigen is not simply more concentrated — it is delivered via a route that circumvents the body's primary attenuation mechanism and deposits it directly into the compromised clearance system.

3.3 The Farmed Salmon Observation: Secondary Metabolic Processing as a Neutralising Mechanism

Farmed Atlantic salmon is commercially fed rations containing rendered bovine products, including blood-based pellets. Despite the bovine-derived content of the feed — and despite salmon being a fatty fish whose lipids would be absorbed via the chylomicron-mediated lymphatic route — consumption of farmed salmon has not triggered an inflammatory response across longitudinal self-provocation trials spanning several decades.

This observation is not easily explained by a simple fat-content or Neu5Gc-load model. If the bovine-derived antigenic load were passing through the salmon and remaining biologically active in its original configuration, a Hamer Loop response would be expected — particularly given the lymphatic delivery route that salmon fat would follow.

The proposed explanation is that secondary metabolic processing by a non-mammalian intermediate host — the salmon — sufficiently alters the bovine glycan structures such that the resulting Neu5Gc is no longer presented in the molecular configuration recognised by the human anti-Neu5Gc antibody response. The glycan scaffolding surrounding the Neu5Gc molecule, which appears to determine its immunogenic potential in this framework, is metabolically transformed during incorporation into salmon tissue, neutralising the bovine antigenic signature.

This finding — taken alongside the porcine lard negative control — points consistently toward a specific molecular configuration of bovine Neu5Gc as the operative trigger, rather than Neu5Gc broadly, animal fat generally, or rendered fat as a category. It also suggests that the antigenic potential of Neu5Gc may be modifiable — a finding with potential implications for food processing and for the development of non-reactive bovine-derived food ingredients.

4. Lymphatic Viscosity and Impaired Clearance

A third mechanism compounds the effects of elevated antigen concentration and direct lymphatic delivery: the physical properties of lipid-rich lymph may further impede clearance of inflammatory mediators in individuals already experiencing Hamer Loop-associated lymphatic functional limitation.

4.1 The Viscosity Hypothesis

High dietary fat intake is associated with transient increases in the lipid content of lymph fluid — a well-established physiological response to fat absorption via the lacteals. In individuals in whom regional lymphatic flow is already functionally limited by active Hamer Loop cycling, the introduction of a high-lipid antigenic load may further impede clearance kinetics through a viscosity-mediated mechanism: elevated lipid content increases lymph viscosity, slowing transit through already-compromised lymphatic vessels and prolonging the dwell time of inflammatory mediators within the interstitial reservoir.

4.2 Consequence for Lesion Characteristics

The combined effect of higher antigenic density (Section 2), direct lymphatic delivery (Section 3), and impaired clearance (Section 4.1) produces a more substantial and longer-lasting inflammatory reservoir than lower-fat dairy exposures. This is consistent with the observation that fat-associated triggers produce lesions that are physically larger, anatomically deeper, and slower to resolve — characteristics that distinguish them qualitatively, not merely quantitatively, from whole-milk-associated lesions.

Within the three-phase lesion classification established in Document 2, fat-dominant exposures are the primary driver of Category 2 (later-phase, sebaceous-microbial transition) and Category 3 (deep cystic, chronic lymphatic sequestration) presentations. The fat delivery mechanisms described in this document provide the underlying explanation for why these deeper lesion categories are associated specifically with sustained and high-fat bovine trigger exposures.

5. Comparative Severity Framework

The following table summarises the proposed relationship between antigen source, delivery mechanism, and observed lesion characteristics across the full range of trigger categories in this framework. It is derived from longitudinal observational data and forms the basis of the severity hierarchy in the Neu5Gc Severity Index (Appendix II).

Trigger Category	Antigen Density	Delivery Route	Amplifiers Present	Typical Lesion Outcome
Whole-food dairy (milk, soft cheese, yoghurt)	Low to moderate	Portal absorption; hepatic first-pass	None	Category 1; small, transient; rapid resolution

Trigger Category	Antigen Density	Delivery Route	Amplifiers Present	Typical Lesion Outcome
Lean red meat (bovine)	Moderate	Portal absorption; standard protein kinetics	None	Category 1 to 2; moderate; resolves within days
Concentrated bovine fats (butter, ghee, tallow, AMF, aged cheese)	High	Chylomicron-mediated lymphatic delivery; bypasses liver	None (fat effect only)	Category 2 to 3; persistent; deeper; slower resolution
Ammonium-potentiated products (any Neu5Gc source)	Variable	Variable; threshold lowered chemically	Ammonium catalyst; NLRP3 priming	Category 2 to 3; deep nodular; prolonged; most severe
Porcine fat (lard)	Present but non-reactive	Chylomicron-mediated (same route as tallow)	None	No response — confirmed negative control
Farmed salmon	Bovine-derived in feed; processed by intermediate host	Chylomicron-mediated	None	No response — bovine antigen neutralised by secondary processing

Table 1. Comparative Severity Framework for Neu5Gc Trigger Categories. The inclusion of negative controls (porcine fat, farmed salmon) is critical: it confirms that the chylomicron delivery route alone is insufficient to trigger the Hamer Loop, and that bovine-specific Neu5Gc in its unmodified configuration is the necessary condition.

6. Practical Implications for Avoidance Strategy

The fat delivery mechanism described in this document has direct implications for the design of elimination protocols and for the clinical advice provided to susceptible individuals.

6.1 Prioritising Elimination of High-Fat Bovine Sources

Complete avoidance of all Neu5Gc sources is necessary to achieve full symptom resolution. However, the severity hierarchy indicates that concentrated dairy fats and rendered bovine fats produce disproportionately severe and persistent responses relative to whole-food sources. For individuals beginning an elimination protocol, prioritising the removal of butter, ghee, AMF, aged and processed cheeses, tallow, and processed meat products containing bovine-derived binders is likely to produce the greatest and most rapid reduction in lesion burden.

6.2 Hidden Sources of Concentrated Bovine Fat

Concentrated bovine fat appears in processed foods in quantities and under label terms that are not intuitively apparent from label reading. The following categories represent the most significant hidden vectors:

- Anhydrous Milk Fat (AMF): appears in confectionery, bakery products, recombined milk powders, and commercial chocolate
- Tallow: appears in commercial frying oils, processed meat binders, cosmetic formulations, and some margarines
- Dairy-derived anti-caking agents: found in powdered foods, spice blends, and pharmaceutical tablet fillers
- Bovine gelatin: present in pharmaceutical capsule shells, confectionery (marshmallows, gummy products), and dessert setting agents — a concentrated collagen-derived Neu5Gc source

The Acnerase application's ingredient detection logic is specifically designed to surface these hidden concentrated fat sources, which represent the upper tier of the Neu5Gc Severity Index and the primary drivers of Category 2 and Category 3 lesion development.

6.3 Re-Exposure Sensitivity Following Elimination

As documented in Section 6.5 of Document 1, complete clearance of systemic trigger load — achieved after approximately 7–10 days of full elimination — produces a marked amplification of re-exposure sensitivity. This phenomenon applies with particular force to concentrated fat sources. An individual who has successfully cleared their system and then inadvertently consumes a product containing AMF, aged cheese, or tallow should expect a response that is more severe than the same exposure would have produced in the chronically loaded state. This is not a sign of worsening disease — it is the immune system operating at full sensitivity in the absence of the chronic tolerance that had been masking its reactivity.

7. Limitations

The mechanisms described in this document are proposed on the basis of longitudinal self-provocation data and known lipid absorption physiology. The specific claim that Neu5Gc is preferentially concentrated in the MFGM fraction of processed dairy fats has support in the glycobiology literature (Samraj et al., 2015; Tangvoranuntakul et al., 2003), but the quantitative relationship between fat concentration and antigenic load in specific processed products has not been directly measured within this study.

The lymphatic viscosity hypothesis is mechanistically plausible but has not been formally tested. No direct measurement of lymph lipid content or lymphatic transit velocity has been performed in association with Hamer Loop provocation events. This hypothesis is offered as a mechanistically coherent explanation for a consistent observational pattern, not as an established physiological finding.

The farmed salmon observation, while compelling in its consistency across decades of provocation trials, is a single-subject finding. The proposed mechanism — secondary metabolic processing neutralising the bovine antigenic signature — is consistent with the broader bovine-specificity evidence but has not been tested by direct analysis of Neu5Gc glycan configuration in salmon tissue from bovine-fed aquaculture sources.

The comparative severity table reflects observational patterns rather than controlled dose-response data. Individual variability in anti-Neu5Gc antibody titre, gut microbiome composition, TRM cell distribution, and intestinal permeability status may substantially influence the severity gradient for any given individual. The hierarchy is a population-level framework derived from one subject's longitudinal record.

8. Conclusion

The central insight of this document is that the route of antigen delivery matters as much as the quantity of antigen delivered. Chylomicron-mediated lymphatic absorption — the pathway followed by concentrated dietary fats — bypasses the hepatic first-pass attenuation that attenuates water-soluble antigen loads and deposits Neu5Gc-bearing glycan structures directly into the compromised clearance system where the Hamer Loop generates its inflammatory reservoir.

The bovine-specificity of this effect — confirmed by the porcine fat and farmed salmon negative controls — is as important as the fat-delivery mechanism itself. Fat amplifies the delivery of an antigen that is already antigen-specific. Remove the bovine-specific antigenic component, and the fat delivery route becomes irrelevant to Hamer Loop activity. This is not a fat sensitivity; it is a bovine Neu5Gc sensitivity that fat delivery makes more severe.

For the susceptible individual, the practical implications are clear: concentrated bovine fat sources are the most impactful first targets for elimination, hidden bovine fat in processed foods is a frequently overlooked trigger category, and re-exposure sensitivity following successful elimination is most acute precisely for the high-fat sources that produced the most severe responses before clearance.

References

Samraj, A. N., Pearce, O. M., et al. (2015). A red meat-derived glycan promotes inflammation and cancer progression. *Proceedings of the National Academy of Sciences*, 112(2), 542–547.

Tangvoranuntakul, P., et al. (2003). Human uptake and incorporation of an immunogenic nonhuman dietary sialic acid. *Proceedings of the National Academy of Sciences*, 100(21), 12045–12050.

Dhar, C., Sasmal, A., & Varki, A. (2019). From 'Serum Sickness' to 'Xenosialitis'. *Frontiers in Immunology*.

Tso, P., & Balint, J. A. (1986). Formation and transport of chylomicrons by enterocytes to the lymphatics. *American Journal of Physiology*, 250(6 Pt 1), G715–G726.

Mather, I. H., & Keenan, T. W. (1998). Origin and secretion of milk lipids. *Journal of Mammary Gland Biology and Neoplasia*, 3(3), 259–273.

Zaramela, L. S., Martino, C., Alisson-Silva, F., et al. (2019). Gut bacteria responding to dietary change encode sialidases that exhibit preference for red meat-associated carbohydrates. *Nature Microbiology*, 4(12), 2082–2089.

Hamer, M. (1981–2026). Longitudinal bedside assessment and pattern recognition. Internal Case Files [Unpublished].