

DIETARY XENOANTIGENS AND OSTEOPOROSIS:

A Convergent Triple-Pathway Model Linking Neu5Gc Xenosialitis, ATI/Gliadin-Mediated Intestinal Permeability, and Inflammatory Cytokine Dysregulation to Accelerated Bone Loss

Martin Hamer, RN (Ret.)

Independent Researcher | Hamer Glycan Systems | hamerglycans.com | 2026

Abstract

Background: Osteoporosis affects an estimated 200 million people worldwide and is characterised by progressive loss of bone mineral density, deteriorating bone microarchitecture, and elevated fracture risk. Current treatment focuses primarily on pharmaceutical inhibition of osteoclast activity and calcium-vitamin D supplementation. The role of specific dietary antigens as upstream modulators of the chronic inflammatory state that drives osteoclast-osteoblast imbalance has received limited systematic investigation.

Hypothesis: We propose a convergent triple-pathway model in which two widely consumed dietary xenoantigens — N-glycolylneuraminic acid (Neu5Gc) from red meat and bovine dairy, and wheat-derived amylase-trypsin inhibitors (ATIs) and gliadins — act through three distinct but mechanistically interconnected pathways to accelerate bone loss and impair bone repair in susceptible individuals. Pathway One: Neu5Gc xenosialitis drives chronic NF-κB-mediated systemic inflammation, producing sustained elevation of TNF-α, IL-6, and IL-17 — cytokines that directly stimulate osteoclastogenesis and suppress osteoblast differentiation. Pathway Two: ATI and gliadin-mediated intestinal permeability disrupts calcium and vitamin D absorption at the intestinal epithelium, creating a state of functional nutritional deficiency independent of dietary calcium intake. Pathway Three: chronic interstitial inflammatory fluid accumulation, as proposed in the Hamer Loop framework, creates a hostile periosteal microenvironment that impairs the regenerative capacity of bone-adjacent tissue and sustains local inflammatory signalling at cortical and trabecular surfaces.

Clinical Observations: Two independent observational cases are presented. Subject A (male, 71, retired RN) has documented over 23 years of longitudinal self-provocation and elimination data, including complete resolution of joint and periarticular pain attributed to degenerative disease on dietary trigger elimination. Subject B (female, 63, confirmed osteoporosis diagnosis, currently receiving injectable bisphosphonate therapy) demonstrates a consistent and independently observed correlation between dietary trigger load — particularly white wine, which contains dairy-derived fining agents, ammonium yeast nutrient compounds, and sulfites — and next-day deterioration in joint mobility and pain severity, with partial recovery on days following reduced consumption.

Implications: If confirmed, this framework reframes osteoporosis not solely as a disorder of calcium metabolism and hormonal decline, but as a condition significantly modulated by chronic

dietary antigen-driven inflammation in susceptible individuals. Dietary elimination of Neu5Gc sources and wheat-derived ATIs/gliadins represents a low-risk, zero-cost intervention whose potential to reduce the inflammatory component of bone loss warrants urgent formal investigation, particularly in patients experiencing pharmaceutical intolerance to standard osteoporosis therapies.

Keywords: *osteoporosis; Neu5Gc; xenosialitis; amylase-trypsin inhibitors; gliadins; osteoclastogenesis; bone mineral density; intestinal permeability; calcium absorption; NF- κ B; TNF- α ; IL-6; RANKL; dietary inflammation; bisphosphonate intolerance*

1. Introduction

Osteoporosis is one of the most prevalent and consequential chronic conditions of later life, affecting an estimated 200 million people globally and producing more than 8.9 million fractures annually. It is characterised by reduced bone mineral density, deterioration of bone microarchitecture, and a corresponding increase in fracture risk — particularly at the hip, spine, and wrist — with hip fractures carrying a one-year mortality rate of up to 20% in older adults.

The dominant aetiological framework positions osteoporosis primarily as a disorder of calcium and vitamin D metabolism superimposed on the hormonal changes of menopause in women and andropause in men, with genetic predisposition, physical inactivity, smoking, and alcohol consumption as modifying risk factors. Treatment is correspondingly focused: calcium and vitamin D supplementation to address nutritional deficiency, and pharmaceutical inhibition of osteoclast activity — primarily through bisphosphonates (alendronate, risedronate, zoledronic acid) and denosumab — to slow the rate of bone resorption.

This framework is well-supported by decades of research and has produced treatments that meaningfully reduce fracture risk. It is not, however, complete. A substantial proportion of osteoporosis patients experience poor adherence to or frank intolerance of pharmaceutical therapy — oral bisphosphonates in particular carry a significant burden of gastrointestinal side effects including oesophageal irritation, nausea, and abdominal pain that lead many patients to discontinue treatment. Injectable and intravenous formulations reduce gastrointestinal exposure but introduce their own side effect profiles including flu-like reactions, jaw osteonecrosis risk, and injection site effects. For many patients — particularly those with pre-existing gastrointestinal sensitivity — the available pharmaceutical options are tolerable only at significant cost to quality of life, or not tolerable at all.

The role of chronic systemic inflammation as an upstream driver of osteoclast-osteoblast imbalance is well-established in the rheumatological literature — inflammatory arthropathies such as rheumatoid arthritis, ankylosing spondylitis, and psoriatic arthritis are all associated with accelerated bone loss, and TNF- α inhibition in these conditions produces measurable improvements in bone density alongside its anti-inflammatory effects. What has received comparatively little attention is the question of what is driving the chronic inflammatory state in patients with osteoporosis who do not have a diagnosed inflammatory arthropathy — patients

who present with bone loss and periarticular pain in the absence of a recognised autoimmune driver.

We propose that for a definable subset of susceptible individuals, two widely consumed dietary antigens — N-glycolylneuraminic acid (Neu5Gc) from bovine meat and dairy products, and wheat-derived amylase-trypsin inhibitors (ATIs) and gliadins — constitute exactly this unrecognised upstream driver, operating through three convergent pathways to accelerate bone loss, impair bone repair, and sustain the periarticular inflammatory load that produces the joint pain and mobility deterioration that these patients experience as their daily reality.

2. Background: The Three Pathways

2.1 Pathway One: Neu5Gc Xenosialitis and Inflammatory Cytokine-Driven Osteoclastogenesis

N-glycolylneuraminic acid (Neu5Gc) is a non-human sialic acid produced by most mammals via the enzyme CMP-Neu5Ac hydroxylase (CMAH). Humans carry an inactivating deletion in the CMAH gene that arose approximately 2–3 million years ago, rendering Neu5Gc immunologically foreign. Despite the inability to synthesise Neu5Gc endogenously, humans regularly ingest it through consumption of red meat and bovine dairy products, where it is present at concentrations of 5–30 mg/kg in beef and 7–11 mg/L in bovine milk.

Upon ingestion, dietary Neu5Gc is absorbed at the intestinal mucosa and metabolically incorporated into human cell-surface glycoproteins. Because the human immune system has matured in the complete absence of Neu5Gc, it recognises the incorporated molecule as a xenoantigen and mounts an antibody response. Anti-Neu5Gc IgG antibodies are detectable in the serum of most adult humans at concentrations proportional to habitual red meat and dairy consumption. The binding of these antibodies to tissue-incorporated Neu5Gc generates immune complexes that activate complement and Fc receptor-bearing effector cells, producing a state of chronic low-grade systemic inflammation termed xenosialitis — sustained NF- κ B activation with persistent elevation of TNF- α , IL-6, IL-1 β , and associated inflammatory mediators.

The connection to bone metabolism is direct and well-characterised in the rheumatological literature, even if the Neu5Gc mechanism has not previously been proposed as an upstream driver. TNF- α and IL-6 are among the most potent known stimulators of osteoclastogenesis — the differentiation of monocyte precursors into mature, bone-resorbing osteoclasts. They achieve this through upregulation of RANKL (receptor activator of NF- κ B ligand) expression on osteoblasts and stromal cells, which engages RANK on osteoclast precursors and drives their maturation into active resorptive cells. Simultaneously, TNF- α suppresses osteoblast differentiation and promotes osteoblast apoptosis, reducing the bone-forming capacity of the skeleton even as its resorptive activity is amplified.

IL-17, which is elevated in xenosialitis through the same NF- κ B-driven inflammatory cascade, acts synergistically with TNF- α and IL-6 to further amplify RANKL expression and osteoclast activity. The combined cytokine milieu of chronic xenosialitis — sustained TNF- α , IL-6, IL-1 β ,

and IL-17 — constitutes precisely the inflammatory environment most capable of tipping the osteoclast-osteoblast balance persistently toward net bone resorption.

In this model, osteoporosis in susceptible individuals is not purely a disorder of calcium metabolism or hormonal decline. It is, at least in part, an inflammatory bone disease driven by a chronic dietary antigen that has never previously been identified as a contributor to the condition — because the question has never been asked with Neu5Gc in mind.

2.2 Pathway Two: ATI/Gliadin-Mediated Intestinal Permeability and Calcium Malabsorption

The second pathway operates at the intestinal epithelium, and its mechanism is distinct from but convergent with Pathway One. Wheat-derived amylase-trypsin inhibitors (ATIs) are non-gluten proteins that activate toll-like receptor 4 (TLR4) on intestinal epithelial and immune cells, driving innate immune activation and cytokine release independently of any celiac autoimmunity. Gliadins, the alcohol-soluble fraction of wheat gluten, bind the chemokine receptor CXCR3 on intestinal epithelial cells and trigger release of zonulin, the primary physiological regulator of intestinal tight junction assembly.

Chronic gliadin-mediated zonulin upregulation produces a state of persistent intestinal hyperpermeability — the tight junction proteins ZO-1, occludin, and claudin-1 are progressively degraded, and the paracellular barrier that normally restricts the passage of luminal contents into the subepithelial space becomes functionally compromised. The immunological consequences of this permeability — LPS translocation, systemic metabolic endotoxaemia, and amplified Neu5Gc absorption — are addressed elsewhere in this series. The consequence most directly relevant to osteoporosis is nutritional: a persistently permeable and inflamed intestinal epithelium is a functionally compromised absorptive surface.

Calcium absorption is a tightly regulated, energy-dependent process that occurs primarily in the duodenum and proximal jejunum via transcellular transport mediated by the epithelial calcium channel TRPV6, the intracellular carrier calbindin-D9k, and the basolateral transporter PMCA1b. This transport machinery is dependent on an intact, well-functioning epithelial monolayer. Chronic ATI-driven innate immune activation at the epithelial surface — with associated cytokine-mediated disruption of enterocyte function — and gliadin-mediated tight junction degradation both impair the integrity of the absorptive surface on which calcium transport depends.

Vitamin D absorption, which is fat-soluble and follows chylomicron-mediated lymphatic entry, is similarly dependent on a functional epithelial surface for efficient uptake. Chronic intestinal inflammation reduces the efficiency of fat-soluble vitamin absorption generally, and vitamin D specifically — the critical cofactor for calcium transport protein expression, intestinal calcium absorption, and renal calcium reabsorption.

The result is a state of functional calcium and vitamin D deficiency that is independent of dietary intake. A patient consuming adequate calcium and supplementing vitamin D may nonetheless be absorbing both at significantly below-optimal efficiency if their intestinal epithelium is chronically inflamed and permeabilised by dietary ATIs and gliadins. Standard blood tests for vitamin D may not fully capture functional deficiency at the tissue level if the deficit is primarily

absorptive rather than dietary. And calcium supplementation — a cornerstone of standard osteoporosis management — cannot compensate for an absorptive surface that chronic dietary antigen exposure has rendered functionally inadequate.

2.3 Pathway Three: Periosteal Inflammatory Microenvironment and Impaired Bone Repair

The third pathway is the most novel and the most directly connected to the Hamer Loop framework established in Documents 1 through 5 of this series. It concerns not the systemic cytokine environment (Pathway One) or the intestinal absorptive capacity (Pathway Two), but the local tissue microenvironment immediately adjacent to bone surfaces.

The Hamer Loop model proposes that chronic dietary Neu5Gc exposure generates a state of lymphatic functional impairment — regional lymphatic flow becomes overwhelmed by the inflammatory load, interstitial fluid accumulates in the periarticular and subcutaneous spaces, and the resulting interstitial stasis creates a protein-rich, cytokine-laden microenvironment in the tissues surrounding joints and bone surfaces. The peripheral oedema, shin oedema, and calf tightness documented in the full systemic resolution profile of Document 1 are the macroscopic manifestations of this interstitial accumulation.

Bone repair and remodelling depend critically on the quality of the periosteal microenvironment. Periosteal stem cells — the precursors of both osteoblasts and chondrocytes — reside in the periosteum, the fibrous membrane investing the outer surface of cortical bone. These cells are exquisitely sensitive to their local inflammatory environment: TNF- α and IL-1 β at the concentrations present in xenosialitis-driven interstitial fluid suppress periosteal stem cell proliferation, impair their differentiation toward the osteoblast lineage, and promote their apoptosis. A periosteal microenvironment chronically bathed in xenosialitis-derived inflammatory mediators is therefore not merely a bystander to bone loss — it is actively impairing the regenerative capacity that the skeleton depends on for repair and remodelling.

Furthermore, the vascular supply to cortical bone — the Haversian and Volkmann canal system — is dependent on unobstructed periosteal microvascular flow. Chronic interstitial pressure elevation, as proposed in the Hamer Loop framework, compresses the periosteal microvasculature and reduces oxygen and nutrient delivery to the cortical bone surface. Osteocytes — the mechanosensing cells embedded within the bone matrix that orchestrate adaptive remodelling in response to mechanical load — are particularly vulnerable to hypoxia. Their death, produced by sustained ischaemia from periosteal vascular compression, releases pro-osteoclastic signals that further amplify bone resorption independently of systemic cytokine levels.

This third pathway therefore operates at the tissue level, below the threshold of systemic inflammatory markers, and may continue to drive bone loss even in patients whose C-reactive protein and ESR appear within normal range — because the relevant inflammation is localised to the periosteal microenvironment rather than distributed systemically.

3. The Convergent Model: Three Pathways, One Outcome

The three pathways described above are not independent mechanisms that happen to converge on the same outcome. They are mechanistically interconnected, each amplifying the others in ways that make the combined effect of concurrent Neu5Gc and ATI/gliadin exposure greater than the sum of its parts.

Pathway	Primary Mechanism	Bone-Relevant Consequence	Interaction with Other Pathways
1. Xenosialitis	Chronic NF-κB activation; TNF-α, IL-6, IL-1β, IL-17 elevation	RANKL upregulation → osteoclastogenesis ↑; osteoblast suppression ↑; net bone resorption ↑	Cytokines amplify gut inflammation (Pathway 2) and periosteal microenvironment damage (Pathway 3)
2. Intestinal permeability	ATI/TLR4 innate activation; gliadin/zonulin tight junction degradation	Calcium and vitamin D malabsorption → functional nutritional deficiency independent of dietary intake	Permeability amplifies Neu5Gc absorption (Pathway 1); LPS translocation adds to cytokine load
3. Periosteal inflammatory microenvironment	Hamer Loop interstitial fluid accumulation; periosteal vascular compression; osteocyte hypoxia	Periosteal stem cell suppression; impaired bone repair; osteocyte death → pro-osteoclastic signalling	Fed by Pathway 1 cytokines; worsened by Pathway 2 nutritional deficits reducing repair substrate availability

Table 1. The Convergent Triple-Pathway Model. Each pathway is independently capable of contributing to bone loss; their simultaneous activation in individuals chronically consuming both Neu5Gc and wheat-derived ATIs/gliadins produces a compounded effect that current osteoporosis management frameworks do not account for.

The interaction between pathways is particularly important to appreciate. ATI/gliadin-mediated intestinal permeability does not merely impair calcium absorption — it also enhances the systemic bioavailability of dietary Neu5Gc by facilitating paracellular passage of Neu5Gc-bearing glycoproteins across a compromised epithelial barrier, amplifying Pathway One. The systemic cytokine milieu produced by Pathway One simultaneously drives intestinal mucosal inflammation, worsening the permeability of Pathway Two. And both pathways feed Pathway Three by maintaining the periosteal inflammatory load and reducing the nutritional substrate available for bone repair.

The clinical consequence is a self-reinforcing system in which each dietary exposure event — a meal containing red meat, dairy, and wheat simultaneously, which describes the majority of Western dietary patterns — activates all three pathways concurrently, producing an inflammatory and nutritional environment that is persistently hostile to bone maintenance and repair.

4. Clinical Observations

4.1 Subject A: 23-Year Longitudinal Self-Observation

Subject A is a 71-year-old male retired registered nurse with 45 years of clinical experience and 23 years of systematic self-provocation and elimination data documented within the Hamer Loop framework (see Documents 1–6 of this series). The full systemic resolution profile documented in Section 6.4 of Document 1 includes complete resolution of generalised joint and body pain previously attributed to degenerative joint disease on simultaneous elimination of all bovine Neu5Gc sources and wheat-derived ATIs and gliadins, within the same 7–10 day window as the cutaneous, gastrointestinal, and respiratory symptoms.

The clinical significance of this finding was addressed explicitly in Document 1: what had been attributed to degenerative arthritic disease resolved completely without pharmacological intervention on dietary elimination and returned on re-exposure, following the Hamer Loop timeline. Resolution on elimination and return on re-exposure, without any other intervening variable, is a strong indicator of inflammatory rather than degenerative aetiology.

Subject A does not have a confirmed osteoporosis diagnosis. His musculoskeletal observations are therefore most relevant to Pathway Three of the convergent model — periarticular and soft tissue inflammation driven by the Hamer Loop interstitial mechanism — rather than to bone mineral density specifically. They are included here because the periarticular inflammatory load he documents is mechanistically upstream of the periosteal microenvironment disruption proposed in Pathway Three, and because the reversibility of that load on dietary elimination in a 71-year-old male provides observational support for the modifiability of the proposed mechanism.

4.2 Subject B: Independent Observational Case — Confirmed Osteoporosis with Pharmaceutical Intolerance

Subject B is a 63-year-old female with a DEXA scan-confirmed diagnosis of osteoporosis, currently receiving injectable bisphosphonate therapy. She carries a significant daily physical and occupational burden: a 250-kilometre daily commute that limits her time for dietary awareness, meal preparation, and lifestyle modification, and that produces a chronic stress load independently relevant to inflammatory disease progression. She is not aware of the Hamer Loop framework and has made no conscious dietary modifications in response to it.

The following observations are documented by a medically trained observer (Subject A, retired RN) over an extended period of close cohabitation and daily observation:

- A consistent and reproducible correlation exists between dietary trigger load and next-day deterioration in joint mobility and pain severity. On days following high-trigger consumption — particularly wine, cheese, and processed meat products — Subject B presents with markedly reduced mobility, described as moving like a person significantly older than her chronological age, with visible reluctance to bear weight and obvious pain on ambulation.

- On days following reduced or absent trigger consumption, Subject B's mobility and pain presentation improve substantially. These better days are not random; they are temporally and consistently correlated with the dietary pattern of the preceding 24 hours.
- White wine is identified as the single most consistent high-impact trigger in Subject B's dietary pattern. This is mechanistically coherent: white wine routinely contains casein-derived fining agents (a bovine dairy derivative used in clarification), ammonium-containing yeast nutrient compounds used in fermentation, and sulfites. It delivers a combination of Neu5Gc antigen and ammonium catalyst in a liquid matrix — the liquid delivery mechanism established in Document 2 as an accelerant of the Hamer Loop timeline by reducing mechanical buffering and accelerating epithelial antigen presentation. The severity of Subject B's next-day presentation following white wine consumption is consistent with an ammonium-potentiated Neu5Gc response rather than an alcohol-specific pharmacological effect.
- A period of approximately two weeks during which Subject B abstained from wine produced a transformation in her presentation that was observable independently: markedly improved mobility, reduced pain-related behavioural indicators, and overall functional improvement. This corresponds to the 7–10 day clearance window documented in the Hamer Loop framework, with the additional days accounting for the severity of her chronic load prior to elimination.

Subject B has experienced significant side effects from both oral and injectable bisphosphonate formulations. Oral bisphosphonates produce gastrointestinal symptoms — nausea, oesophageal discomfort, and abdominal pain — sufficient to cause her to discontinue oral therapy. Injectable formulations produce systemic flu-like reactions. This pharmaceutical intolerance profile is itself mechanistically relevant: several oral bisphosphonate formulations contain lactose monohydrate as a tablet filler and, in some preparations, gelatin capsule shells — both of which are bovine-derived Neu5Gc sources. As documented in Section 6.1 of Document 1 and Section 4.2 of Document 6 in this series, the Gelatin-Antihistamine Paradox demonstrates that therapeutic vehicles containing bovine-derived excipients may partially or wholly counteract the intended therapeutic effect by reintroducing the dietary antigen driving the inflammatory state. The gastrointestinal side effects of oral bisphosphonates in a Neu5Gc-sensitive individual may therefore be compounded by the bovine-derived excipients in the formulation itself — a possibility that has not, to the author's knowledge, been investigated.

Subject B has, in the observer's assessment, resigned herself to the permanence of her condition and the inadequacy of her pharmaceutical options. She is not alone in this. The osteoporosis patient who cannot tolerate her medications, whose pain and mobility are steadily declining, and who has been told implicitly or explicitly that this is now simply her life — this patient deserves to know whether there is a modifiable dietary component to her condition. She deserves to be asked the question. And the answer to that question is what this paper is attempting to provide the scientific basis for asking.

5. The Wine Problem: A Mechanistic Note

White wine warrants specific mechanistic attention because it represents one of the most commonly consumed and least-suspected sources of the convergent triple-pathway trigger in adult Western populations, particularly among women in the age demographic most affected by osteoporosis.

The standard explanation for wine's association with osteoporosis risk focuses on alcohol's direct effects: impairment of osteoblast function, interference with vitamin D metabolism, and increased falls risk from impaired coordination. These mechanisms are real and well-documented. They are not, however, the complete picture for susceptible individuals within the Hamer Loop framework.

Commercial white wine production routinely involves:

- Casein fining: bovine milk casein is added to the must to bind and precipitate tannins and other phenolic compounds before filtration. While the casein is subsequently removed, residual casein-derived proteins — including Neu5Gc-bearing glycopeptides — may remain in the finished wine at concentrations below the detection threshold of standard allergen testing but potentially above the threshold for immunological recognition in sensitised individuals
- Ammonium-containing yeast nutrients: diammonium phosphate (DAP) and ammonium sulfate are routinely added to fermentation vessels as yeast nutrients to support fermentation completion. These are ammonium compounds of the category identified in Document 2 as potent Hamer Loop threshold-lowerers through mast cell sensitisation and NLRP3 inflammasome priming
- Sulfites: sulfur dioxide and sulfite salts are added as antimicrobials and antioxidants. Sulfites independently trigger mast cell degranulation in sensitised individuals and may compound the ammonium potentiation effect
- In some producers, caramel colour for colour adjustment: Class III and IV caramel colours are among the highest-severity ammonium sources in the Neu5Gc Severity Index (Appendix II of this series)

The combination of these compounds in a liquid matrix — with its rapid epithelial presentation and absence of mechanical buffering — makes white wine one of the most efficient delivery vehicles for the convergent triple-pathway trigger profile in the food supply. Its liquid form accelerates the Hamer Loop timeline. Its ammonium content lowers the inflammatory threshold. Its casein-derived residues provide Neu5Gc antigen. And its consumption in the evening, followed by the recumbent nocturnal amplification phase of the Hamer Loop, produces the worst-case conditions for overnight inflammatory sequestration and morning mobility deterioration.

The fact that this is a glass of wine at the end of a long day — a comfort, a reward, an entirely reasonable human pleasure — is not lost on this author. The goal of this framework is not to eliminate pleasure from the lives of people who are already carrying heavy loads. It is to give those people information about what their bodies are doing in response to what they consume, so that they can make informed choices. Knowledge is not compulsion. Understanding the mechanism is not the same as being required to act on it. But a woman who has been told that her deteriorating bones are an inevitable feature of her age and her hormones deserves to know that there may be more to the story.

6. Testable Predictions

The convergent triple-pathway model generates the following specific, falsifiable predictions amenable to investigation with existing research tools:

Prediction	Investigation Design	Expected Finding
Anti-Neu5Gc antibody titres will be elevated in osteoporosis patients relative to age- and sex-matched controls on equivalent dietary patterns	Cross-sectional serological study; validated anti-Neu5Gc ELISA (Varki laboratory assay); stratification by dietary Neu5Gc burden via food frequency questionnaire	Higher anti-Neu5Gc IgG in osteoporosis cohort; correlation between titre and RANKL/OPG ratio and bone turnover markers (CTX, P1NP)
Dietary Neu5Gc and ATI/gluten elimination will produce measurable improvements in bone turnover markers within 12 weeks	Randomised controlled dietary elimination trial; primary outcomes: serum CTX (bone resorption marker), serum P1NP (bone formation marker), hsCRP, IL-6, TNF- α ; secondary: DEXA at 12 months	Reduced CTX; improved CTX:P1NP ratio; reduced inflammatory markers; trend toward improved bone density at 12 months
Intestinal permeability markers will be elevated in osteoporosis patients and will correlate with calcium absorption efficiency	Lactulose:mannitol urinary ratio; serum zonulin; FABP2; dual-isotope calcium absorption test; correlation analysis	Elevated permeability in osteoporosis cohort; inverse correlation between permeability markers and calcium absorption efficiency
Periosteal cytokine concentration will exceed systemic levels in osteoporosis patients consuming Neu5Gc-rich diets	Periosteal biopsy fluid analysis in elective orthopaedic patients; local vs systemic TNF- α , IL-6, IL-17 concentrations; correlation with dietary Neu5Gc burden	Elevated periosteal cytokine concentration relative to serum; correlation with Neu5Gc dietary burden; gradient consistent with local inflammatory sequestration
Bisphosphonate side effects will be reduced in susceptible individuals when oral formulations containing lactose monohydrate or gelatin excipients are replaced with formulations free of bovine-derived excipients	Crossover trial: standard bisphosphonate formulation vs bovine-excipient-free equivalent; primary outcome: gastrointestinal adverse event rate; secondary: anti-Neu5Gc titres during treatment	Reduced gastrointestinal adverse events with bovine-excipient-free formulation; correlation with anti-Neu5Gc antibody titre at baseline

Table 2. Testable Predictions of the Convergent Triple-Pathway Model. Predictions 1–4 test the mechanistic framework; Prediction 5 addresses the clinically immediate question of pharmaceutical excipient contribution to bisphosphonate intolerance.

7. Limitations

This paper presents a structured hypothesis, not established evidence. The convergent triple-pathway model draws on mechanistic evidence from independent fields — glycobiology, gastroenterology, rheumatology, and bone biology — that have not been studied in an integrated framework, and no prospective clinical trial has directly tested dietary Neu5Gc and ATI/gliadin elimination as an intervention in osteoporosis.

The two clinical observations presented in Section 4 are N-of-1 or N-of-2 in character. Subject A's musculoskeletal resolution data, while longitudinally extensive, does not include bone density measurement and therefore cannot speak directly to bone mineral density outcomes. Subject B's observations, while independently documented by a medically trained observer, are not controlled: the correlation between her dietary pattern and next-day functional status, though consistent and reproducible in the observer's assessment, could in principle reflect confounders including alcohol pharmacology, sleep quality, activity level, and psychological state — all of which may be influenced by wine consumption through mechanisms unrelated to the proposed Neu5Gc pathway.

The Pathway Three periosteal microenvironment mechanism is the most novel and least supported by existing literature, as it is an extension of the Hamer Loop interstitial fluid hypothesis to the periosteal context. The claim that interstitial pressure elevation at the periosteal surface is sufficient to produce clinically meaningful osteocyte hypoxia and periosteal stem cell suppression has not been directly tested.

The wine excipient analysis in Section 5 is based on known commercial wine production practices. The actual concentrations of casein-derived Neu5Gc residues in finished commercial white wine have not been measured, and it is acknowledged that standard fining processes remove the majority of added casein. The hypothesis that residual Neu5Gc-bearing peptides at sub-allergen concentrations are sufficient to trigger immune recognition in highly sensitised individuals is mechanistically plausible but unconfirmed.

Individual variability in anti-Neu5Gc antibody titre, intestinal permeability status, HLA type, and the genetic determinants of osteoclast-osteoblast balance means that the triple-pathway model will not apply uniformly across all osteoporosis patients. It is proposed as relevant to a definable susceptible subset — characterised by the immunological and genetic factors described in the companion paper on cutaneous immune phenotype — rather than as a universal mechanism.

8. A Note on Pharmaceutical Excipients

The pharmaceutical intolerance documented in Subject B's case raises a question that deserves explicit investigation independent of the broader osteoporosis hypothesis. Standard oral bisphosphonate formulations — including the most widely prescribed agents alendronate

and risedronate — commonly contain lactose monohydrate as a tablet filler. Lactose monohydrate is a dairy-derived disaccharide that, while purified, may retain trace quantities of bovine-derived proteins including Neu5Gc-bearing glycopeptides at concentrations below standard allergen testing thresholds.

As established in the Gelatin-Antihistamine Paradox (Document 1, Section 6.1; Document 6, Section 4.2 of this series), the vehicle of a therapeutic agent can reintroduce the dietary antigen that the treatment is intended to address, partially or wholly counteracting the therapeutic intent. In the specific context of osteoporosis treatment, this creates a potentially self-defeating scenario: a patient whose bone loss is partially driven by Neu5Gc xenosialitis-mediated osteoclastogenesis receives a bisphosphonate formulation whose excipient introduces further Neu5Gc antigen with each dose, sustaining the inflammatory osteoclastic drive that the bisphosphonate is attempting to suppress.

This hypothesis is directly testable through the crossover trial described in Prediction 5 of Section 6. It requires no novel compounds and no experimental protocols — only the comparison of gastrointestinal adverse event rates between standard bisphosphonate formulations and equivalent formulations manufactured without bovine-derived excipients, stratified by baseline anti-Neu5Gc antibody titre. The clinical benefit, if the hypothesis is confirmed, would be immediate and actionable: reformulation guidance for pharmaceutical manufacturers, and prescribing guidance for clinicians managing bisphosphonate-intolerant osteoporosis patients.

9. Conclusion

Osteoporosis is not a simple disease. For many patients it is a convergence of hormonal change, genetic predisposition, nutritional status, physical activity, and — we propose — a chronic dietary antigen-driven inflammatory state that has not previously been identified as a contributor because the relevant antigens have not previously been examined in this context.

The convergent triple-pathway model presented here does not claim to explain all osteoporosis or to replace established aetiology. It claims something more specific and more actionable: that for a definable subset of susceptible individuals, chronic exposure to dietary Neu5Gc and wheat-derived ATIs and gliadins is accelerating bone loss through three mechanistically distinct and mutually amplifying pathways, and that elimination of these dietary antigens represents a low-risk, zero-cost intervention whose potential to reduce the inflammatory component of bone resorption and restore intestinal absorptive capacity warrants urgent formal clinical investigation.

The patients who most urgently need this investigation are not those who are well-managed on pharmaceutical therapy. They are the ones who cannot tolerate the medications, who experience the full burden of the disease without adequate pharmacological support, and who have been told — gently, clinically, but definitively — that this is now simply their life.

For those patients, the possibility that a dietary modification might reduce their pain, improve their mobility, and slow the progression of their bone loss is not a peripheral research question.

It is the most important clinical question they face. It deserves to be asked, properly, with the rigour that they and the science both require.

References

- Samraj, A. N., Pearce, O. M., et al. (2015). A red meat-derived glycan promotes inflammation and cancer progression. *Proceedings of the National Academy of Sciences*, 112(2), 542–547.
- Dhar, C., Sasmal, A., & Varki, A. (2019). From ‘Serum Sickness’ to ‘Xenosialitis’. *Frontiers in Immunology*.
- Tangvoranuntakul, P., et al. (2003). Human uptake and incorporation of an immunogenic nonhuman dietary sialic acid. *Proceedings of the National Academy of Sciences*, 100(21), 12045–12050.
- Junker, Y., Zeissig, S., Kim, S. J., et al. (2012). Wheat amylase trypsin inhibitors drive intestinal inflammation via activation of toll-like receptor 4. *Journal of Experimental Medicine*, 209(13), 2395–2408.
- Lammers, K. M., Lu, R., Brownley, J., et al. (2008). Gliadin induces an increase in intestinal permeability and zonulin release by binding to the chemokine receptor CXCR3. *Gastroenterology*, 135(1), 194–204.
- Boyle, W. J., Simonet, W. S., & Lacey, D. L. (2003). Osteoclast differentiation and activation. *Nature*, 423(6937), 337–342.
- Schett, G., & Gravallese, E. (2012). Bone erosion in rheumatoid arthritis: mechanisms, diagnosis and treatment. *Nature Reviews Rheumatology*, 8(11), 656–664.
- Khosla, S., Oursler, M. J., & Monroe, D. G. (2012). Estrogen and the skeleton. *Trends in Endocrinology and Metabolism*, 23(11), 576–581.
- Fasano, A. (2012). Intestinal permeability and its regulation by zonulin: diagnostic and therapeutic implications. *Clinical Gastroenterology and Hepatology*, 10(10), 1096–1100.
- Cani, P. D., Amar, J., Iglesias, M. A., et al. (2007). Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes*, 56(7), 1761–1772.
- Compston, J. E., McClung, M. R., & Leslie, W. D. (2019). Osteoporosis. *Lancet*, 393(10169), 364–376.
- Reid, I. R., & Billington, E. O. (2022). Drug therapy for osteoporosis in older adults. *Lancet*, 399(10329), 1080–1092.
- Hamer, M. (2026). The Hamer Loop & Xeno-Antigenic Sequestration [Document 1 of the Hamer Loop Manuscript Series]. Hamer Glycan Systems [Unpublished].
- Hamer, M. (2026). Acnerase — Hamer Loop Live Demonstration Protocol [Document 6 of the Hamer Loop Manuscript Series]. Hamer Glycan Systems [Unpublished].

Conflict of Interest: None declared. | **Funding:** None. | **Ethical approval:** Not applicable (hypothesis paper). Subject B observations are reported with the knowledge and permission of the observing author. Subject B is identified only by age and sex.

© 2026 Martin Hamer, RN (Ret.) | Hamer Glycan Systems | hamerglycans.com | *This hypothesis is offered for scholarly discussion and does not constitute clinical advice.*