

A focused model selection criterion for selecting among parametric and nonparametric models

Martin Jullum

Norwegian Computing Center


jullum@nr.no

May 23, 2017


Outline

- 1 Motivation and idea
- 2 FIC derivation
- 3 Data types and situations
- 4 Properties
- 5 Summary
- 6 Bonus


I.i.d. data: Cumulative distribution



Time series: Covariance structure



Survival analysis: Survival curves





A focused model selection criterion for selecting among parametric and nonparametric models

Model selection

- Consider i.i.d. data Y_1, \dots, Y_n from some unknown underlying distribution G
- Parametric approach
 - Restrict G to a specific parametric family F_θ
 - Estimate parameter θ (e.g. by ML) and use $\hat{G} = F_{\hat{\theta}}$
- Nonparametric approach
 - Let the data speak for themselves, no structural assumptions
 - Estimate G by ecdf:
 $\hat{G}_n(y) = n^{-1} \#\{Y_i \leq y\}.$
 - Smoother versions: Local kernel smoothers
- $K + 1$ different appropriate models – which one should we trust?



Model selection

- Consider i.i.d. data Y_1, \dots, Y_n from some unknown underlying distribution G
- Parametric approach
 - Restrict G to a specific parametric family F_θ
 - Estimate parameter θ (e.g. by ML) and use $\hat{G} = F_{\hat{\theta}}$
- Nonparametric approach
 - Let the data speak for themselves, no structural assumptions
 - Estimate G by ecdf:
 $\hat{G}_n(y) = n^{-1} \#\{Y_i \leq y\}$.
 - Smoother versions: Local kernel smoothers
- $K + 1$ different appropriate models – which one should we trust?



Model selection

- Consider i.i.d. data Y_1, \dots, Y_n from some unknown underlying distribution G
- Parametric approach
 - Restrict G to a specific parametric family F_θ
 - Estimate parameter θ (e.g. by ML) and use $\hat{G} = F_{\hat{\theta}}$
- Nonparametric approach
 - Let the data speak for themselves, no structural assumptions
 - Estimate G by ecdf:
 $\hat{G}_n(y) = n^{-1} \#\{Y_i \leq y\}$.
 - Smoother versions: Local kernel smoothers
- $K + 1$ different appropriate models – which one should we trust?



Classical parametric model selection

- Information criteria

- $AIC = 2 \log\text{-likelihood}_{\max} - 2 \dim(\theta)$
- $BIC = 2 \log\text{-likelihood}_{\max} - (\log n) \dim(\theta)$
- DIC, GIC, TIC, etc...

- Select the model optimizing the information criterion
- Cannot handle nonparametrics

Classical parametric model selection

- Information criteria

- $AIC = 2 \log\text{-likelihood}_{\max} - 2 \dim(\theta)$
- $BIC = 2 \log\text{-likelihood}_{\max} - (\log n) \dim(\theta)$
- DIC, GIC, TIC, etc...

- Select the model optimizing the information criterion
- Cannot handle nonparametrics

Classical parametric model selection

- Information criteria
 - $AIC = 2 \log\text{-likelihood}_{\max} - 2 \dim(\theta)$
 - $BIC = 2 \log\text{-likelihood}_{\max} - (\log n) \dim(\theta)$
 - DIC, GIC, TIC, etc...
- Select the model optimizing the information criterion
- Cannot handle nonparametrics

Classical parametric model selection

- Information criteria
 - $AIC = 2 \log\text{-likelihood}_{\max} - 2 \dim(\theta)$
 - $BIC = 2 \log\text{-likelihood}_{\max} - (\log n) \dim(\theta)$
 - DIC, GIC, TIC, etc...
- Select the model optimizing the information criterion
- Cannot handle nonparametrics

Goodness of fit measures

- Measures the quality of the parametric fit compared to nonparametric
 - Cramér–von Mises: $\int (\hat{G}_n(y) - F(y; \hat{\theta}))^2 dF(y; \hat{\theta})$
 - Kolmogorov–Smirnov: $\sup_y |\hat{G}_n(y) - F(y; \hat{\theta})|$
 - Categorical data: Pearson's chi-squared: $\sum_{j=1}^k \frac{(N_j - n f_j(\hat{\theta}))^2}{n f_j(\hat{\theta})}$
- Problematic to use for selecting between nonparametric and k parametric models
 - How to set thresholds for rejecting the parametric models?
 - How to penalize parametric model complexity?

Goodness of fit measures

- Measures the quality of the parametric fit compared to nonparametric
 - Cramér–von Mises: $\int (\hat{G}_n(y) - F(y; \hat{\theta}))^2 dF(y; \hat{\theta})$
 - Kolmogorov–Smirnov: $\sup_y |\hat{G}_n(y) - F(y; \hat{\theta})|$
 - Categorical data: Pearson's chi-squared: $\sum_{j=1}^k \frac{(N_j - n f_j(\hat{\theta}))^2}{n f_j(\hat{\theta})}$
- Problematic to use for selecting between nonparametric and k parametric models
 - How to set thresholds for rejecting the parametric models?
 - How to penalize parametric model complexity?

Parametrics or nonparametrics

- No appropriate criterion for model selection among parametrics and nonparametrics in the literature
- Different models have strengths and weaknesses on different parts of the data space
- Why you are doing the analysis should reflect your choice of model

Our proposed solution

A focused or interest driven model selection criterion for selection among a set of parametric and nonparametric models

Parametrics or nonparametrics

- No appropriate criterion for model selection among parametrics and nonparametrics in the literature
- Different models have strengths and weaknesses on different parts of the data space
- Why you are doing the analysis should reflect your choice of model

Our proposed solution

A focused or interest driven model selection criterion for selection among a set of parametric and nonparametric models

Parametrics or nonparametrics

- No appropriate criterion for model selection among parametrics and nonparametrics in the literature
- Different models have strengths and weaknesses on different parts of the data space
- Why you are doing the analysis should reflect your choice of model

Our proposed solution

A focused or interest driven model selection criterion for selection among a set of parametric and nonparametric models

Focus parameter

- A general population quantity of interest – not(!) a model specific parameter
- A quantity μ mapping the distribution G to a scalar (multivariate situation later)

Examples


- Expectation:
$$\mu = E_G(Y_i) = \int y \, dG(y)$$
- $\Pr(Y_i > 2)$: $\mu = 1 - G(2)$
- Interquartile range:
$$\mu = G^{-1}(3/4) - G^{-1}(1/4)$$

Focus parameter

- A general population quantity of interest – not(!) a model specific parameter
- A quantity μ mapping the distribution G to a scalar (multivariate situation later)

Examples

- Expectation:
$$\mu = E_G(Y_i) = \int y dG(y)$$
- $\Pr(Y_i > 2)$: $\mu = 1 - G(2)$
- Interquartile range:
$$\mu = G^{-1}(3/4) - G^{-1}(1/4)$$




Focus parameter

- A general population quantity of interest – not(!) a model specific parameter
- A quantity μ mapping the distribution G to a scalar (multivariate situation later)

Examples

- Expectation:
$$\mu = E_G(Y_i) = \int y dG(y)$$
- $\Pr(Y_i > 2)$: $\mu = 1 - G(2)$
- Interquartile range:
$$\mu = G^{-1}(3/4) - G^{-1}(1/4)$$




Focus parameter

- A general population quantity of interest – not(!) a model specific parameter
- A quantity μ mapping the distribution G to a scalar (multivariate situation later)


Examples

- Expectation:
$$\mu = E_G(Y_i) = \int y dG(y)$$
- $\Pr(Y_i > 2)$: $\mu = 1 - G(2)$
- Interquartile range:
$$\mu = G^{-1}(3/4) - G^{-1}(1/4)$$



Simple illustration

- I.i.d. univariate observations
- Focus parameter of interest:
 $\mu = G^{-1}(1/2)$, the median of the unknown data generating distribution G
- Gaussian or nonparametric?
- Nonparametric sample median $\hat{G}_n^{-1}(0.5)$ or the Gaussian alternative $\bar{Y} = n^{-1} \sum_{i=1}^n Y_i$?



Criterion idea

- Model selection problem \rightarrow Best estimator for μ
 - Compare estimated performance of $\hat{\mu}_{\text{pm},1}, \dots, \hat{\mu}_{\text{pm},k}$ and $\hat{\mu}_{\text{np}}$
 - Select model with best performing estimator $\hat{\mu}_M$
- Performance measure:

$$\text{risk} = \text{mse}(\hat{\mu}_M) = \text{E} \{ (\hat{\mu}_M - \mu_{\text{true}})^2 \} = \text{bias}^2(\hat{\mu}_M) + \text{Var}(\hat{\mu}_M)$$

Basic idea: Focused information criterion (FIC)

- Estimate the mean squared error (mse) as squared bias + variance:

$$\text{FIC}(M) = \widehat{\text{mse}}(\hat{\mu}_M) = \widehat{\text{bias}}^2(\hat{\mu}_M) + \widehat{\text{Var}}(\hat{\mu}_M)$$

- Choose the model and estimator with the smallest FIC score

Criterion idea

- Model selection problem \rightarrow Best estimator for μ
 - Compare estimated performance of $\hat{\mu}_{\text{pm},1}, \dots, \hat{\mu}_{\text{pm},k}$ and $\hat{\mu}_{\text{np}}$
 - Select model with best performing estimator $\hat{\mu}_M$
- Performance measure:

$$\text{risk} = \text{mse}(\hat{\mu}_M) = \text{E} \{ (\hat{\mu}_M - \mu_{\text{true}})^2 \} = \text{bias}^2(\hat{\mu}_M) + \text{Var}(\hat{\mu}_M)$$

Basic idea: Focused information criterion (FIC)

- Estimate the mean squared error (mse) as squared bias + variance:

$$\text{FIC}(M) = \widehat{\text{mse}}(\hat{\mu}_M) = \widehat{\text{bias}}^2(\hat{\mu}_M) + \widehat{\text{Var}}(\hat{\mu}_M)$$

- Choose the model and estimator with the smallest FIC score

FIC construction: Derivation details (I)

- Let (generically) $\mu_{0,\text{pm}} = T(F_{\theta_0})$ be the least false focus parameter value with θ_0 be the least false parameter value of θ

Under weak regularity conditions:

- Nonparametric: $\sqrt{n}(\hat{\mu}_{\text{np}} - \mu_{\text{true}}) \rightarrow_d \sim N(0, v_{\text{np}})$
- Parametric $\sqrt{n}(\hat{\mu}_{\text{pm}} - \mu_{0,\text{pm}}) \rightarrow_d \sim N(0, v_{\text{pm}})$

Large sample approximate risks

$$\text{Nonparametric: } \text{mse}(\hat{\mu}_{\text{np}}) \approx \widetilde{\text{mse}}(\hat{\mu}_{\text{np}}) = 0 + \frac{1}{n}v_{\text{np}}$$

$$\text{Parametric: } \text{mse}(\hat{\mu}_{\text{pm}}) \approx \widetilde{\text{mse}}(\hat{\mu}_{\text{pm}}) = b^2 + \frac{1}{n}v_{\text{pm}}$$

where $b = \mu_{0,\text{pm}} - \mu_{\text{true}}$

FIC construction: Derivation details (I)

- Let (generically) $\mu_{0,\text{pm}} = T(F_{\theta_0})$ be the least false focus parameter value with θ_0 be the least false parameter value of θ

Under weak regularity conditions:

- Nonparametric: $\sqrt{n}(\hat{\mu}_{\text{np}} - \mu_{\text{true}}) \rightarrow_d \sim N(0, v_{\text{np}})$
- Parametric $\sqrt{n}(\hat{\mu}_{\text{pm}} - \mu_{0,\text{pm}}) \rightarrow_d \sim N(0, v_{\text{pm}})$

Large sample approximate risks

$$\text{Nonparametric: } \text{mse}(\hat{\mu}_{\text{np}}) \approx \widetilde{\text{mse}}(\hat{\mu}_{\text{np}}) = 0 + \frac{1}{n}v_{\text{np}}$$

$$\text{Parametric: } \text{mse}(\hat{\mu}_{\text{pm}}) \approx \widetilde{\text{mse}}(\hat{\mu}_{\text{pm}}) = b^2 + \frac{1}{n}v_{\text{pm}}$$

where $b = \mu_{0,\text{pm}} - \mu_{\text{true}}$

FIC construction: Derivation details (II)

- The convergence in distribution also holds jointly:

$$\sqrt{n} \begin{pmatrix} \hat{\mu}_{\text{np}} - \mu_{\text{true}} \\ \hat{\mu}_{\text{pm}} - \mu_{0,\text{pm}} \end{pmatrix} \rightarrow_d N_2 \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} v_{\text{np}} & v_c \\ v_c & v_{\text{pm}} \end{pmatrix} \right)$$

- $\Rightarrow \sqrt{n}(\hat{b} - b) \rightarrow_d N(0, v_b)$, with $\hat{b} = \hat{\mu}_{\text{pm}} - \hat{\mu}_{\text{np}}$ and $v_b = v_{\text{pm}} + v_{\text{np}} - 2v_c$
- $\Rightarrow E\{(\hat{b})^2\} = (E\{\hat{b}\})^2 + \text{Var}(\hat{b}) \approx b^2 + \frac{1}{n}v_b$

General estimates of risk components

- Squared parametric bias: b^2 estimated by $\max\{0, \hat{b}^2 - \frac{1}{n}\hat{v}_b\}$
- Variances: v_{np} estimated by \hat{v}_{np} , and v_{pm} estimated by \hat{v}_{pm}

FIC scores

Nonparametric: $\text{FIC}(\hat{\mu}_{\text{np}}) = \widehat{\text{mse}}(\hat{\mu}_{\text{np}}) = 0 + \frac{1}{n}\hat{v}_{\text{np}}$

Parametric: $\text{FIC}(\hat{\mu}_{\text{pm}}) = \widehat{\text{mse}}(\hat{\mu}_{\text{pm}}) = \max\{0, \hat{b}^2 - \frac{1}{n}\hat{v}_b\} + \frac{1}{n}\hat{v}_{\text{pm}}$

FIC construction: Derivation details (II)

- The convergence in distribution also holds jointly:

$$\sqrt{n} \begin{pmatrix} \hat{\mu}_{np} - \mu_{\text{true}} \\ \hat{\mu}_{pm} - \mu_{0,pm} \end{pmatrix} \rightarrow_d N_2 \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} v_{np} & v_c \\ v_c & v_{pm} \end{pmatrix} \right)$$

- $\Rightarrow \sqrt{n}(\hat{b} - b) \rightarrow_d N(0, v_b)$, with $\hat{b} = \hat{\mu}_{pm} - \hat{\mu}_{np}$ and $v_b = v_{pm} + v_{np} - 2v_c$
- $\Rightarrow E\{(\hat{b})^2\} = (E\{\hat{b}\})^2 + \text{Var}(\hat{b}) \approx b^2 + \frac{1}{n}v_b$

General estimates of risk components

- Squared parametric bias: b^2 estimated by $\max\{0, \hat{b}^2 - \frac{1}{n}\hat{v}_b\}$
- Variances: v_{np} estimated by \hat{v}_{np} , and v_{pm} estimated by \hat{v}_{pm}

FIC scores

Nonparametric: $\text{FIC}(\hat{\mu}_{np}) = \widehat{\text{mse}}(\hat{\mu}_{np}) = 0 + \frac{1}{n}\hat{v}_{np}$

Parametric: $\text{FIC}(\hat{\mu}_{pm}) = \widehat{\text{mse}}(\hat{\mu}_{pm}) = \max\{0, \hat{b}^2 - \frac{1}{n}\hat{v}_b\} + \frac{1}{n}\hat{v}_{pm}$

FIC construction: Derivation details (II)

- The convergence in distribution also holds jointly:

$$\sqrt{n} \begin{pmatrix} \hat{\mu}_{\text{np}} - \mu_{\text{true}} \\ \hat{\mu}_{\text{pm}} - \mu_{0,\text{pm}} \end{pmatrix} \rightarrow_d N_2 \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} v_{\text{np}} & v_c \\ v_c & v_{\text{pm}} \end{pmatrix} \right)$$

- $\Rightarrow \sqrt{n}(\hat{b} - b) \rightarrow_d N(0, v_b)$, with $\hat{b} = \hat{\mu}_{\text{pm}} - \hat{\mu}_{\text{np}}$ and $v_b = v_{\text{pm}} + v_{\text{np}} - 2v_c$
- $\Rightarrow E\{(\hat{b})^2\} = (E\{\hat{b}\})^2 + \text{Var}(\hat{b}) \approx b^2 + \frac{1}{n}v_b$

General estimates of risk components

- Squared parametric bias: b^2 estimated by $\max\{0, \hat{b}^2 - \frac{1}{n}\hat{v}_b\}$
- Variances: v_{np} estimated by \hat{v}_{np} , and v_{pm} estimated by \hat{v}_{pm}

FIC scores

Nonparametric: $\text{FIC}(\hat{\mu}_{\text{np}}) = \widehat{\text{mse}}(\hat{\mu}_{\text{np}}) = 0 + \frac{1}{n}\hat{v}_{\text{np}}$

Parametric: $\text{FIC}(\hat{\mu}_{\text{pm}}) = \widehat{\text{mse}}(\hat{\mu}_{\text{pm}}) = \max\{0, \hat{b}^2 - \frac{1}{n}\hat{v}_b\} + \frac{1}{n}\hat{v}_{\text{pm}}$

Averaged Focused Information Criterion (AFIC)

- Focus on a (weighted) set of focus parameters simultaneously
- Performance measure: $\text{risk} = \int \mathbb{E} [(\hat{\mu}(t) - \mu_{\text{true}}(t))^2] dW(t)$,
for some cumulative weight function W

AFIC scheme

$$\text{Nonparametric: AFIC}(\hat{\mu}_{\text{np}}) = \int \frac{1}{n} V_{\text{np}}(t; \hat{G}_n) dW(t)$$

$$\begin{aligned} \text{Parametric: AFIC}(\hat{\mu}_{\text{pm}}) = \max & \left[0, \int \{b(t; \hat{\theta}, \hat{G}_n)^2 - \frac{1}{n} V_{\text{b}}(t; \hat{\theta}, \hat{G}_n)\} dW(t) \right] \\ & + \int \frac{1}{n} V_{\text{pm}}(t; \hat{\theta}) dW(t) \end{aligned}$$

The criterion selects the model with the smallest AFIC score

- Estimate W empirically if it depends on unknown quantities

Averaged Focused Information Criterion (AFIC)

- Focus on a (weighted) set of focus parameters simultaneously
- Performance measure: $\text{risk} = \int \mathbb{E} [(\hat{\mu}(t) - \mu_{\text{true}}(t))^2] dW(t)$,
for some cumulative weight function W

AFIC scheme

$$\text{Nonparametric: AFIC}(\hat{\mu}_{\text{np}}) = \int \frac{1}{n} V_{\text{np}}(t; \hat{G}_n) dW(t)$$

$$\begin{aligned} \text{Parametric: AFIC}(\hat{\mu}_{\text{pm}}) = \max & \left[0, \int \{b(t; \hat{\theta}, \hat{G}_n)^2 - \frac{1}{n} V_{\text{b}}(t; \hat{\theta}, \hat{G}_n)\} dW(t) \right] \\ & + \int \frac{1}{n} V_{\text{pm}}(t; \hat{\theta}) dW(t) \end{aligned}$$

The criterion selects the model with the smallest AFIC score

- Estimate W empirically if it depends on unknown quantities

Averaged Focused Information Criterion (AFIC)

- Focus on a (weighted) set of focus parameters simultaneously
- Performance measure: $\text{risk} = \int \mathbb{E} [(\hat{\mu}(t) - \mu_{\text{true}}(t))^2] dW(t)$,
for some cumulative weight function W

AFIC scheme

$$\text{Nonparametric: AFIC}(\hat{\mu}_{\text{np}}) = \int \frac{1}{n} V_{\text{np}}(t; \hat{G}_n) dW(t)$$

$$\begin{aligned} \text{Parametric: AFIC}(\hat{\mu}_{\text{pm}}) = \max & \left[0, \int \{b(t; \hat{\theta}, \hat{G}_n)^2 - \frac{1}{n} V_{\text{b}}(t; \hat{\theta}, \hat{G}_n)\} dW(t) \right] \\ & + \int \frac{1}{n} V_{\text{pm}}(t; \hat{\theta}) dW(t) \end{aligned}$$

The criterion selects the model with the smallest AFIC score

- Estimate W empirically if it depends on unknown quantities

Averaged Focused Information Criterion (AFIC)

- Focus on a (weighted) set of focus parameters simultaneously
- Performance measure: $\text{risk} = \int \mathbb{E} [(\hat{\mu}(t) - \mu_{\text{true}}(t))^2] dW(t)$,
for some cumulative weight function W

AFIC scheme

$$\text{Nonparametric: AFIC}(\hat{\mu}_{\text{np}}) = \int \frac{1}{n} V_{\text{np}}(t; \hat{G}_n) dW(t)$$

$$\begin{aligned} \text{Parametric: AFIC}(\hat{\mu}_{\text{pm}}) = \max & \left[0, \int \{b(t; \hat{\theta}, \hat{G}_n)^2 - \frac{1}{n} V_{\text{b}}(t; \hat{\theta}, \hat{G}_n)\} dW(t) \right] \\ & + \int \frac{1}{n} V_{\text{pm}}(t; \hat{\theta}) dW(t) \end{aligned}$$

The criterion selects the model with the smallest AFIC score

- Estimate W empirically if it depends on unknown quantities

Data types and situations (I)

General setup requirements

- Some data generating mechanism G
- Focus parameters which can be written as a functional T of the data generating mechanism G : $\mu = T(G)$
- Focus parameter μ estimated by plug-in for each model M :
 $\hat{\mu}_M = T(\hat{G}_M)$

As seen: Standard i.i.d. situation

- G is cumulative distribution function
- Nonparametric estimation: Cumulative distribution function
 $\hat{G}_n = n^{-1} \# \{Y_i \leq y\}$ or smoothed versions
- Parametric estimation: Whatever you can think of
- Typical focus parameters: Smooth functions of means and quantiles

Data types and situations (II)

Dependency modelling in stationary Gaussian time series models

- G = Unknown spectral measure/distribution
- Nonparametric estimation: Integrated periodogram
 $\hat{G}_n(\omega) = \int_{-\pi}^{\omega} I_n(u) \, du$ for I_n the periodogram
- Parametric estimation: Typically ARMA-models, but also more general parametric models for the spectral distribution
- Typical focus parameters: Differences in spectral distribution, covariance lags and correlation lags

Data types and situations (III)

Censoring

- Semiparametric Cox regression vs. fully parametric proportional hazard regression models
 - $G = (A(\cdot), \beta)$ corresponding to the hazard rate function: $\alpha(s) \exp(x^t \beta)$, with α unspecified baseline hazard and A its cumulative
 - Semiparametric Cox regression: $\hat{\beta}_{\text{cox}}$ via Cox's partial likelihood, $\hat{A}_{\text{cox}}(\cdot) =$ Breslow estimator
 - Parametric estimation: Joint ML estimation of θ, β , with a parametric hazard rate function α_θ (exponential, Weibull, Gompertz, ...)
 - Typical focus parameters: Survival probabilities, quantiles and cumulative hazards, conditional on covariate values
- Without covariates: Nelson–Aalen or Kaplan–Meier estimators vs. parametric survival models

FIC/AFIC asymptotics

Consider one of the parametric model $M : F_\theta$ with least false model specification F_{θ_0} and least false focus parameter value $\mu_{0,\text{pm}}$:

- If M is biased ($\mu_{\text{true}} \neq \mu_{0,\text{pm}}$)
 - $\Pr(\text{FIC/AFIC selects } M) \rightarrow 0$
- If M is fully correct ($G = F_{\theta_0}$)
 - $\Pr(\text{FIC selects } M \text{ over nonparametric}) \rightarrow \chi_1^2(2) \approx 0.843$
 - No general results for AFIC (depends on μ and weight W)
 - AFIC for parametric with focus on all $\mu(y) = G(y)$ and $W = F_{\theta_0}$
 - Re-invents the Cramér-von Mises goodness-of-fit test, with a threshold value found by the theory itself

\Rightarrow If all parametric candidate models are biased:

$\Pr(\text{FIC/AFIC selects nonparametric}) \rightarrow 1$

- Insurance mechanism against parametric misspecification

FIC/AFIC asymptotics

Consider one of the parametric model $M : F_\theta$ with least false model specification F_{θ_0} and least false focus parameter value $\mu_{0,\text{pm}}$:

- If M is biased ($\mu_{\text{true}} \neq \mu_{0,\text{pm}}$)
 - $\Pr(\text{FIC/AFIC selects } M) \rightarrow 0$
- If M is fully correct ($G = F_{\theta_0}$)
 - $\Pr(\text{FIC selects } M \text{ over nonparametric}) \rightarrow \chi_1^2(2) \approx 0.843$
 - No general results for AFIC (depends on μ and weight W)
 - AFIC for parametric with focus on all $\mu(y) = G(y)$ and $W = F_{\theta_0}$
 - Re-invents the Cramér-von Mises goodness-of-fit test, with a threshold value found by the theory itself

⇒ If all parametric candidate models are biased:

$\Pr(\text{FIC/AFIC selects nonparametric}) \rightarrow 1$

- Insurance mechanism against parametric misspecification

FIC/AFIC asymptotics

Consider one of the parametric model $M : F_\theta$ with least false model specification F_{θ_0} and least false focus parameter value $\mu_{0,\text{pm}}$:

- If M is biased ($\mu_{\text{true}} \neq \mu_{0,\text{pm}}$)
 - $\Pr(\text{FIC/AFIC selects } M) \rightarrow 0$
- If M is fully correct ($G = F_{\theta_0}$)
 - $\Pr(\text{FIC selects } M \text{ over nonparametric}) \rightarrow \chi_1^2(2) \approx 0.843$
 - No general results for AFIC (depends on μ and weight W)
 - AFIC for parametric with focus on all $\mu(y) = G(y)$ and $W = F_{\theta_0}$
 - Re-invents the Cramér-von Mises goodness-of-fit test, with a threshold value found by the theory itself

\Rightarrow If all parametric candidate models are biased:

$\Pr(\text{FIC/AFIC selects nonparametric}) \rightarrow 1$

- Insurance mechanism against parametric misspecification

Summary

- Focus driven model selection with a nonparametric alternative
- Rank models according to $\hat{\mu}_M$'s estimated risk
- Robustifies parametric model selection by including the nonparametric candidate model
- AFIC allows several focus parameters to be handled simultaneously
- FIC and AFIC **Are** model selection schemes, but may also justify the use of different significance levels in hypothesis testing of models

AFIC curiosity 1

- Consider count data $(N_1, \dots, N_k), (p_1, \dots, p_k), \sum p_j = 1, \sum N_j = n$
- Focus on all p_j , with weight $1/p_j$
- Direct comparison between parametrics and nonparametrics reduces to

$$X_n = n \sum \frac{(\hat{p}_{np,j} - \hat{p}_{pm,j})^2}{\hat{p}_{np,j}} \text{ vs. } 2df$$

- Implicit test level for test of pm true with $df=1, \dots, 10$:
0.157, 0.135, 0.112, 0.092, 0.075, 0.062, 0.051, 0.042, 0.035.

AFIC curiosity 1

- Consider count data $(N_1, \dots, N_k), (p_1, \dots, p_k), \sum p_j = 1, \sum N_j = n$
- Focus on all p_j , with weight $1/p_j$
- Direct comparison between parametrics and nonparametrics reduces to

$$X_n = n \sum \frac{(\hat{p}_{\text{np},j} - \hat{p}_{\text{pm},j})^2}{\hat{p}_{\text{np},j}} \text{ vs. } 2\text{df}$$

- Implicit test level for test of pm true with $\text{df}=1, \dots, 10$:
0.157, 0.135, 0.112, 0.092, 0.075, 0.062, 0.051, 0.042, 0.035.

AFIC curiosity 1

- Consider count data $(N_1, \dots, N_k), (p_1, \dots, p_k), \sum p_j = 1, \sum N_j = n$
- Focus on all p_j , with weight $1/p_j$
- Direct comparison between parametrics and nonparametrics reduces to

$$X_n = n \sum \frac{(\hat{p}_{\text{np},j} - \hat{p}_{\text{pm},j})^2}{\hat{p}_{\text{np},j}} \text{ vs. } 2\text{df}$$

- Implicit test level for test of pm true with $\text{df}=1, \dots, 10$:
0.157, 0.135, 0.112, 0.092, 0.075, 0.062, 0.051, 0.042, 0.035.

AFIC curiosity 2

- Focus on the complete $G(y)$, weighted by $W(y) = F(y; \theta_0)$
- Direct comparison between parametrics and nonparametrics reduces to

$$\text{CvM}_n = \int n(\hat{G}_n(y) - F(y; \hat{\theta}))^2 dF(y; \hat{\theta}) \text{ vs. } \kappa$$

- If $G = F \sim N(\xi, \sigma^2)$: $\Pr(\text{Select pm}) \rightarrow 1 - 0.062$
- Replacing $dW(y)$ by $1 dy$ gives $\Pr(\text{Select pm}) \rightarrow 1 - 0.049$

AFIC curiosity 2

- Focus on the complete $G(y)$, weighted by $W(y) = F(y; \theta_0)$
- Direct comparison between parametrics and nonparametrics reduces to

$$\text{CvM}_n = \int n(\hat{G}_n(y) - F(y; \hat{\theta}))^2 dF(y; \hat{\theta}) \text{ vs. } \kappa$$

- If $G = F \sim N(\xi, \sigma^2)$: $\Pr(\text{Select pm}) \rightarrow 1 - 0.062$
- Replacing $dW(y)$ by $1 dy$ gives $\Pr(\text{Select pm}) \rightarrow 1 - 0.049$

AFIC curiosity 2

- Focus on the complete $G(y)$, weighted by $W(y) = F(y; \theta_0)$
- Direct comparison between parametrics and nonparametrics reduces to

$$\text{CvM}_n = \int n(\hat{G}_n(y) - F(y; \hat{\theta}))^2 dF(y; \hat{\theta}) \text{ vs. } \kappa$$

- If $G = F \sim N(\xi, \sigma^2)$: $\Pr(\text{Select pm}) \rightarrow 1 - 0.062$
- Replacing $dW(y)$ by $1 dy$ gives $\Pr(\text{Select pm}) \rightarrow 1 - 0.049$

AFIC curiosity 2

- Focus on the complete $G(y)$, weighted by $W(y) = F(y; \theta_0)$
- Direct comparison between parametrics and nonparametrics reduces to

$$\text{CvM}_n = \int n(\hat{G}_n(y) - F(y; \hat{\theta}))^2 dF(y; \hat{\theta}) \text{ vs. } \kappa$$

- If $G = F \sim N(\xi, \sigma^2)$: $\Pr(\text{Select pm}) \rightarrow 1 - 0.062$
- Replacing $dW(y)$ by $1 dy$ gives $\Pr(\text{Select pm}) \rightarrow 1 - 0.049$

Illustration: Egyptian Roman era lifelengths

- Data: Lifelengths of 82 men and 59 women - Pearson (1902)
- $\mu_1 = G_{\text{men}}^{-1}(0.5) - G_{\text{women}}^{-1}(0.5)$
- $\mu_2 = G_{\text{men}}^{-1}(0.9) - G_{\text{women}}^{-1}(0.9)$

Illustration: Egyptian Roman era lifelengths

- Data: Lifelengths of 82 men and 59 women - Pearson (1902)
- $\mu_1 = G_{\text{men}}^{-1}(0.5) - G_{\text{women}}^{-1}(0.5)$
- $\mu_2 = G_{\text{men}}^{-1}(0.9) - G_{\text{women}}^{-1}(0.9)$


Illustration: Egyptian Roman era lifelengths

- Data: Lifelengths of 82 men and 59 women - Pearson (1902)
- $\mu_1 = G_{\text{men}}^{-1}(0.5) - G_{\text{women}}^{-1}(0.5)$
- $\mu_2 = G_{\text{men}}^{-1}(0.9) - G_{\text{women}}^{-1}(0.9)$

Illustration: Egyptian Roman era lifelengths

- Data: Lifelengths of 82 men and 59 women - Pearson (1902)
- $\mu_1 = G_{\text{men}}^{-1}(0.5) - G_{\text{women}}^{-1}(0.5)$
- $\mu_2 = G_{\text{men}}^{-1}(0.9) - G_{\text{women}}^{-1}(0.9)$

Model fit MEN



Model fit WOMEN




Illustration: Egyptian Roman era lifelengths




Illustration: Egyptian Roman era lifelengths

