

Modelling the formation of the GD-1 stellar stream inside a host with a fermionic dark matter core-halo distribution.

Martín F. Mestre^{1,2,*}, Carlos R. Argüelles^{1,2}, and Daniel D. Carpintero^{1,2}

¹ Instituto de Astrofísica de La Plata (CONICET-UNLP), Paseo del Bosque S/N, La Plata (1900), Buenos Aires, Argentina

² Facultad de Ciencias Astronómicas y Geofísicas de La Plata (UNLP), Paseo del Bosque S/N, La Plata (1900), Buenos Aires, Argentina

Received February 29, 2023; accepted February 30, 2023

ABSTRACT

Context. Stellar streams are a consequence of the tidal forces produced by a host galaxy on its satellites (i.e. globular clusters and dwarf spheroidals). As the self-gravity of stellar streams is almost negligible, they constitute excellent probes of the gravitational potential of the host galaxy. For this reason, some Milky Way stellar streams have been used to put constraints on the dark matter (DM) total mass and shape, under empirical DM distributions (i.e. NFW, logarithmic, etc). Within the set of non-empirical DM distributions, there exists a DM model deduced from first principles by means of the maximization of the coarse-grained entropy for self-gravitating fermions, the MEPP, after Maximum Entropy Production Principle. MEPP profiles have a rich variety of behaviours, being a large subfamily characterized by a dense core of mass $\approx 4.1 \times 10^6 M_\odot$ (that can mimick a black hole regarding the orbits of the S-stars at Sagittario A*) and an extended halo with the plasticity of a Burkert profile.

Aims. In this work we attempt to model the GD-1 stellar stream using a spherical MEPP distribution which, at the same time, is in sintony with previous fits to the S-stars.

Methods. For that purpose we used a genetic algorithm in order to fit both the stream orbit's initial conditions and the fermionic halo. We modelled the barionic potential with a bulge and two disks (thin and thick) with fixed parameters according to the recent literature. The stream observable is 6D phase-space data from the Gaia DR2 survey.

Results. We were able to find good fits for a 1D continuous subfamily of models parametrized by the fermion mass going from 56 keV/c² to 360 keV/c², respectively corresponding to core radii going from 10³ to 10 Schwarzschild radii. For smaller and larger values of the fermion mass, there is no solution that simultaneously fits the GD-1 stream and the S-stars. The solutions have a virial radius of 28 kpc and a virial mass of $1.4 \times 10^{11} M_\odot$, the latter being at 2σ from previous Milky Way DM halo mass estimates using the Sagittarius stream. We do not assume the velocity of the local standard of rest (v_{lsr}) and the result gives $v_{lsr} \approx 244$ km/s, in agreement with recent independent estimates.

Key words. Galaxy: kinematics and dynamics – Galaxy: stellar streams – dark matter

1. Introduction

Stellar streams probe the acceleration field produced by the Milky Way (MW). This information together with barionic measurements help in making claims about the dark matter (DM) halo.

2. Methodology

In this section we explain the observables and methods used in this research. The exact pipeline applied in order to obtain the results and plots of this paper can be found at the following GitHub repository: https://github.com/martinmestre/stream-fit/blob/main/pipeline_paper/.

2.1. Observables and assumed measurements

The main observables used in this project are materialized by the polynomial fits performed by ? on the GD-1 stream using astrometry (Gaia DR2), photometry and high-

precision spectroscopy datasets together with the analysis of the STREAMFINDER algorithm. Those polinomials are the following:

$$\phi_2 = 0.008367\phi_1^3 - 0.05332\phi_1^2 - 0.07739\phi_1 - 0.02007, \quad (1)$$

$$D = -4.302\phi_1^5 - 11.54\phi_1^4 - 7.161\phi_1^3 + 5.985\phi_1^2 + 8.595\phi_1 + 10.36, \quad (2)$$

$$\tilde{\mu}_\alpha = 3.794\phi_1^3 + 9.467\phi_1^2 + 1.615\phi_1 - 7.844, \quad (3)$$

$$\mu_\delta = -1.225\phi_1^3 + 8.313\phi_1^2 + 18.68\phi_1 - 3.95, \quad (4)$$

$$v_h = 90.68\phi_1^3 + 204.5\phi_1^2 - 254.2\phi_1 - 261.5, \quad (5)$$

$$v_h = 90.68\phi_1^3 + 204.5\phi_1^2 - 254.2\phi_1 - 261.5, \quad (6)$$

with ϕ_1 and ϕ_2 in radians, D in kpc, $\tilde{\mu}_\alpha = \mu_\alpha \cos \delta$ and μ_δ in mas/yr and v_h in km/s. These quantities correspond respectively to longitude and latitude in GD-1 celestial frame of reference (?), heliocentric distance, proper motion in right ascension and declination, and heliocentric radial velocity. The domain of this polinomials is limited to $-90 < \phi_1 [^\circ] < 10$. To have our observable data points we sample the domain with 100 points ($\phi_1^{(i)}$ for $i = 1, \dots, 100$) and evaluate the polinomials there in order to have a complete set of observables ($\phi_1^{(i)}, \phi_2^{(i)}, D^{(i)}, \tilde{\mu}_\alpha^{(i)}, \mu_\delta^{(i)}, v_h^{(i)}$). In one

* e-mail: mmestre@fcaglp.unlp.edu.ar

of the experiments we will consider an observable of a different nature and related to the density concentration at the origin, of size $\lesssim 1$ milliparsec, characteristic of the fermionic halo distribution. By convention, the core radius is defined as the radius when the circular velocity reaches its first maximum. The constraint for the mass of the core of the distribution is assumed to be $m_{\text{core}} = 3.5 \times 10^6 M_{\odot}$ in agreement with ???.

The assumed measurements used in this paper are the galactocentric distance of the sun $R_{\odot} = 8.122$ kpc (?) and the sun's peculiar velocity $v_{\odot p} = (11.1, 12.24, 7.25)$ km/s (?).

2.2. The fermionic DM model

Our DM model consists of a spherical and isotropic distribution of fermions at finite temperature in hydrostatic equilibrium subject to the law of General Relativity (GR), i.e. T.O.V equations, as defined in ? with the particularity that we have made a few change of variables in order to make the numerical computation more robust. The system of equations thus obtained is as follows:

$$\frac{dv}{d\zeta} = \frac{1}{2} \left[e^z + e^{2\zeta} \frac{P(r)}{\rho_{\text{rel}} c^2} \right] [1 - e^z]^{-1}, \quad (7)$$

$$\frac{dz}{d\zeta} = -1 + e^{(2\zeta-z)} \frac{\rho(r)}{\rho_{\text{rel}}}, \quad (8)$$

where

$$\zeta = \ln(r/R), \quad (9)$$

$$z = \ln \psi, \quad \psi = \frac{M_{\text{DM}}(r) R}{M}, \quad (10)$$

$$\rho(r) = \frac{4\rho_{\text{rel}}}{\sqrt{\pi}} \int_1^{\infty} \epsilon^2 [\epsilon^2 - 1]^{1/2} f(r, \epsilon) d\epsilon, \quad (11)$$

$$P(r) = \frac{4\rho_{\text{rel}} c^2}{3\sqrt{\pi}} \int_1^{\infty} [\epsilon^2 - 1]^{3/2} f(r, \epsilon) d\epsilon, \quad (12)$$

$$R^2 = \frac{c^2}{8\pi G \rho_{\text{rel}}}, \quad (13)$$

$$M = 4\pi R^3 \rho_{\text{rel}}, \quad (14)$$

$$\rho_{\text{rel}} = \frac{G\pi^{3/2} m^4 c^3}{h^3}, \quad (15)$$

where G is the Gravitational constant, c is the speed of light, h is the Planck constant, m is the fermion mass, $M_{\text{DM}}(r)$ is DM mass enclosed up to radius r , $\rho(r)$ is the density, $P(r)$ is the pressure and $v(r)$ is the metric exponent, i.e. we use the convention $g_{00} = e^{v(r)}$. The phase-space function (f) is given by a Fermi-Dirac distribution with energy cutoff:

$$f(r, \epsilon) = \begin{cases} \frac{1 - e^{[\epsilon - \epsilon_c(r)]/\beta(r)}}{1 + e^{[\epsilon - \alpha(r)]/\beta(r)}} & \epsilon \leq \epsilon_c(r) \\ 0 & \epsilon > \epsilon_c(r), \end{cases} \quad (16)$$

where $\alpha(r)$ is the chemical potential (including rest mass), $\epsilon_c(r)$ is the cutoff energy (including rest mass) and $\beta(r)$ is the temperature variable.

The above equations should be complemented with two thermodynamic equilibrium conditions given in ? and ? together with the condition of energy conservation along the geodesic given in ?:

$$\frac{1}{\alpha} \frac{d\alpha}{dr} = \frac{1}{\beta} \frac{d\beta}{dr} = \frac{1}{\epsilon_c} \frac{d\epsilon_c}{dr} = -\frac{1}{2} \frac{dv}{dr}. \quad (17)$$

It is not possible to integrate this equations from $r = 0$ because the right hand the change of variables $\zeta(r)$ is divergent at the origin. Nevertheless, it is possible to reach the following approximations for the initial conditions at a value $r_{\text{min}} \gtrsim 0$:

$$v(r_{\text{min}}) \approx \frac{1}{6} \frac{\rho_0}{\rho_{\text{rel}}} \left[\frac{r_{\text{min}}}{R} \right]^2 \equiv \tau, \quad (18)$$

$$\psi(r_{\text{min}}) \approx \frac{1}{3} \frac{\rho_0}{\rho_{\text{rel}}} \left[\frac{r_{\text{min}}}{R} \right]^2 = 2\tau, \quad (19)$$

which implies

$$\frac{r_{\text{min}}}{R} = \sqrt{6\tau \frac{\rho_{\text{rel}}}{\rho_0}}, \quad (20)$$

where $\rho_0 \equiv \rho(0)$. It can be seen that the right hand side of Eq. ?? does not depend on the metric so we can add a constant v_0 to the solution in order to satisfy a condition of continuity with the Schwarzschild metric at the border of the fermion distribution, obtaining:

$$v_0 = 2 \ln \left(\frac{\beta_b}{\beta_0} \sqrt{1 - \psi_b} \right),$$

where ψ_b and β_b are quantities evaluated at the border.

The system of equations thus obtained has four free parameters: m , $\alpha_0 = \alpha(0)$, $\beta_0 = \beta(0)$, and $\epsilon_{c0} = \epsilon_c(0)$ but in practice we will use the following related quantities as parameters: m , β_0 , $\theta_0 = (\alpha_0 - 1)/\beta_0$, and $W_0 = (\epsilon_{c0} - 1)/\beta_0$.

Equations ??-?? are solved using a PYTHON (?) script¹ that makes use of the NumPy (?) and SciPy (?) libraries. We tried all the solvers available in the latter package but the only one that could properly integrate these equations was the LSODA algorithm. We used relative and absolute tolerance parameters respectively given by $\text{rtol} = 5 \times 10^{-14}$ and $\text{atol} = 0$.

2.3. Milky Way and stream models

We model our Galaxy with a combination of the fermionic dark halo recently described, whose parameters will be determined in this work, plus a barionic component fixed and identical to the one in Model I of ?. Here we name this full Galaxy model as Fermionic-MW.

Appart from our model, for some intermediary tasks, we will make use of the Galactic model fitted by ? which consists in the MWPotential2014 model with an axisymmetric NFW profile. This fitted model corresponds to a circular velocity at the position of the sun of $v_c(R_{\odot}) = 244 \pm 4 \text{ km s}^{-1}$ and a z -flattening of the DM density distribution of $q_{\rho} = 0.82^{+0.25}_{-0.13}$. Here we name this Galaxy model as NFW-MW.

As GD-1 is a dynamically cold stream, with its stars keeping a large degree of correlation, its almost one-dimensional distribution in phase-space can be well approximated with the orbit of its progenitor. Another facts in this direction, are that there has not been any observation of the tidal arms feature nor of the progenitor's position (????).

In the code, the orbit model is computed starting from initial conditions in spherical equatorial coordinates, ICRS frame: α (RA), δ (declination), D , μ_{α} , μ_{δ} , v_h . The code uses the Astropy ecosystem (???) in order to transform the initial condition to galactocentric coordinates assuming a Galactocentric reference frame with the sun at the position $\mathbf{x}_{\odot} = (-R_{\odot}, 0, 0)$ and a sun's velocity given by the sum of the circular velocity at the position of the sun and the sun's peculiar velocity:

¹ model_def.py

$v_\odot = (11.1, v_c(R_\odot) + 12.24, 7.25)$. The circular velocity is dependent on the model and the position and given by

$$\begin{aligned} v_c^2(R_\odot) &= R_\odot \|\nabla\Phi(\mathbf{x})\|_{\mathbf{x}=\mathbf{x}_\odot} \\ &= R_\odot \|\nabla\Phi_B(\mathbf{x})\|_{\mathbf{x}=\mathbf{x}_\odot} + G \frac{M_{\text{DM}}(R_\odot)}{R_\odot}, \end{aligned} \quad (21)$$

where Φ is the total potential, Φ_B is the potential generated by the three barionic components and $M_{\text{DM}}(r)$ was defined in Eq. (??). We have used the spherical symmetry property of the DM distribution in order to relate the acceleration with the enclosed mass.

A second step is to integrate the orbit forwards and backwards in time during a time interval of $\Delta t = 0.2$ Gyr, starting in both cases from a given initial condition for the progenitor. In the next sections we will explain how this initial condition was chosen for some simulations and fitted for others. The integrator used is a Runge-Kutta of order eight (DOP853 called from *SciPy*'s `solve_ivp` function) with relative and absolute tolerance parameters respectively given by $\text{rtol} = 5 \times 10^{-14}$ and $\text{atol} = 0.5 \times 10^{-14}$.

After having the stream's orbit integrated, we transform the orbit to both the ICRS and the GD-1 frame of coordinates. For the latter we used the `GD1Koposov10` class defined in the *Gala* package (??) which uses the transformation matrix defined by ?. After these two transformation we obtain the orbit expressed in the observable variables $(\phi_1, \phi_2, D, \tilde{\mu}_\alpha, \mu_\delta, v_h)$ used in Eqs. (??)-(??). Finally, we build interpolators of these variables as a function of ϕ_1 that will be used, together with the observed data defined in Sec. ??, to evaluate the following χ^2 stream function:

$$\chi_{\text{stream}}^2 = \chi_{\phi_2}^2 + \chi_D^2 + \chi_{\tilde{\mu}_\alpha}^2 + \chi_{\mu_\delta}^2 + \chi_{v_h}^2, \quad (22)$$

$$\chi_\eta^2 = \frac{1}{\sigma_\eta^2} \sum_{i=1}^{100} (\eta^{(i)} - \eta(\phi_1^{(i)}))^2, \quad (23)$$

where η represents each of the four corresponding independent variables $(\phi_2, D, v_h, \tilde{\mu}_\alpha, \mu_\delta)$ and σ_η is the average dispersion of the stream data points taken by ocular inspection from Figs. 1,3,4 of ?: $\sigma_{\phi_2} = 0.5^\circ$, $\sigma_D = 1.5$ kpc, $\sigma_{v_h} = 10$ km/s and $\sigma_{\tilde{\mu}_\alpha} = \sigma_\delta = 2$ mas/yr. Thus χ_{stream}^2 measures the departure of the model from the stream observations. For some fits we will also consider the departure of the model from a dark mass constraint in the core of the distribution:

$$\chi_{\text{core}}^2 = \frac{(m - m_{\text{core}})^2}{\sigma_m^2}, \quad (24)$$

where m_{core} was defined in Sec. ?? and σ_m was tested between 10% and 50% of m_{core} , so sometimes we will use the following compound function:

$$\chi_{\text{full}}^2 = \chi_{\text{stream}}^2 + \chi_{\text{core}}^2. \quad (25)$$

Note that in order to compute the estimated core mass of each model, m in Eq.(??), we must find the first local maximum of the expression for the circular velocity in GR:

$$v_{\text{DM}}(r) = c \sqrt{\frac{r}{2} \frac{dv(r)}{dr}}. \quad (26)$$

2.4. Optimization algorithms

Our objective consists in fitting our MW model by minimizing the χ^2 function given by Eqs. (??,??). In order

to do so we use two optimization algorithms for different purposes that will be explained in Sec. ???. One of them is *SciPy*'s `optimize.differential_evolution` algorithm. This genetic architecture was used with metaparameters given by `strategy="best2bin"`, `maxiter=200`, `popsiz=200`, `tol=5 \times 10^{-8}` and `atol=0`. This method can be run in parallel with shared memory, so a convergence test was made in a cluster's node with 24 processors, varying the values of the metaparameters (e.g. halving and doubling the values of `maxiter` and `popsiz`, setting `strategy="best1bin"` and combinations).

The other algorithm is `LN_NELDERMEAD` (???) from the *NLOpt.jl* (?) package for the *JULIA* (?) language. We used default values for the all its metaparameters except for `reltol=5 \times 10^{-5}`.

3. Results and discussion

Initially, we searched for a good enough initial condition (IC) of the progenitor orbit for our Fermionic-MW. For that, we fitted the initial conditions in the (fixed) NFW-MW model defined in Sec. ?? by using the χ_{stream}^2 function defined in Eq. (??) and the genetic algorithm specified in Sec. ???. The genetic algorithm needs to be provided with bounds for the variables so we used boxes with a centre close to the middle point of the observable data, i.e. $\eta^{(51)}$ that corresponds to $\alpha = 149.24^\circ$, $\delta = 36.61^\circ$, $D = 7.69$ kpc, $\tilde{\mu}_\alpha = -5.70$ mas/yr, $\mu_\delta = -12.48$ mas/yr and $v_h = -18.81$ km/s, and with generous sides of length equal to their absolute values (except for α where we used a side of length $\alpha/5$).

It should be said that the NFW-MW model was fitted by ? with a set of Gaia DR2 data different from the polynomials used here. Any point along the stream is equally good as an IC. The genetic algorithm converged to a solution given approximately by $\alpha = 149.17^\circ$, $\delta = 36.51^\circ$, $D = 7.95$ kpc, $\tilde{\mu}_\alpha = -6.94$ mas/yr, $\mu_\delta = -12.49$ mas/yr and $v_h = -18.33$ km/s with a value of $\chi_{\text{stream}}^2 \approx 17.52$. The orbit that corresponds to this IC is displayed in the observable space in Fig. ?? (top: ϕ_2 , bottom: D) and Fig. ?? (top: $\tilde{\mu}_\alpha$, middle: μ_δ , bottom: v_h) with a dashed yellow line. The thick solid black line shows the corresponding observable η data while the thin solid ones demarkate the corresponding regions at $1\sigma_\eta$.

From previous works (???) we know that the fermionic DM model parameters can be approximately grouped in two sets with different functionality. One group is formed by ϵ and β_0 that control the core of the distribution in the sense that for given values of m_{core} (Sec. ??) and ϵ , it is possible to find a value of β_0 in agreement; with a very small influence of the other two parameters: θ_0 and W_0 . This implies a partial degeneracy between ϵ and β_0 . The other group is formed by θ_0 and W_0 , which have small effect on the core region but determine the distribution behaviour in the halo region. Additionally, there is also an approximate degeneracy between these two variables in the sense that it is the parameter $\omega_0 = W_0 - \theta_0$ which mainly controls the halo profile, as will be further shown in this paper.

As a starting point for our three-step multiparameter search, we took the result from ? who found that for $(\epsilon, \beta_0, \theta_0, W_0) = (56, 1.1977 \times 10^{-5}, 37.765, 66.3407)$ their MW model (DM+barions) satisfies both the geodesic motion of S2 and G2 at Sagittarius A*, i.e. $m_{\text{core}} = 3.5 \times 10^5 M_\odot$, and the rotation curve from ?. As a first step, we fixed values for $\epsilon = 56$ and fixed the IC that was recently fitted for the NFW-MW model, in order to perform a genetic minimization of the χ_{full}^2 function in the window: $(\theta_0, \omega_0, \beta_0) \in [35, 40] \times [25, 30] \times [1.2 \times 10^{-5}, 1.3 \times 10^{-5}]$. The algorithm converged to $(\theta_0, \omega_0, \beta_0) \approx (36.06, 27.34, 1.258 \times 10^{-5})$ with a value of $\chi_{\text{stream}}^2 \approx 21.96$ and

$\chi_{\text{core}}^2 \approx 6.05 \times 10^{-6}$. Finally, the third step consisted in fixing the Fermionic-MW parameters while floating the ICs of the orbit with the genetic scheme, which gave us very similar to the FMW-MW case: $\alpha = 148.88^\circ$, $\delta = 36.34^\circ$, $D = 7.96$ kpc, $\tilde{\mu}_\alpha = -6.87$ mas/yr, $\mu_\delta = -12.49$ mas/yr and $v_h = -16.05$ km/s. The corresponding orbit is displayed in the observable space in Figs. ??-?? with a dotted green line where it can be noticed that both the Fermionic-MW and NFW-MW models fit the GD-1 stream very well. We have not performed any statistically rigorous comparison between both models to determine which one is more consistent with the data. In fact, here we used a fixed NFW-MW model that was fitted by ? with another set of Gaia DR2 observations, as has previously been stated.

We would like to give a glimpse of how badly behaved is the χ_{stream}^2 function so we have evaluated it for fixed values of $(\epsilon, \beta_0) = (56, 1.25 \times 10^{-5})$ in a bidimensional grid spanning the box $(\theta_0, \omega_0) \in [35.25, 37.25] \times [26.5, 28.5]$. Fig. ?? shows this grid colored by the value of their corresponding χ_{stream}^2 .

4. Conclusions

Acknowledgements. Thank to people, institutions and codes used. This work used computational resources from CCAD – Universidad Nacional de Córdoba (<https://ccad.unc.edu.ar/>), which are part of SNCAD – MinCyT, República Argentina. MFM thanks Juan Ignacio Rodríguez for his great support with hardware and software issues regarding the IALP server and personal computers. The figures presented were made with the *AlgebraOfGraphics.jl* (<https://aog.makie.org/dev/>) and *Makie.jl* (?) packages.

References

- Argüelles, C. R., Krut, A., Rueda, J. A., & Ruffini, R. 2019, *Physics of the Dark Universe*, 24, 100278
- Argüelles, C. R., Mestre, M. F., Becerra-Vergara, E. A., et al. 2022, *MNRAS*, 511, L35
- Argüelles, C. R., Krut, A., Rueda, J. A., & Ruffini, R. 2018, *Physics of the Dark Universe*, 21
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, *apj*, 935, 167
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Becerra-Vergara, E. A., Argüelles, C. R., Krut, A., Rueda, J. A., & Ruffini, R. 2020, *A&A*, 641, A34
- Becerra-Vergara, E. A., Argüelles, C. R., Krut, A., Rueda, J. A., & Ruffini, R. 2021, *MNRAS*, 505, L64
- Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. 2017, *SIAM review*, 59, 65
- Box, M. J. 1965, *The Computer Journal*, 8, 42
- Danisch, S. & Krumbiegel, J. 2021, *Journal of Open Source Software*, 6, 3349
- de Boer, T. J. L., Belokurov, V., Koposov, S. E., et al. 2018, *Monthly Notices of the Royal Astronomical Society*, 477, 1893
- GRAVITY Collaboration, Abuter, R., Amorim, A., et al. 2018, *A&A*, 615, L15
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, *Nature*, 585, 357
- Ibata, R., Thomas, G., Famaey, B., et al. 2020, *The Astrophysical Journal*, 891, 161
- Johnson, S. G. 2007, The NLOpt nonlinear-optimization package, <https://github.com/stevengj/nlopt>
- Klein, O. 1949, *Rev. Mod. Phys.*, 21, 531
- Koposov, S. E., Rix, H.-W., & Hogg, D. W. 2010, *The Astrophysical Journal*, 712, 260
- Krut, A., Argüelles, C. R., Chavanis, P. H., Rueda, J. A., & Ruffini, R. 2023, *ApJ*, 945, 1
- Malhan, K. & Ibata, R. A. 2019, *MNRAS*, 486, 2995
- Malhan, K., Ibata, R. A., Goldman, B., et al. 2018, *Monthly Notices of the Royal Astronomical Society*, 478, 3862
- Merafina, M. & Ruffini, R. 1989, *A&A*, 221, 4
- Nelder, J. A. & Mead, R. 1965, *The Computer Journal*, 7, 308
- Pouliasis, E., Di Matteo, P., & Haywood, M. 2017, *A&A*, 598, A66
- Price-Whelan, A., Sipőcz, B., Lenz, D., et al. 2020, *adrm/gala*: v1.3

- Price-Whelan, A. M. 2017, *The Journal of Open Source Software*, 2
- Price-Whelan, A. M. & Bonaca, A. 2018, *The Astrophysical Journal Letters*, 863, L20
- Schönrich, R., Binney, J., & Dehnen, W. 2010, *Monthly Notices of the Royal Astronomical Society*, 403, 1829
- Shere, K. 1974, *Commun. ACM*, 17, 471
- Sofue, Y. 2013, *Publications of the Astronomical Society of Japan*, 65, 118
- Tolman, R. C. 1930, *Phys. Rev.*, 35, 904
- Van Rossum, G. & Drake Jr, F. L. 1995, *Python tutorial* (Centrum voor Wiskunde en Informatica Amsterdam, The Netherlands)
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature Methods*, 17, 261




Fig. 1. Stream fits in observable space: sky position (top: ϕ_2) and heliocentric distance (bottom: D).




Fig. 2. Stream fits in observable space: proper motion (top: $\tilde{\mu}_\alpha$, middle: μ_δ) and heliocentric velocity (bottom: v_h).

Fig. 3. χ^2_{stream} function for