

ASSIGNMENT 1

Advanced Algorithms and Datastructures

Authors:

Jenny-Margrethe Vej (rwj935)

Martin Gielsgaard Grünbaum (wrk272)

Martin Nicklas Jørgensen (tzk173)

May 12, 2014

1 Exercise 1: *b*-flow

A flow is a *b*-flow if its satisfies the following

$$\sum_{e \in \delta^-(v)} x_e - \sum_{e \in \delta^+(v)} x_e = b_v, \forall v \in V \quad (1)$$

$$0 \leq x_e \leq u_e, \forall e \in E \quad (2)$$

Below we have illustrated the *b*-flows for figure (a). We see that each node satisfies equation 1 and 2, giving us the *b*-flows

$$v_2 v_4 = 2$$

$$v_5 v_1 = 3$$

$$v_5 v_3 = 4$$

which is illustrated in figure 2.




Figure 1: (a)




Figure 2: (a) *b*-flow

In figure (b) we can only satisfy equation 1 and 2 with some of the nodes. Due to the fact that vertex v_4 has no outgoing edges and we do not allow negative flows, we can not fulfil the demand of -2. Because of that, we have no *b*-flow here.




Figure 3: (b)




Figure 4: (b) *b*-flow

2 Exercise 2: An application of MCFP: rectilinear planar embedding

2.1 Exercise 2.1

The x_{vf} values for all vertices and faces of [1, Figure 3] can be found in Table 1. The z_{fg} values

| x_{vf} | a | b | c | d | e |
|----------|-----|-----|-----|-----|-----|
| v_1 | 0 | 1 | 1 | 0 | 0 |
| v_2 | 0 | 0 | 1 | 1 | 0 |
| v_3 | 1 | 0 | 1 | 1 | 1 |
| v_4 | 0 | 0 | 0 | -1 | 1 |
| v_5 | 1 | 0 | 0 | 0 | -1 |
| v_6 | 1 | 1 | 0 | 1 | 1 |
| v_7 | 0 | 0 | 0 | 0 | 0 |

Table 1: x_{vf} -values for all vertex/face combinations.

for the same graph can be found in Table 2. There are a total of 13 breakpoints in [1, Figure 3].

| z_{fg} | a | b | c | d | e |
|----------|-----|-----|-----|-----|-----|
| a | | 0 | 0 | 0 | 0 |
| b | 2 | | 1 | 1 | 0 |
| c | 1 | 1 | | 0 | 0 |
| d | 0 | 1 | 0 | | 2 |
| e | 4 | 0 | 0 | 0 | |

Table 2: z_{fg} values for all faces, values for empty sets are not displayed

This corresponds to the sum of all values of Table 2, as we would expect.

A drawing of a rectilinear layout for [1, Figure 2] can be seen in Figure 5.




Figure 5: Graph in rectilinear form.

2.2 Exercise 2.2

Let B be the set of all boundary cycles and f_e the external boundary cycle. In the following, we make use of the fact that given two boundary cycles x and y , inner turns from x to y (z_{xy}) will be outer turns from y to x (z_{yx}). We do not verify the truth of this statement. The constraints can be expressed as:

$$\forall f \in B \setminus \{f_e\} : \sum_v x_{vf} + \sum_{b \in B \setminus \{f\}} z_{fb} - z_{bf} = 4 \quad (3)$$

$$\sum_v x_{vf_e} + \sum_{b \in B \setminus \{f_e\}} z_{f_e b} - z_{bf_e} = -4 \quad (4)$$

Boundary cycle a is the external boundary cycle, so Eq 4 must hold for a :

$$\begin{aligned} & \sum_v x_{va} + \sum_{b \in B \setminus \{a\}} z_{ab} - z_{ba} \\ &= 3 + z_{ab} - z_{ba} + z_{ac} - z_{ca} + z_{ad} - z_{da} + z_{ae} - z_{ea} \\ &= 3 + 0 - 2 + 0 - 1 + 0 - 0 + 0 - 4 \\ &= 3 - 2 - 1 - 4 = -4 \end{aligned}$$

Boundary cycle e is an internal boundary cycle, so Eq 3 must hold for e specifically:

$$\begin{aligned} & \sum_v x_{vf} + \sum_{b \in B \setminus f} z_{fb} - z_{bf} \\ &= 2 + z_{ea} - z_{ae} + z_{eb} - z_{be} + z_{ec} - z_{ce} + z_{ed} - z_{de} \\ &= 2 + 4 - 0 + 0 - 0 + 0 - 0 + 0 - 2 \\ &= 2 + 4 - 2 = 4 \end{aligned}$$

2.3 Exercise 2.3

The assumption is necessary since only 4 edges can be connected to any single node, when the edges are limited to moving in two directions. From each node, an edge can go either up or down, or left or right.

To show that Equation 1 from [1] holds true we will divide it into the three different cases and show them separately.

- $\sum_f x_{vf} = 0$ if v has degree 2
Any vertex v of degree 2 has 2 edges. This means that v is part of exactly 2 boundary cycles, let's call them f and g . This means that our sum can be defined as

$$\sum_f x_{vf} = x_{vf} + x_{vg}.$$

Since v is of second degree, there is only two possible “kinds” of configurations of the edges, either they form a 180 degree angle, or they form a 90 and a 270 degree angle. In the case of the 180 degree angles we have $x_{vf} = x_{vg} = 0$. The last case is the 90/270 degree case, this

menas one of them will be an innerturn and one will be an outer turn. Because of this one cycle must have an x value of 1 and the other must have an x value of -1 giving a summation of 0, proving the initial statement..

- $\sum_f x_{vf} = 2$ if v has degree 3
Having a degree of 3 means that the edge configuratin can be only one way, 2 edges will go either vertical or horizontal while the last edge is perpendicular to those. This configuration leaves 2 inner turns, and no outer turns. Naming the cycles f, g and h where f and g form inner turns with v , the sum can be written and calculated like this

$$\sum_f x_{vf} = x_{vf} + x_{vg} + x_{vh} = 1 + 1 + 0 = 2$$

which shows the second part of the initial sum is true.

- $\sum_f x_{vf} = 4$ if v has degree 4
If v is uf degree 4, there is exactly 4 edges connected to v and v must be part of 4 cycles. In order to have space for these 4 edges all edges must be inner turns, the sum can then be written out as so

$$\sum_f x_{vf} = 1 + 1 + 1 + 1 = 4$$

This shows that Equation 1 from [1] holds in all three cases for all vertices.

2.4 Exercise 2.4

The objective function $\sum_{f \in B} \sum_{g \in B \setminus \{f\}} z_{fg} + z_{gf}$ expresses the total number of breakpoints, which we wish to minimize.

$$\text{minimize } \sum_{f \in B} \sum_{g \in B \setminus \{f\}} z_{fg} + z_{gf}$$

subject to

$$\forall f \in B \setminus \{f_e\} : \sum_v x_{vf} + \sum_{b \in B \setminus \{f\}} z_{fb} - z_{bf} = 4$$

$$\sum_v x_{vf_e} + \sum_{b \in B \setminus \{f_e\}} z_{f_e b} - z_{b f_e} = -4$$

$$\sum_{u \in V} \sum_g x_{ug} \mod 2 = 0$$

$$\sum_{u \in V} \sum_g x_{ug} \geq 0$$

$$\sum_{u \in V} \sum_g x_{ug} \leq 4$$

$$z_{fg}, z_{gf} \geq 0$$

where the sum \sum_g is over all boundary cycles g containing vertex u .

2.5 Exercise 2.5

We are unsure how to solve this exercise. Vertices and faces of the planar graph would correspond to vertices in an MCFP, such that an edge in the MCFP is either of the form (v, f) (vertex to face) or (f, g) (face to different face). Flow should correspond to some measure of inner turns (e.g. z_{fg} for edges (f, g) and x_{vf} for edges (v, f)). Demands should be of the form $-4 + \sum_v x_{vf}$ for the external face and $4 + \sum_v x_{vf}$ for the internal faces. We are unsure of capacity, except that there is zero capacity between faces which are not adjacent.

3 Exercise 3: Reduction to MCFP

Unfortunately we did not manage to break this exercise, even though we have tried for several hours. If we could get some help with it, we would appreciate it very much, and we would be happy to solve it for the re-handin.

3.1 Exercise 3.1

3.2 Exercise 3.2

3.3 Exercise 3.3

3.4 Exercise 3.4

3.5 Exercise 3.5 (Optional)

References

- [1] Noy Rotbart and Christian Wulff-Nilsen. Minimum-cost flow, advanced algorithms 2014 assignment 1, 2014.