

# Complex Analysis: Homework 13

Martín Prado

November 17, 2024

Universidad de los Andes – Bogotá Colombia

## Exercise 1.

Determine whether the following products converge:

$$(a) \prod_{n=1}^{\infty} \left(1 + \frac{(-1)^n}{n}\right), \quad (b) \prod_{n=1}^{\infty} \left(1 + \frac{(-1)^n}{\sqrt{n}}\right).$$

**Note:** The first term in both products is 0, so we are going to skip and show that the tails from  $n = 2$  forward converge (or diverge) for both cases.

### Solution Item (a)

We are going to prove that  $\sum_{n=2}^{\infty} \log \left(1 + \frac{(-1)^n}{n}\right)$  converges. In fact, if  $M$  is even,

$$\begin{aligned} \sum_{n=2}^M \log \left(1 + \frac{(-1)^n}{n}\right) &= \sum_{k=1}^{M/2} \log \left(\frac{2k+1}{2k}\right) + \log \left(\frac{2k}{2k+1}\right) \\ &= \sum_{k=1}^{M/2} \log \left(\frac{2k+1}{2k} \cdot \frac{2k}{2k+1}\right) \\ &= 0, \end{aligned}$$

and if  $M = 2K + 1$  is odd,

$$\begin{aligned} \sum_{n=2}^M \log \left(1 + \frac{(-1)^n}{n}\right) &= \log \left(\frac{2K}{2K+1}\right) + \sum_{k=1}^{(M-1)/2} \log \left(\frac{2k+1}{2k}\right) + \log \left(\frac{2k}{2k+1}\right) \\ &= \log \left(\frac{2K}{2K+1}\right). \end{aligned}$$

Since  $\log$  is continuous at 1 and  $\frac{2K}{2K+1} \rightarrow 1$  when  $K \rightarrow \infty$ , we conclude that  $\log\left(\frac{2K}{2K+1}\right)$  converges to 0. Since the entire log-series converges to 0, it must follow that

$$\prod_{n=2}^{\infty} \left(1 + \frac{(-1)^n}{n}\right) = \exp \left[ \sum_{n=2}^{\infty} \log \left(1 + \frac{(-1)^n}{n}\right) \right] = 1.$$

### Solution Item (b)

Now, we are going to prove that  $\sum_{n=2}^{\infty} \log \left(1 + \frac{(-1)^n}{\sqrt{n}}\right)$  diverges. In fact, we are going to show that the subsequence  $\sum_{n=1}^M \log \left(1 + \frac{(-1)^n}{\sqrt{n}}\right)$  with  $M = 2k$  diverges to  $-\infty$ . Note that

$$\begin{aligned} \log \left(1 + \frac{1}{\sqrt{2n}}\right) + \log \left(1 - \frac{1}{\sqrt{2n+1}}\right) &= \log \left[ \left(1 + \frac{1}{\sqrt{2n}}\right) \cdot \left(1 - \frac{1}{\sqrt{2n+1}}\right) \right] \\ &= \log \left[ 1 + \frac{\sqrt{2n+1} - \sqrt{2n} - 1}{\sqrt{2n} \cdot \sqrt{2n+1}} \right] \end{aligned}$$

By the limit comparison test  $\log(1+x) = x - O(x^2) \approx x$  for  $x$  near to 0. Then, we can compare the series of  $a_n = \log \left[ 1 + \frac{\sqrt{2n+1} - \sqrt{2n} - 1}{\sqrt{2n} \cdot \sqrt{2n+1}} \right]$  with the series of  $b_n = \frac{\sqrt{2n+1} - \sqrt{2n} - 1}{\sqrt{2n} \cdot \sqrt{2n+1}}$ .

**Remark:** Note that for  $n \geq 1$ ,  $\sqrt{2n+1} - \sqrt{2n} - 1 < 0$ . The limit comparison test only applies if  $a_n, b_n > 0$ , but for our case,

$$a_n = \log \left[ 1 + \frac{\sqrt{2n+1} - \sqrt{2n} - 1}{\sqrt{2n} \cdot \sqrt{2n+1}} \right] < 0 \text{ and } b_n = \frac{\sqrt{2n+1} - \sqrt{2n} - 1}{\sqrt{2n} \cdot \sqrt{2n+1}} < 0$$

and thus, the same argument can be applied to  $-a_n, -b_n > 0$  to conclude that  $-\sum_n a_n$  diverges because  $-\sum_n b_n$  does.

Since  $a_n = \log(1+b_n)$  and  $b_n \rightarrow 0$  when  $n \rightarrow \infty$ , by uniqueness of limit and then L-hôpital rule,

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{x \rightarrow 0} \frac{\log(1+x)}{x} = \lim_{x \rightarrow 0} \frac{1}{1+x} = 1.$$

Then, we can also compare  $b_n$  with  $c_n = \frac{-1}{2n}$  because

$$\begin{aligned} \frac{b_n}{c_n} &= \frac{\sqrt{2n+1} - \sqrt{2n} - 1}{\sqrt{2n} \cdot \sqrt{2n+1}} \cdot (-2n) \\ &= \underbrace{\sqrt{\frac{2n}{2n+1}}}_{\rightarrow 1} + \underbrace{\frac{2n}{\sqrt{2n+1}} - \sqrt{2n}}_{\rightarrow 0} \\ &\rightarrow 1, \quad n \rightarrow \infty \end{aligned}$$

Finally, since  $\sum_{n=1}^{\infty} c_n = \sum_{n=1}^{\infty} \frac{-1}{2n}$  diverges to  $-\infty$ , it follows that  $\sum_{n=1}^{\infty} b_n$  diverges, and thus,  $\sum_{n=1}^{\infty} a_n$  diverges to  $-\infty$  too. Therefore,

$$\begin{aligned} \sum_{n=2}^{2k} \log \left( 1 + \frac{(-1)^n}{\sqrt{n}} \right) &= \sum_{n=1}^k \log \left[ 1 + \frac{\sqrt{2n+1} - \sqrt{2n-1}}{\sqrt{2n} \cdot \sqrt{2n+1}} \right] \\ &= \sum_{n=1}^k a_n \rightarrow -\infty, \quad M \rightarrow \infty. \end{aligned}$$

That implies that  $\prod_{n=2}^{\infty} \left( 1 + \frac{(-1)^n}{\sqrt{n}} \right)$  has a subsequence that diverges to 0.

## Exercise 2.

Prove the class theorem: Let  $(X, d)$  be a compact metric space and let  $g_n : X \rightarrow \mathbb{C}$  be continuous functions such that  $\sum_{n=1}^{\infty} |g_n|$  converges uniformly. Define  $f_n : X \rightarrow \mathbb{C}$  by

$$f_n(x) = \prod_{j=1}^n (1 + g_j(x)).$$

We already know that for every  $x \in X$ , the product  $\prod_{n=1}^{\infty} (1 + g_j(x))$  is absolutely convergent. Then

$$f : X \rightarrow \mathbb{C}, \quad f(x) := \lim_{n \rightarrow \infty} f_n(x)$$

is well-defined.

Show that (a)  $f_n \rightarrow f$  uniformly and (b) that there exists  $N \in \mathbb{N}$  such that for all  $x \in X$ ,

$$f(x) = 0 \iff g_n(x) = -1 \text{ for some } n \leq N.$$

## Solution Item (a)

**Claim 1:** for any real number  $x$ ,  $x + 1 \leq e^x$ .

*Proof:* The function  $F(x) = e^x - x - 1$  has derivative  $F'(x) = e^x - 1$  which has a critical point at  $x = 0$ . The function is convex because  $F''(x) = e^x > 0$  and thus,  $x = 0$  is a global minimum of  $F$ . Since  $F(0) = 0$ , it follows that for any  $x \in \mathbb{R}$ ,  $F(x) \geq F(0) = 0$ , and thus,  $e^x - x - 1 \geq 0$ .

.

□

**Claim 2:** For an absolutely convergent sequence  $(a_n)_{n \in \mathbb{N}}$ , that is  $\sum_{n=1}^{\infty} |a_n| < \infty$ ,

$$\left| \prod_{k=1}^n (1 + a_k) - 1 \right| \leq \prod_{k=1}^n (1 + |a_k|) - 1.$$

*Proof:* By the triangle inequality, any polynomial  $P(a_1, \dots, a_n)$  satisfies

$$|P(a_1, \dots, a_n)| \leq P(|a_1|, \dots, |a_n|).$$

Therefore, by taking the polynomial  $P_n(a_1, \dots, a_n) = \prod_{k=1}^n (1 + a_k) - 1$ , we obtain the desired result.  $\square$

**Claim 3:** For an absolutely convergent sequence  $(a_n)_{n \in \mathbb{N}}$ ,

$$\left| \prod_{k=1}^{\infty} (1 + a_k) - 1 \right| \leq \exp \left( \sum_{k=1}^{\infty} |a_k| \right) - 1.$$

*Proof:* we know that both  $\lim_n \prod_{k=1}^n (1 + a_k)$  and  $\lim_n \sum_{k=1}^n |a_k|$  exist from the hypothesis that  $a_n$  is absolutely convergent. Use **Claim 2** and **Claim 1** to conclude that

$$\left| \prod_{k=1}^n (1 + a_k) - 1 \right| \leq \prod_{k=1}^n (1 + |a_k|) - 1 \leq \exp \left( \sum_{k=1}^n |a_k| \right) - 1.$$

Therefore, after taking limits on both sides we obtain the desired result. In fact, this exact same argument also works on the tails:

$$\left| \prod_{k=N+1}^{\infty} (1 + a_k) - 1 \right| \leq \exp \left( \sum_{k=N+1}^{\infty} |a_k| \right) - 1, \quad \forall N \in \mathbb{N}.$$

Now, note that we can factorize the first  $n$  product terms of  $f_n$  from  $f$

$$f(z) - f_n(z) = \left( \frac{f(z)}{f_n(z)} - 1 \right) \cdot f_n(z) = \underbrace{\left( \prod_{j=n+1}^{\infty} (1 + g_j(z)) - 1 \right)}_{(1)} \cdot \underbrace{\left( \prod_{j=1}^n (1 + g_j(z)) \right)}_{(2)}.$$

For (1) apply **Claim 3** to conclude that

$$\left| \prod_{j=n+1}^{\infty} (1 + g_j(z)) - 1 \right| \leq \exp \left( \sum_{j=n+1}^{\infty} |g_j(z)| \right) - 1.$$

Since  $\sum_{k=n+1}^{\infty} |g_j(z)|$  converges uniformly to 0 (tail of a uniformly convergent sequence), for any  $\varepsilon > 0$ , there exists  $N \in \mathbb{N}$  such that  $\sum_{k=n+1}^{\infty} |g_j(z)| < \varepsilon$  for every  $z \in X$ ,  $n \geq N$ , and thus,

$$\exp \left( \sum_{k=n+1}^{\infty} |g_j(z)| \right) - 1 < \underbrace{e^{\varepsilon} - 1}_{\approx 0} \quad \forall z \in X, \quad n \geq N.$$

so it follows that  $\prod_{j=n+1}^{\infty} (1 + g_j(z)) - 1$  converges uniformly to 0.

On the other hand, for (2), since  $h := \sum_{j=1}^{\infty} |g_j|$  is the uniform limit of continuous functions on a compact set, it follows that there exists  $M > 0$  such that  $h(z) < M$  for every  $z \in X$ . In fact, since  $|g_j| \geq 0$ , it follows that the sequence  $h_n := \sum_{j=1}^n |g_j|$  is increasing and  $h_n \leq h < M$ . Then, by **Claim 1**

$$\begin{aligned} \left| \prod_{j=1}^n 1 + g_j(z) \right| &= \prod_{j=1}^n |1 + g_j(z)| \\ &\leq \prod_{j=1}^n 1 + |g_j(z)| \\ &\leq \exp \left( \sum_{j=1}^n |g_j(z)| \right) \\ &< e^M \quad \forall z \in X. \end{aligned}$$

Finally, (1) converges uniformly to 0 and (2) is uniformly bounded, so it follows that  $|f_n(z) - f(z)|$  converges uniformly to 0.

### Solution Item (b)

Since  $X$  is compact and  $f$  is holomorphic, the number of zeros of  $f$  is finite, otherwise, the set of zeros would have an accumulation point on  $X$ . If  $\{z_1, \dots, z_p\}$  is the set of zeros of  $f$ , the goal is to find  $\forall j \leq p$ ,  $N_j$  for which  $g_{N_j}(z_j) = -1$  and then take  $N = \max_j N_j$ .

Let  $z$  be one of those zeros and assume for the sake of contradiction that  $g_j(z) \neq -1$  for every  $j \in \mathbb{N}$ . Then, we would obtain a contradiction with the fact that for a convergent product  $\prod_{j=1}^{\infty} a_j$  to be zero, one of the elements in the sequence is zero (otherwise, the product must diverge to 0), so let  $a_j = (1 + g_j(z))$  to obtain the contradiction. Therefore, for every  $j$ , there exists  $N_j \in \mathbb{N}$  for which  $g_{N_j}(z_j) = -1$ .

If  $g_n(z) = -1$  for some  $n \leq N$ , then it's clear from the pointwise convergence that  $z$  is a zero of  $f$ :

$$f(z) = (1 - g_n(z)) \times \prod_{j \neq n} (1 - g_j(z)).$$

### Exercise 3.

Let  $U \subset \mathbb{C}$  be open and let  $g_n : U \rightarrow \mathbb{C}$  be holomorphic functions such that  $\sum_{n=1}^{\infty} |g_n|$  converges compactly in  $U$ . Define

$$f_n(x) = \prod_{j=1}^n (1 + g_j(x)).$$

(a) Show that  $(f_n)_{n \in \mathbb{N}}$  converges compactly to a holomorphic function  $f : U \rightarrow \mathbb{C}$ .

(b) Let  $z_0 \in U$ . Show that  $f(z_0) = 0$  if and only if there exists  $j \in \mathbb{N}$  such that  $g_j(z_0) = -1$ , that there are finitely many such  $j$ , and that the order of the zero  $z_0$  for  $f$  is equal to the sum of the multiplicities of  $z_0$  as a zero of all the functions  $1 + g_j$ .

### Solution Part (a)

Let  $K \subseteq U$  be a compact set. Then,  $\sum_{n=1}^{\infty} |g_n|$  converges uniformly in  $K$ , so apply exercise 2 to conclude that  $f_n$  converges uniformly to a function  $f_K$  in  $K$ . By Weierstrass theorem,  $f_K$  must be holomorphic.

Now for any  $z \in U$  let  $K_z \subset U$  be a compact set that contains  $z$ . Define  $f(z) = f_{K_z}(z)$ .  $f$  is well defined because if we take another compact set  $K'_z$  that contains  $z$ , then by uniqueness of limit,  $f_{K_z}(w) = f_{K_z \cap K'_z}(w) = f_{K'_z}(w)$  for every  $w \in K_z \cap K'_z$ .

Fix  $z \in U$ . We want to show that there exists a neighborhood of  $z$  for which  $f$  is holomorphic. Since the choosing of the compact set  $K_z$  doesn't affect the value of  $f(z)$ , let  $\varepsilon > 0$  such that  $\overline{B_\varepsilon(z)} \subset U$  and let  $K_w = \overline{B_\varepsilon(z)}$  for every  $w \in B_\varepsilon(z)$ . With this choosing of  $K_w = K_z$  we get that  $f(w) = f_{K_z}(w)$ , and since  $f_{K_z}$  is holomorphic, it follows that  $f$  is holomorphic at  $B_\varepsilon(z)$ . Thus,  $f$  is holomorphic.

Finally,  $f_n$  converges compactly to  $f$  because for any compact set  $K$ ,  $f_n$  converges uniformly to  $f_K$ , and again, by uniqueness of limit,  $f(z) = f_K(z)$  for every  $z \in K$ .

### Solution Part (b)

If  $f(z_0) = 0$  for  $z_0 \in U$ , then there exists  $j \in \mathbb{N}$  for which  $1 + g_j(z_0) = 0$ , otherwise we would get the same contradiction we formulated at exercise 2(b).

If for some reason there exists an infinite number of  $j \in \mathbb{N}$  for which  $1 + g_j(z_0) = 0$ , then for every  $n \in \mathbb{N}$ , the tail  $\prod_{j=n+1}^{\infty} (1 + g_j(z_0)) = 0 \not\rightarrow 1$  (doesn't converge to 1), so the product doesn't converge to  $f$  according to the definition.

Now, let  $J = \{j_1, \dots, j_q\}$  be the set of indices for which  $g_{j_i}(z_0) = -1$ . Then, for each of this indices, we can factorize the zeros of order say  $m_i$  to obtain  $1 + g_{j_i}(z) = (z - z_0)^{m_i} \cdot h_i(z)$  for some holomorphic function that doesn't vanish at  $z_0$ . Then,

$$\begin{aligned} f(z) &= \prod_{j \in J} (1 + g_j(z)) \times \prod_{j \notin J} (1 + g_j(z)) \\ &= (z - z_0)^{m_1 + \dots + m_q} \cdot \prod_{i=1}^q h_i(z) \times \prod_{j \in \mathbb{N} \setminus J} (1 + g_j(z)). \end{aligned}$$

From the definition of  $h_i$ , we know that  $\prod_{i=1}^q h_i(z_0) \neq 0$  and we defined  $J$  in such way that there are no indices outside of  $J$  for which  $1 + g_j(z_0) = 0$ . Therefore,  $f$  has a zero of order  $m_1 + \dots + m_q$  at  $z_0$ .

### Exercise 4.

Let  $U \subset \mathbb{C}$  be a region, let  $f_n : U \rightarrow \mathbb{C}$  be holomorphic functions, and assume that  $\prod_{j=1}^{\infty} f_j$  converges absolutely and compactly in  $U$ . Show that

$$\frac{f'}{f} = \sum_{j=1}^{\infty} \frac{f'_j}{f_j}$$

where the sum on the right side converges compactly in its domain.

### Solution

$f$  is holomorphic because is the compact limit of holomorphic functions (Weierstrass theorem).

Let  $g_n = \prod_{j=1}^n f_j$ , since  $(g_n)_{n \in \mathbb{N}}$  is a sequence of holomorphic functions that converges compactly to  $f$ , by Weierstrass' theorem, the sequence of derivatives  $(g'_n)_{n \in \mathbb{N}}$  also converge compactly to  $f'$ . Now, by the product rule

$$\begin{aligned} \frac{g'_n}{g_n} &= \frac{\sum_{k=1}^n f'_k \times \prod_{j \neq k}^n f_j}{\prod_{j=1}^n f_j} \\ &= \sum_{k=1}^n f'_k \times \frac{\prod_{j \neq k}^n f_j}{\prod_{j=1}^n f_j} \\ &= \sum_{k=1}^n \frac{f'_k}{f_k}. \end{aligned}$$

Then domain of  $\frac{f'}{f}$  is  $U_0$  which is equal to  $U$  excluding the zeros of  $f$  (which are isolated). For every  $n \in \mathbb{N}$ ,  $g_n$  cannot have more zeros than  $f$ , otherwise there would be a contradiction with

$$\underbrace{f(z)}_{\neq 0} = \underbrace{g_n(z)}_{=0} \times \underbrace{\prod_{j=n+1}^{\infty} f_j(z)}_{< \infty}.$$

Finally,  $\frac{g'_n}{g_n}$  is defined on  $U_0$ , and in every compact set of  $U_0$ ,  $\frac{1}{g_n}$  converges compactly to  $\frac{1}{f}$  because:

For any  $K \subset U_0$ , there exists a constant  $M > 0$  for which  $|g_n(z)|, |f(z)| < M$  for every  $z \in K$ , and thus,

$$\sup_{z \in K} \left| \frac{1}{g_n(z)} - \frac{1}{f(z)} \right| = \sup_{z \in K} \left| \frac{g_n(z) - f(z)}{g_n(z)f(z)} \right| < \frac{1}{M^2} \cdot \sup_{z \in K} |g_n(z) - f(z)| \rightarrow 0.$$

Therefore,  $\frac{g'_n}{g_n}$  converges compactly to  $\frac{f'}{f}$ .