

Exercise 2, TFY4235 Computational physics

Martin a. Johnsrud

Introduction

This report documents the simulation of magnons, as described in [1].

Theory

The Hamiltonian in question is, in units of the coupling constant J ,

$$\mathcal{H}(S; d_z, a, B) = -\frac{1}{2} \sum_{\langle i, j \rangle, a} S_{i,a} S_{j,a} - d_z \sum_j (S_{j,3})^2 - \sum_{j,a} B_{j,a} S_{j,a}.$$

S are the spins, the dynamical variables, $i \in \{1, \dots, N\}$ is the site index, a is vector component index. d_z , a and $B_{i,a}$ are respectively the inisotropy strength, the coupling strength, the magnetic moment and the external magnetic field. The magnetic field is given in units of $\mu/J[B_0]$, where μ magnetic moment. These appear as parameters in the Hamiltonian. The dynamics of the system at zero temprature is governd by the Landau-Lifshitz-Gilbert equation,

$$\begin{aligned} \frac{d}{dt} S_{j,a} &= -\frac{1}{(1 + \alpha^2)} \left[\sum_{bc} \varepsilon_{abc} S_{j,b} H_{j,c} + \alpha \sum_b (S_{j,b} S_{j,b} H_{j,a} - S_{j,b} H_{j,b} S_{j,a}) \right], \\ H_{k,b} &= -\frac{\partial \mathcal{H}}{\partial S_{k,b}} = \frac{1}{2} \sum_{\langle i, j \rangle, a} (S_{i,a} \delta_{j,k} \delta_{a,b} + S_{j,a} \delta_{i,k} \delta_{a,b}) + 2d_z \sum_j S_{j,3} \delta_{b,3} \delta_{j,k} + \sum_{j,a} B_{j,a} \delta_{k,b}. \end{aligned}$$

The time units are given by γ . The triple product identity $\vec{A} \times (\vec{B} \times \vec{C}) = (\vec{A} \cdot \vec{B})\vec{C} - (\vec{A} \cdot \vec{C})\vec{B}$ has been used for the convinience of implementation. The first sum of the H -term can be written as

$$\frac{1}{2} \sum_{\langle i, j \rangle, a} (S_{i,a} \delta_{j,k} \delta_{a,b} + S_{j,a} \delta_{i,k} \delta_{a,b}) = \frac{1}{2} \sum_{\langle i, j \rangle} (S_{i,b} \delta_{j,k} + S_{j,b} \delta_{i,k}) = \frac{1}{2} \sum_{\langle j, i \rangle} 2S_{i,b} \delta_{j,k} = \sum_{j \in \text{NN}_k} S_{i,b},$$

where NN_k are the set of nearest neghbourss of lattice point k . This gives the expression for the effective field

$$H_{k,b} = \sum_{j \in \text{NN}_k} S_{i,b} + 2d_z S_{j,3} \delta_{j,3} + B_{k,b}.$$

References

- [1] Exercise 2, 2021, tfy4235 computational physics.