

Wave function of the isotropic potential

Martin Johnsrud

October 17, 2019

The Schrödinger equation

The Schrödinger equation is, in all generality

$$i\hbar\partial_t\Psi(t,\vec{r}) = \hat{H}\Psi(t,\vec{r}), \quad (1)$$

where the hamiltonian \hat{H} depends on the problem. This text examines an isotropic potential, where the hamiltonian is

$$\hat{H} = -\frac{\hbar^2}{2m} + \nabla V(r), \quad (2)$$

meaning the potential V only depends on the distance from origo, r . The first step is to use separation of variables, and assume the wave equation can be written in the form

$$\Psi(t,\vec{r}) = \psi(\vec{r})\phi(t).$$

inserting this into (1) we obtain

$$i\hbar\psi(\vec{r})\partial_t\phi(t) = \phi(t)\left(-\frac{\hbar^2}{2m} + \nabla V(r)\right)\psi(\vec{r}).$$

Carelessly dividing the equation with the wave function gives us

$$\frac{1}{\phi(t)}i\hbar\partial_t\phi(t) = \frac{1}{\psi(\vec{r})}\left(-\frac{\hbar^2}{2m} + \nabla V(r)\right)\psi(\vec{r}) = E, \quad (3)$$

where E is a constant with a carefully selected name. We can be certain that each side of the equation is a constant, as one side only depends on t , and the other side on \vec{r} . If they are not constant, we could vary one side, while not changing the other, violating the equality. This is the time independent Schrödinger equation.

The time dependent side

$$\partial_t\phi(t) = \frac{-iE}{\hbar}\phi(t)$$

is solved by recognizing that the only function that is its own derivative, up to a constant, is the exponential function. This yields

$$\phi(t) = \exp(-iEt/\hbar).$$

We could add a constant here, but as it will be multiplied with the solution for $\psi(\vec{r})$, we will let that function contain the constant instead.

The time independent Schrödinger equation

Solving for $\phi(\vec{r})$ is considerably more work. As the problem is spherically symmetric, we will choose spherical coordinates (r, ϕ, θ) . The laplacian in spherical coordinates is¹

$$\nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin^2(\theta)} \left(\frac{\partial^2}{\partial \phi^2} + \sin(\theta) \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial}{\partial \theta} \right) \right),$$

meaning we can split it up in to two parts, one affecting the r coordinate, and ϕ, θ coordinates. Noticing that the similarity between the laplacian and the angular momentum operator, we write

$$\nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) - \frac{1}{r^2 \hbar^2} \hat{L}^2$$

Making use of separation of varibales again, we assume

$$\psi(r, \phi, \theta) = R(r)\Phi(\phi)\Theta(\theta) = R(r)Y(\phi, \theta).$$

Taking the \vec{r} -dependent part of (3),

$$\left(-\frac{\hbar}{2m} \nabla^2 + V(r) \right) \psi(r, \phi, \theta) = E \psi(r, \phi, \theta), \quad (4)$$

we get the time independent Schrödinger equation. Seeing that the hamiltonian have some parts affecting the r coordinate, and something affecting the ϕ, θ coordinates, we can pull the functions through the operators and divide though with $\psi(r, \phi, \theta)$ and obtain

$$\begin{aligned} -\frac{1}{Y(\phi, \theta)} \frac{\hat{L}^2}{2m} Y(\phi, \theta) + \frac{1}{R(r)} \left[\frac{\hbar^2}{2m} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + r^2 (V(r) - E) \right] R(r) \\ = -\frac{1}{Y(\phi, \theta)} \frac{\hat{L}^2}{2m} Y(\phi, \theta) + \frac{1}{R(r)} \hat{H}_r R(r) \end{aligned}$$

This equation now has the dimensions of energy. To make it dimensionless, we introduce the variables

$$q = \frac{r}{a}, \quad \epsilon = \frac{2mE}{\hbar^2}, \quad \nu(q) = \frac{2mV(r/a)}{\hbar^2}, \quad \hat{\Lambda} = \frac{\hat{L}}{\hbar}$$

where a, m is constant choosen from the problem with dimensions length and mass. This makes the partial derivative

$$\frac{\partial}{\partial q} = \frac{\partial r}{\partial q} \frac{\partial}{\partial r} = a \frac{\partial}{\partial r}$$

¹Laplacian

dimensionless as well. This makes TUSL

$$\begin{aligned} -\frac{1}{Y(\phi, \theta)} \hat{\Lambda}^2 Y(\phi, \theta) + \left[\frac{\partial}{\partial q} \left(q^2 \frac{\partial}{\partial q} \right) + q^2 (\nu(q) - \epsilon) \right] Q(q) \\ = -\frac{1}{Y(\phi, \theta)} \hat{\Lambda}^2 Y(\phi, \theta) + \frac{1}{Q(q)} \hat{H}_q Q(q) = 0. \end{aligned} \quad (5)$$

As a mathematical sidenote, we can write this in a more mathematical way

$$\nabla^2 F(r, \phi, \theta) = G(r) F(r, \phi, \theta),$$

with $F = -\hbar/2m\psi$ and $G = E - V$, making these problems mathematically equivalent, the only difference is the physical interpretation.

Spherical harmonics

We will first focus on the angular dependent part of the equation. Writing out the angular momentum operator in (5), we get

$$-\frac{1}{Y(\phi, \theta)} \left[\frac{1}{\sin^2(\theta)} \left(\frac{\partial^2}{\partial \phi^2} + \sin(\theta) \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial}{\partial \theta} \right) \right) \right] Y(\phi, \theta) + \frac{\hat{H}_q Q(q)}{Q(q)} = 0.$$

This means we can use the process of separation of variables again, and get

$$\frac{1}{\Phi} \frac{\partial^2}{\partial \phi^2} \Phi = -\frac{1}{\Theta} \sin(\theta) \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial}{\partial \theta} \right) \Theta - \sin^2(\theta) \frac{\hat{H}_q Q}{Q} = -m^2, \quad (6)$$

where $-m^2$ is the separation constant. First, taking the Φ part of the equation, we get

$$\frac{\partial^2 \Phi}{\partial \phi^2} = -m^2 \Phi,$$

giving us the general solution $\Phi(\phi) = A_1 e^{im\phi} + A_2 e^{-im\phi}$. One of the constant can be absorbed into the Θ -function, so that A_1 can be assumed to be 1, without loss of generality. To have a well defined wave function, we have to impose the boundary condition $\Phi(\phi) = \Phi(\phi + 2\pi)$. From this it follows

$$\begin{aligned} e^{im\phi} + A e^{-im\phi} &= e^{im\phi + i2\pi m} + A e^{-im\phi - 2\pi i m} \\ \implies e^{im\phi} (1 - e^{i2\pi m}) &= A e^{-im\phi} (e^{-i2\pi m} - 1), \quad \forall A \in \mathbb{C} \\ \implies e^{i2\pi m} &= e^{-i2\pi m} = 1 \implies m \in \mathbb{N}, \end{aligned}$$

that is, m is a whole number. Seeing that m can be both positive and negative, we will only use the solution

$$\Phi(\phi) = e^{im\phi}, \quad m \in \mathbb{N}$$

as superpositions of this solution, combined with $\Theta(\theta), R(r)$ covers the whole solution space.

Separating (6) the other way yields

$$-\frac{1}{\Theta} \frac{1}{\sin(\theta)} \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial}{\partial \theta} \right) \Theta(\theta) + \frac{m^2}{\sin^2(\theta)} = \frac{\hat{H}_q Q}{Q} = \lambda$$

Here, we perform the sneaky substitution $\cos(\theta) = x$, giving us

$$\frac{\partial}{\partial \theta} = \frac{\partial x}{\partial \theta} \frac{\partial}{\partial x} = -\sin(\theta) \frac{\partial}{\partial x} = \sqrt{1-x^2} \frac{\partial}{\partial x}$$

This gives us the general legendre differential equation

$$\left[\frac{\partial}{\partial x} \left((1-x^2) \frac{\partial}{\partial x} \right) + l(l+1) - \frac{m^2}{1-x^2} \right] P_m^l(x) = 0, \quad (7)$$

where $\Theta^{l,m}(\theta) = P_m^l(\cos(\theta))$. Solving for $P_m^l(x)$, show in the pdf of legendre functions ¹. This results in

$$P_m^l(x) = \frac{1}{l^2 l!} (1-x^2)^{m/2} \frac{d^{l+m}}{dx^{l+m}} (1-x^2)^l, \quad l \in \mathbb{N}^+, \quad m \in \{-l, \dots, 0, \dots, l\}.$$

The full solution to the spherical problem then becomes

$$Y(\phi, \theta) = C P_m^l(\cos(\theta)) e^{im\phi},$$

known as the spherical harmonics.

The radial equation

The last equation is

$$\left[\frac{\partial}{\partial q} \left(q \frac{\partial}{\partial q} \right) + q^2(\nu(q) - \epsilon) \right] Q(q) = l(l+1)Q(q)$$

rewriting the equation,

$$\left[\frac{1}{q^2} \frac{\partial}{\partial q} \left(q^2 \frac{\partial}{\partial q} \right) - \frac{l(l+1)}{q^2} + \nu(q) \right] Q(q) = \epsilon Q(q)$$

Defining $\rho(q) = qQ(q)$, and seeing that

$$\frac{1}{q} \frac{\partial^2}{\partial q^2} \rho(q) = \frac{1}{q} \frac{\partial}{\partial q} \left(Q(q) + q \frac{\partial}{\partial q} Q(q) \right) = \frac{2}{q} \frac{\partial}{\partial q} Q(q) + \frac{\partial^2}{\partial q^2} Q(q) = \frac{1}{q^2} \frac{\partial}{\partial q} \left(q^2 \frac{\partial}{\partial q} \right)$$

¹https://github.com/martkjoh/KlasMek/blob/master/legendre_functions/legendre_functions.pdf