

Exercise 11 - TFY4345 Classical Mechanics

2020

1 Binding energy of the deuteron


[This is a short one]

Deuteron can be split by gamma rays in a nuclear experiment in the reaction $\gamma + {}^2\text{H} \rightarrow p + n$. Calculate the energy required for the gamma rays for this process to occur, in electron volts. The quantities needed are

- Mass of a proton (${}^1\text{H} = p$): 1.007825u
- Mass of a neutron (n): 1.008665u
- Mass of deuteron (${}^2\text{H}$): 2.014102
- 1 u = 931.5 MeV c^{-2}

2 Frequency shift on a rotating disk

A radioactive ${}^{57}\text{Co}$ element is situated on the periphery of a rotating disk. The peripheral velocity is u . The radiation from the cobalt is received by an observer located in the center of the disk. Let f_0 be the eigenfrequency of the radiation in the inertial system where the element is momentarily at rest. Find the frequency f of the radiation observed by the observer in the center.



3 Fast moving particle in two inertial frames


[Exam Dec. 2019]

We shall consider a particle with rest mass m seen from two different inertial reference systems. In the reference system S , the particle has velocity

$$\mathbf{u} = (u_x, u_y, u_z).$$

The reference system S' is moving along the z axis with a velocity v relative to the system S . In this frame, the particle has the velocity

$$\mathbf{u}' = (u'_x, u'_y, u'_z).$$



Find the explicit relationship between \mathbf{u} and \mathbf{u}' , i.e. derive the transformation that give \mathbf{u}' from the components of \mathbf{u} . [Hint] Start with the relationship between the two frames, the Lorentz transformation. What is the definition of the components u_i, u'_j ?

4 Lorentz transformation of energy and momentum

We continue the analysis of the particle in the last exercise. The relations derived there are Einstein's velocity addition formulas,

$$\begin{aligned}u'_x &= \frac{u_x}{\gamma(1 - vu_z/c^2)} \\u'_y &= \frac{u_y}{\gamma(1 - vu_z/c^2)} \\u'_z &= \frac{u_z - v}{1 - vu_z/c^2}\end{aligned}$$

(The difference between these, and the ones derived in the exercise 10, is that they do not assume the third reference system e.g. the particle moves along the z -axis).

a) Use these to show that

$$\frac{1}{\sqrt{1 - u'^2/c^2}} = \gamma \frac{1 - vu_z/c^2}{\sqrt{1 - u^2/c^2}}.$$

Here, $\gamma = 1/\sqrt{1 - (v/c)^2}$, $u^2 = |\mathbf{u}|^2 = u_x^2 + u_y^2 + u_z^2$ and $u'^2 = |\mathbf{u}'|^2 = u_x'^2 + u_y'^2 + u_z'^2$.

b) The energy E and 3-momentum \mathbf{p} of the particle in the reference system S are

$$E = \frac{mc^2}{\sqrt{1 - (u/c)^2}}, \quad \mathbf{p} = \frac{m\mathbf{u}}{\sqrt{1 - (u/c)^2}},$$

while in the S' system they are

$$E' = \frac{mc^2}{\sqrt{1 - (u'/c)^2}}, \quad \mathbf{p}' = \frac{m\mathbf{u}'}{\sqrt{1 - (u'/c)^2}}.$$

Use the equation derived in a) to find the transformation rule of energy of momentum, i.e. express E' and \mathbf{p}' in terms of E and \mathbf{p} .