

# Chiral Perturbation Theory

Martin Johnsrud

September 30, 2021

## 1 Effective Pion Lagrangian

The technique used in  $\chi$ PT to obtain the effective Lagrangian of the pion relies on a “theorem”, as formulated by Weinberg:

[I]f one writes down the most general possible Lagrangian, including all terms consistent with assumed symmetry principles, and then calculates matrix elements with this Lagrangian to any given order of perturbation theory, the result will simply be the most general possible S-matrix consistent with analyticity, perturbative unitarity, cluster decomposition and the assumed symmetry principles. [1]

In other words, if we write down the most general Lagrange density consistent with symmetries of the underlying theory, it will result in the most general S-matrix consistent with that theory, and important physical assumptions. This leaves a Lagrange density with infinitely many terms, and infinitely many free parameters. To be able to use this theory for anything one must have a method for ordering the terms in order of importance. As described in [2], by rescaling the external momenta  $p_\mu \rightarrow tp_\mu$  and quark masses  $m_i \rightarrow t^2 m_i$ , each term in the Lagrangian obtains a factor  $t^D$ . The Lagrangian is then expanded as  $\mathcal{L} = \sum_D \mathcal{L}_D$ , where  $\mathcal{L}_D$  contains all terms with a factor  $t^D$ .

In our case, the underlying theory is QCD with two quarks, up and down, with mass matrix

$$M = \begin{pmatrix} m_u & 0 \\ 0 & m_d \end{pmatrix}. \quad (1)$$

In the isospin limit,  $m_u = m_d$ , the theory is invariant under global transformations by elements of the group  $G' = \text{SU}(2)_L \times \text{SU}(2)_R \times \text{U}(1)_V$ . All terms involving only pions are trivially invariant under  $\text{U}(1)_V$ , (HVVORFOR?) so we focus on the  $G = \text{SU}(2)_L \times \text{SU}(2)_R$  subgroup. This symmetry is spontaneously broken if the quark field has a non-zero ground state expectation value  $\langle \bar{q}q \rangle$ , leaving only a subgroup  $H = \text{SU}(2)_V \subseteq G$  of symmetry transformations of the vacuum state. The Goldstone manifold  $G/H = \text{SU}(2)_A$  is a three-dimensional Lie group, and therefore results in three (pseudo) Goldstone bosons, the pions. There exists an isomorphism from a subset  $S \subseteq M_1$  of the set of all Goldstone-fields

$$M_1 = \{ \pi_a : \mathcal{M}_4 \longrightarrow \mathbb{R}^3 | \pi_a \text{ smooth} \}$$

close to the ground state, into fields taking values in the Goldstone manifold  $G/H$ . (BEVISE?)(HVA ER ISOMORFISME HER?). The  $\chi$ PT effective Lagrangian will be constructed using this map, through the parametrization

$$\begin{aligned} \Sigma : \mathcal{M}_4 &\longrightarrow \text{SU}(2), \\ x &\longrightarrow \Sigma(x) = A_\alpha(U(x)\Sigma_0 U(x))A_\alpha, \end{aligned} \quad (2)$$

where

$$\Sigma_0 = \mathbb{1}, A_\alpha = \exp\left(\frac{i\alpha}{2}\tau_1\right), U(x) = \exp\left(i\frac{\tau_a\pi_a(x)}{2f}\right).$$

$\tau_a$  are the  $\text{SU}(2)$  generators, i.e. Pauli matrices, as described in Appendix A.  $\pi_a$ , where  $a \in \{1, 2, 3\}$ , are the pion fields. These are real fields, meaning  $\pi_a^\dagger = \pi_a$ .

## Leading order Lagrangian

The leading order Lagrangian in  $\chi$ PT is [2, 3]

$$\mathcal{L}_2 = \frac{f^2}{4} \text{Tr} [\nabla_\mu \Sigma (\nabla^\mu \Sigma)^\dagger] + \frac{f^2}{4} \text{Tr} [\chi^\dagger \Sigma + \Sigma^\dagger \chi]. \quad (3)$$

$\chi$  and  $f$  are the free parameters of the theory.  $f$  is the pion decay constant, while  $\chi = 2B_0M$ . Here,  $M$  is the mass matrix Eq. (1), and  $B_0$  is related to the quark condensate through  $f^2 B_0 = -\langle \bar{u}u \rangle$ . The covariant derivative is defined by

$$\nabla_\mu \Sigma = \partial_\mu \Sigma - i[v_\mu, \Sigma], \quad (\nabla_\mu \Sigma)^\dagger = \partial_\mu \Sigma^\dagger - i[v_\mu, \Sigma^\dagger], \quad v_\mu = \frac{1}{2} \mu_I \delta_\mu^0 \tau_3,$$

where  $\mu_I$  is the isospin chemical potential. To get the series expansion of  $\Sigma$  in powers of  $\pi/f$ , we start by using the fact that  $\tau_a^2 = \mathbb{1}$  to write

$$A_\alpha = \sum_n \frac{1}{n!} \left( \frac{i\alpha}{2} \tau_1 \right)^n = \sum_n \left[ \frac{1}{(2n)!} \left( \frac{i\alpha}{2} \right)^{(2n)} + \frac{\tau_1}{(2n+1)!} \left( \frac{i\alpha}{2} \right)^{(2n+1)} \right] = \mathbb{1} \cos \frac{\alpha}{2} + i\tau_1 \sin \frac{\alpha}{2}. \quad (4)$$

The series expansion of  $U$  is

$$U = \exp \left( \frac{i\pi_a \tau_a}{2f} \right) = 1 + \frac{i\pi_a \tau_a}{2f} + \frac{1}{2} \left( \frac{i\pi_a \tau_a}{2f} \right)^2 + \frac{1}{6} \left( \frac{i\pi_a \tau_a}{2f} \right)^3 + \frac{1}{24} \left( \frac{i\pi_a \tau_a}{2f} \right)^4 + \mathcal{O}((\pi/f)^5),$$

which we use to calculate the expansion of the inner part of  $\Sigma$ , as given in Eq. (2),

$$\begin{aligned} U \Sigma_0 U &= \left( 1 + \frac{i\pi_a \tau_a}{2f} + \frac{1}{2} \left( \frac{i\pi_a \tau_a}{2f} \right)^2 + \frac{1}{6} \left( \frac{i\pi_a \tau_a}{2f} \right)^3 + \frac{1}{24} \left( \frac{i\pi_a \tau_a}{2f} \right)^4 \right) \\ &\times \left( 1 + \frac{i\pi_a \tau_a}{2f} + \frac{1}{2} \left( \frac{i\pi_a \tau_a}{2f} \right)^2 + \frac{1}{6} \left( \frac{i\pi_a \tau_a}{2f} \right)^3 + \frac{1}{24} \left( \frac{i\pi_a \tau_a}{2f} \right)^4 \right) + \mathcal{O}((\pi/f)^5) \\ &= 1 + \frac{i\pi_a \tau_a}{f} + 2 \left( \frac{i\pi_a \tau_a}{2f} \right)^2 + \frac{4}{3} \left( \frac{i\pi_a \tau_a}{2f} \right)^3 + \frac{2}{3} \left( \frac{i\pi_a \tau_a}{2f} \right)^4 + \mathcal{O}((\pi/f)^5). \end{aligned}$$

The symmetry of  $\pi_a \pi_b$  means that

$$(\pi_a \tau_a)^2 = \pi_a \pi_b \frac{1}{2} \{\tau_a, \tau_b\} = \pi_a \pi_a, \quad (\pi_a \tau_a)^3 = \pi_a \pi_a \pi_b \tau_b, \quad (\pi_a \tau_a)^4 = \pi_a \pi_a \pi_b \pi_b.$$

This gives us the expression

$$U \Sigma_0 U = 1 + i \frac{\pi_a \tau_a}{f} - \frac{\pi_a^2}{2f^2} - i \frac{\pi_a^2 \pi_b \tau_b}{6f^3} + \frac{\pi_a^2 \pi_b^2}{24f^4} + \mathcal{O}((\pi/f)^5).$$

We combine this result with Eq. (4) to get an expression for  $\Sigma$  up to  $\mathcal{O}((\pi/f)^5)$

$$\begin{aligned} \Sigma &= \left( \cos \frac{\alpha}{2} + i\tau_1 \sin \frac{\alpha}{2} \right) \left( 1 + i \frac{\pi_a \tau_a}{f} - \frac{\pi_a^2}{2f^2} - i \frac{\pi_a^2 \pi_b \tau_b}{6f^3} + \frac{\pi_a^2 \pi_b^2}{24f^4} \right) \left( \cos \frac{\alpha}{2} + i\tau_1 \sin \frac{\alpha}{2} \right) \\ &= \left( 1 + i \frac{\pi_a \tau_a}{f} - \frac{\pi_a^2}{2f^2} - i \frac{\pi_a^2 \pi_b \tau_b}{6f^3} + \frac{\pi_a^2 \pi_b^2}{24f^4} \right) \cos^2 \frac{\alpha}{2} \\ &\quad - \left( 1 + i \frac{\pi_a}{f} \tau_1 \tau_a \tau_1 - \frac{\pi_a^2}{2f^2} - i \frac{\pi_a^2 \pi_b}{6f^3} \tau_1 \tau_b \tau_1 + \frac{\pi_a^2 \pi_b^2}{24f^4} \right) \sin^2 \frac{\alpha}{2} \\ &\quad + i \left( 2\tau_1 + i \frac{\pi_a}{f} \{\tau_1, \tau_a\} - 2\tau_1 \frac{\pi_a^2}{2f^2} - i \frac{\pi_a^2 \pi_b}{6f^3} \{\tau_1, \tau_b\} + 2\tau_1 \frac{\pi_a^2 \pi_b^2}{24f^4} \right) \sin \frac{\alpha}{2} \cos \frac{\alpha}{2}. \end{aligned}$$

Using trigonometric identities and the commutator,

$$\cos^2 \frac{\alpha}{2} - \sin^2 \frac{\alpha}{2} = \cos \alpha, \quad 2 \cos \frac{\alpha}{2} \sin \frac{\alpha}{2} = \sin \alpha, \quad \tau_1 \tau_a \tau_1 = -\tau_a + 2\delta_{1a} \tau_1,$$

the final expression of  $\Sigma$  to  $\mathcal{O}((\pi/f)^5)$  is

$$\Sigma = \left(1 - \frac{\pi_a^2}{2f^2} + \frac{\pi_a^2 \pi_b^2}{24f^4}\right) (\cos \alpha + i\tau_1 \sin \alpha) + \left(\frac{\pi_a}{f} - \frac{\pi_b^2 \pi_a}{6f^3}\right) \left(i\tau_a - 2i\delta_{a1}\tau_1 \sin^2 \frac{\alpha}{2} - \delta_{a1} \sin \alpha\right). \quad (5)$$

The kinetic term in the  $\chi$ PT Lagrangian is

$$\nabla_\mu \Sigma (\nabla^\mu \Sigma)^\dagger = \partial_\mu \Sigma \partial^\mu \Sigma^\dagger - i (\partial_\mu \Sigma [v^\mu, \Sigma^\dagger] - \text{h.c.}) - [v_\mu, \Sigma] [v_\mu, \Sigma^\dagger]. \quad (6)$$

Using Eq. (5) we find the expansion of the constitutive parts of the kinetic term to be

$$\begin{aligned} \partial_\mu \Sigma &= \left(\frac{-1}{f^2} + \frac{\pi_b^2}{6f^4}\right) (\cos \alpha + i\tau_1 \sin \alpha) (\pi_a \partial_\mu \pi_a) \\ &\quad + \left(\frac{\partial_\mu \pi_a}{f} - \frac{\pi_b^2 \partial_\mu \pi_a + 2\pi_a \pi_b \partial_\mu \pi_b}{6f^3}\right) \left(i\tau_a - 2i\delta_{a1}\tau_1 \sin^2 \frac{\alpha}{2} - \delta_{a1} \sin \alpha\right) \\ &= \left[\left(\frac{-1}{f^2} + \frac{\pi_b^2}{6f^4}\right) (\pi_a \partial_\mu \pi_a) \cos \alpha - \left(\frac{\partial_\mu \pi_1}{f} - \frac{\pi_b^2 \partial_\mu \pi_1 + 2\pi_1 \pi_b \partial_\mu \pi_b}{6f^3}\right) \sin \alpha\right] \\ &\quad - \left[\left(\frac{-1}{f^2} + \frac{\pi_b^2}{6f^4}\right) (\pi_a \partial_\mu \pi_a) \sin \alpha - \left(\frac{\partial_\mu \pi_1}{f} - \frac{\pi_b^2 \partial_\mu \pi_1 + 2\pi_1 \pi_b \partial_\mu \pi_b}{6f^3}\right) 2 \sin^2 \frac{\alpha}{2}\right] i\tau_1 \\ &\quad + \left(\frac{\partial_\mu \pi_a}{f} - \frac{\pi_b^2 \partial_\mu \pi_a + 2\pi_a \pi_b \partial_\mu \pi_b}{6f^3}\right) i\tau_a, \end{aligned} \quad (7)$$

and

$$\begin{aligned} [v_\mu, \Sigma] &= \frac{1}{2} \mu_I \delta_\mu^0 \left[ \left(1 - \frac{\pi_a^2}{2f^2} + \frac{\pi_a^2 \pi_b^2}{24f^4}\right) i \sin \alpha [\tau_3, \tau_1] + \left(\frac{\pi_a}{f} - \frac{\pi_b^2 \pi_a}{6f^3}\right) \left(i [\tau_a, \tau_3] - 2i\delta_{a1} \sin^2 \frac{\alpha}{2} [\tau_3, \tau_1]\right) \right] \\ &= -\mu_I \delta_\mu^0 \left\{ \left(1 - \frac{\pi_a^2}{2f^2} + \frac{\pi_a^2 \pi_b^2}{24f^4}\right) \tau_2 \sin \alpha + \left(\frac{\pi_a}{f} - \frac{\pi_b^2 \pi_a}{6f^3}\right) \left[(\delta_{a1}\tau_2 - \delta_{a2}\tau_1) - 2\delta_{a1}\tau_2 \sin^2 \frac{\alpha}{2}\right] \right\} \\ &= -\mu_I \delta_\mu^0 \left\{ \left[\left(1 - \frac{\pi_a^2}{2f^2} + \frac{\pi_a^2 \pi_b^2}{24f^4}\right) \sin \alpha + \left(\frac{\pi_1}{f} - \frac{\pi_b^2 \pi_1}{6f^3}\right) \cos \alpha\right] \tau_2 - \left(\frac{\pi_2}{f} - \frac{\pi_b^2 \pi_2}{6f^3}\right) \tau_1 \right\}. \end{aligned} \quad (8)$$

Combining Eq. (7) and Eq. (8) gives the following terms <sup>1</sup>

$$\begin{aligned} \text{Tr}\{\partial_\mu \Sigma \partial^\mu \Sigma^\dagger\} &= \frac{2}{f^2} \partial_\mu \pi_a \partial^\mu \pi_a + \frac{2}{3f^4} [(\pi_a \partial_\mu \pi_a)(\pi_b \partial^\mu \pi_b) - (\pi_a \partial_\mu \pi_b)(\pi_b \partial^\mu \pi_a)], \\ -i \text{Tr}\{\partial^\mu \Sigma [v_\mu, \Sigma^\dagger] - \text{h.c.}\} &= 4\mu_I \frac{\partial_0 \pi_2}{f} + 8\mu_I \frac{\pi_3}{3f^3} \sin \alpha (\pi_2 \partial_0 \pi_3 - \pi_3 \partial_0 \pi_2) \sin \alpha \\ &\quad + \left(\frac{4\mu_I}{f^2} \cos \alpha - \frac{8\mu_I \pi_1}{3f^3} \sin \alpha - \frac{4\mu_I \pi_a \pi_a}{3f^4} \cos \alpha\right) (\pi_1 \partial_0 \pi_2 - \pi_2 \partial_0 \pi_1), \\ -\text{Tr}\{[v_\mu, \Sigma] [v^\mu, \Sigma^\dagger]\} &= \mu_I^2 \left[2 \sin^2 \alpha + \left(\frac{2}{f} - \frac{4\pi_a \pi_a}{3f^3}\right) \pi_1 \sin 2\alpha + \left(\frac{2}{f^2} - \frac{2\pi_a \pi_a}{3f^4}\right) \pi_a \pi_b k_{ab}\right], \\ \text{Tr}\{\Sigma + \Sigma^\dagger\} &= 4 \cos \alpha - \frac{4\pi_1}{f} \sin \alpha - \frac{2\pi_a \pi_a}{f^2} \cos \alpha + \frac{2\pi_1 \pi_a \pi_a}{3f^3} \sin \alpha + \frac{(\pi_a \pi_a)^2}{6f^4} \cos \alpha, \end{aligned}$$

where  $k_{ab} = \delta_{a1}\delta_{b1} \cos 2\alpha + \delta_{a2}\delta_{b2} \cos^2 \alpha - \delta_{a3}\delta_{b3} \sin^2 \alpha$ . If we write the Lagrangian as show in Eq. (3) as  $\mathcal{L}_2 = \mathcal{L}_2^{(0)} + \mathcal{L}_2^{(1)} + \mathcal{L}_2^{(2)} + \dots$ , where  $\mathcal{L}_2^{(n)}$  contains all terms of order  $\mathcal{O}((\pi/f)^n)$ , then the result of the series

<sup>1</sup>The scripts used to aid the calculation of the Lagrangian is available at <https://github.com/martkjoh/prosjektoppgave>

expansion is

$$\mathcal{L}_2^{(0)} = f^2 \left( 2B_0 m \cos \alpha + \frac{1}{2} \mu^2 \sin^2 \alpha \right), \quad (9)$$

$$\mathcal{L}_2^{(1)} = f(\mu_I^2 \cos \alpha - 2B_0 m) \pi_1 \sin \alpha + f \mu_I \partial_0 \pi_2 \sin \alpha, \quad (10)$$

$$\mathcal{L}_2^{(2)} = \frac{1}{2} \partial_\mu \pi_a \partial^\mu \pi_a + \mu_I \cos \alpha (\pi_1 \partial_0 \pi_2 - \pi_2 \partial_0 \pi_1) - B_0 m \pi_a \pi_a \cos \alpha + \frac{1}{2} \mu_I^2 \pi_a \pi_b k_{ab}, \quad (11)$$

$$\begin{aligned} \mathcal{L}_2^{(3)} &= \frac{\pi_a \pi_a \pi_1}{6f} (2B_0 m \sin \alpha - 2\mu_I^2 \sin 2\alpha) \\ &\quad - \frac{2\mu_I}{3f} [\pi_1 (\pi_1 \partial_0 \pi_2 - \pi_2 \partial_0 \pi_1) + \pi_3 (\pi_3 \partial_0 \pi_2 - \pi_2 \partial_0 \pi_3)] \sin \alpha, \end{aligned} \quad (12)$$

$$\begin{aligned} \mathcal{L}_2^{(4)} &= \frac{1}{6f^2} \left\{ \frac{1}{2} B_0 m (\pi_a \pi_a)^2 \cos \alpha - [(\pi_a \pi_a)(\partial_\mu \pi_b \partial^\mu \pi_b) - (\pi_a \partial_\mu \pi_a)(\pi_b \partial^\mu \pi_b)] \right\} \\ &\quad - \frac{\mu_I \pi_a \pi_a}{3f^2} \left[ (\pi_1 \partial_0 \pi_2 - \pi_2 \partial_0 \pi_1) \cos \alpha + \frac{1}{2} \mu_I \pi_a \pi_b k_{ab} \right]. \end{aligned} \quad (13)$$

## Equation of motion and redundant terms

Changing the field parametrization that appear in the Lagrangian does not affect any of the physics, as it corresponds to a change of variables in the path integral [2, 4, 5]. However, a change of variables can result in new terms in the Lagrangian. As a result of this, terms that on the face of it appear independent may be redundant. These terms can be eliminated by using the classical equation of motion. In this section we show first the derivation of the equation of motion, then use this result to identify redundant terms which need not be included in the most general Lagrangian.

We derive the equation of motion for the leading order Lagrangian using the principle of least action. Choosing the parametrization  $\Sigma = \exp(i\pi_a \tau_a)$ , a variation  $\pi_a \rightarrow \pi_a + \delta\pi_a$  results in a variation in  $\Sigma$ ,  $\delta\Sigma = i\tau_a \delta\pi_a \Sigma$ . The variation of the leading order action,

$$S_2 = \int_\Omega d^4x \mathcal{L}_2, \quad (14)$$

when varying  $\pi_a$  is

$$\delta S = \int_\Omega dx \frac{f^2}{4} \text{Tr} \{ (\nabla_\mu \delta\Sigma)(\nabla^\mu \Sigma)^\dagger + (\nabla_\mu \Sigma)(\nabla^\mu \delta\Sigma)^\dagger + \chi \delta\Sigma^\dagger + \delta\Sigma \chi^\dagger \}.$$

Using the properties of the covariant derivative to do partial integration, as show in Appendix B, as well as  $\delta(\Sigma\Sigma^\dagger) = (\delta\Sigma)\Sigma^\dagger + \Sigma(\delta\Sigma)^\dagger = 0$ , the variation of the action can be written

$$\begin{aligned} \delta S &= \frac{f^2}{4} \int_\Omega dx \text{Tr} \{ -\delta\Sigma \nabla^2 \Sigma^\dagger + (\nabla^2 \Sigma)(\Sigma^\dagger \delta\Sigma \Sigma^\dagger) - \chi(\Sigma^\dagger \delta\Sigma \Sigma^\dagger) + \delta\Sigma \chi^\dagger \} \\ &= \frac{f^2}{4} \int_\Omega dx \text{Tr} \{ \delta\Sigma \Sigma^\dagger [(\nabla^2 \Sigma)\Sigma^\dagger - \Sigma \nabla^2 \Sigma^\dagger - \chi \Sigma^\dagger + \Sigma \chi^\dagger] \} \\ &= i \frac{f^2}{4} \int_\Omega dx \text{Tr} \{ \tau_a [(\nabla^2 \Sigma)\Sigma^\dagger - \Sigma \nabla^2 \Sigma^\dagger - \chi \Sigma^\dagger + \Sigma \chi^\dagger] \} \delta\pi_a = 0. \end{aligned}$$

As the variation is arbitrary, the equation of motion to leading order is

$$\text{Tr} \{ \tau_a [(\nabla^2 \Sigma)\Sigma^\dagger - \Sigma \nabla^2 \Sigma^\dagger - \chi \Sigma^\dagger + \Sigma \chi^\dagger] \} = 0. \quad (15)$$

This may be rewritten as a matrix equation. Using that

$$\text{Tr} \{ (\nabla_\mu \Sigma) \Sigma^\dagger \} = \text{Tr} \{ i\tau_a (\partial_\mu \pi_a) \Sigma \Sigma^\dagger \} - i \text{Tr} \{ [v_\mu, \Sigma] \Sigma^\dagger \} = 0,$$

we can see that  $\text{Tr} \{ (\nabla^2 \Sigma) \Sigma^\dagger - \Sigma \nabla^2 \Sigma^\dagger \} = 0$ , and the equation of motion may be written as

$$\mathcal{O}_{\text{EOM}}^{(2)}(\Sigma) = (\nabla^2 \Sigma) \Sigma^\dagger - \Sigma \nabla^2 \Sigma^\dagger - \chi \Sigma^\dagger + \Sigma \chi^\dagger - \frac{1}{2} \text{Tr} \{ \chi \Sigma^\dagger - \Sigma \chi^\dagger \} = 0. \quad (16)$$

The next step in eliminating redundant terms is to change the parametrization of  $\Sigma$  by  $\Sigma(x) \rightarrow \Sigma'(x)$ . Here,  $\Sigma(x) = e^{iS(x)}\Sigma'(x)$ ,  $S(x) \in \mathfrak{su}(2)$ . This change leads to a new Lagrange density,  $\mathcal{L}[\Sigma] = \mathcal{L}[\Sigma'] + \Delta\mathcal{L}[\Sigma']$ . We are free to choose  $S(x)$ , as long  $\Sigma'$  still obeys the required transformation properties. Any terms in the Lagrangian  $\Delta\mathcal{L}$  due to a reparametrization can be neglected, as argued earlier. When demanding that  $\Sigma'$  obey the same symmetries as  $\Sigma$ , the most general transformation to second order in Weinberg's power counting scheme is [2]

$$S_2 = i\alpha_2 [(\nabla^2\Sigma')\Sigma'^\dagger - \Sigma'(\nabla^2\Sigma')^\dagger] + i\alpha_2 \left[ \chi\Sigma'^\dagger - \Sigma'\chi^\dagger - \frac{1}{2} \text{Tr}\{\chi\Sigma'^\dagger - \Sigma'\chi^\dagger\} \right]. \quad (17)$$

$\alpha_1$  and  $\alpha_2$  are arbitrary real numbers. As Eq. (17) is to second order,  $\Delta\mathcal{L}$  is fourth order in Weinberg's power counting scheme. To leading order is given by

$$\begin{aligned} \mathcal{L}_2[e^{iS_2}\Sigma'] &= \frac{f^2}{4} \text{Tr}\{[\nabla_\mu(1+iS_2)\Sigma'][\nabla^\mu\Sigma'^\dagger(1-iS_2)]\} + \frac{f^2}{4} \text{Tr}\{\chi\Sigma'^\dagger(1-iS_2) + (1+iS_2)\Sigma'\chi^\dagger\} \\ &= \mathcal{L}[\Sigma'] + i\frac{f^2}{4} \text{Tr}\{[\nabla_\mu(S_2\Sigma')][\nabla^\mu\Sigma']^\dagger - [\nabla_\mu\Sigma'][\nabla^\mu(\Sigma'^\dagger S_2)]\} - i\frac{f^2}{4} \text{Tr}\{\chi\Sigma'^\dagger S_2 - S_2\Sigma'\chi^\dagger\} \end{aligned}$$

Using the properties of the covariant derivative, as described in Appendix B, we may use the product rule and partial integration to write the difference between the two Lagrangians to fourth order as

$$\begin{aligned} \Delta\mathcal{L}[\Sigma'] &= i\frac{f^2}{4} \text{Tr}\{(\nabla_\mu S_2)(\Sigma'\nabla^\mu\Sigma'^\dagger - (\nabla^\mu\Sigma')\Sigma'^\dagger)\} - i\frac{f^2}{4} \text{Tr}\{\chi\Sigma'^\dagger S_2 - S_2\Sigma'\chi^\dagger\} \\ &= i\frac{f^2}{4} \text{Tr}\{S_2[\Sigma'^\dagger\nabla^2\Sigma' - (\nabla^2\Sigma')\Sigma'^\dagger - \chi\Sigma'^\dagger + \Sigma'\chi^\dagger]\}. \end{aligned}$$

Using the equation of motion Eq. (16), and the fact that  $\text{Tr}\{S_2\} = 0$ , this difference can be written as

$$\Delta\mathcal{L}[\Sigma'] = \frac{f^2}{4} \text{Tr}\{iS_2\mathcal{O}_{\text{EOM}}^{(2)}(\Sigma')\}. \quad (18)$$

Any term that can be written in the form of Eq. (18) for arbitrary  $\alpha_1, \alpha_2 \in \mathbb{R}$  is redundant, as we argued earlier, and may therefore be discarded.  $\Delta\mathcal{L}_2$  is of fourth order, and it can thus be used to remove terms from  $\mathcal{L}_4$  or higher order.

## Next to leading order Lagrangian

The next to leading order Lagrangian density is, assuming no external fields

$$\begin{aligned} \mathcal{L}_4 &= \frac{l_1}{4} \text{Tr}\{\nabla_\mu\Sigma(\nabla^\mu\Sigma)^\dagger\}^2 + \frac{l_2}{4} \text{Tr}\{\nabla_\mu\Sigma(\nabla_\nu\Sigma)^\dagger\} \text{Tr}\{\nabla^\mu\Sigma(\nabla^\nu\Sigma)^\dagger\} + \frac{l_3 + h_1 - h_3}{16} \text{Tr}\{\chi\Sigma^\dagger + \Sigma\chi^\dagger\}^2 \\ &+ \frac{l_4}{4} \text{Tr}\{\nabla_\mu\Sigma(\nabla^\mu\Sigma)^\dagger\} \text{Tr}\{\chi\Sigma^\dagger + \Sigma\chi^\dagger\} + \frac{h_1 - h_3 - l_4 - l_7}{16} \text{Tr}\{\chi\Sigma^\dagger - \Sigma\chi^\dagger\}^2 + \frac{h_1 + h_2 - l_4}{4} \text{Tr}\{\chi\chi^\dagger\} \\ &- \frac{h_1 - h_3 - l_4}{8} \text{Tr}\{(\chi\Sigma^\dagger)^2 + (\Sigma\chi^\dagger)^2\} \end{aligned} \quad (19)$$

To  $\mathcal{L}_4$  to  $\mathcal{O}((\pi/f)^3)$ , we use the result from Eq. (7) and Eq. (8), up to and including  $\mathcal{O}((\pi/f)^2)$ , which gives

$$\begin{aligned} \text{Tr}\{\partial_\mu\Sigma\partial_\nu\Sigma^\dagger\} &= 2\frac{\partial_\mu\pi_a\partial_\nu\pi_a}{f^2} \\ -i\text{Tr}\{\partial_\mu\Sigma[v_\nu, \Sigma^\dagger] - \text{h.c.}\} &= \frac{2\mu_I\pi_2}{f}(\delta_\mu^0\partial_\nu + \delta_\nu^0\partial_\mu)\sin\alpha + \frac{2\mu_I}{f^2}[\pi_1(\delta_\mu^0\partial_\nu + \delta_\nu^0\partial_\mu)\pi_2 - \pi_2(\delta_\mu^0\partial_\nu + \delta_\nu^0\partial_\mu)\pi_1]\cos\alpha \\ -\text{Tr}\{[v_\nu, \Sigma][v_\nu, \Sigma^\dagger]\} &= 2\mu_I^2\delta_\mu^0\delta_\nu^0\left[\sin^2\alpha + \frac{\pi_1}{f}\sin 2\alpha + \frac{\pi_a\pi_b}{f^2}k_{ab}\right]. \end{aligned}$$

Using the form of the Pauli matrices, we can write  $\chi$  as

$$\chi = 2B_0M = 2B_0(\bar{m}\mathbf{1} + \Delta m\tau_3),$$

where  $\bar{m} = (m_u + m_d)/2$ ,  $\Delta m = (m_u - m_d)/2$ , which gives

$$\begin{aligned}\chi\Sigma^\dagger + \Sigma\chi^\dagger &= 4B_0 \left\{ (\bar{m} + \Delta m\tau_3) \left[ \left(1 - \frac{\pi_a^2}{2f^2}\right) \cos\alpha - \frac{\pi_1}{f} \sin\alpha \right] \right. \\ &\quad \left. + \Delta m \left[ \left(1 - \frac{\pi_a^2}{2f^2}\right) \tau_2 \sin\alpha + \frac{\pi_a}{f} (\delta_{a1}\tau_2 \cos\alpha - \delta_{a2}\tau_1) \right] \right\}, \\ \chi\Sigma^\dagger - \Sigma\chi^\dagger &= -4iB_0 \left\{ \bar{m} \left[ \left(1 - \frac{\pi_a^2}{2f^2}\right) \tau_1 \sin\alpha + \frac{\pi_a}{f} \left( \tau_a - 2\delta_{1a}\tau_1 \sin^2 \frac{\alpha}{2} \right) \right] + \Delta m \frac{\pi_3}{f} \right\}.\end{aligned}$$

Combining these results gives all the terms in  $\mathcal{L}_4$ , to  $\mathcal{O}((\pi/f)^3)$ :

$$\begin{aligned}\text{Tr}\{\nabla_\mu\Sigma(\nabla^\mu\Sigma)^\dagger\}^2 &= \text{Tr}\{\partial_\mu\Sigma\partial^\mu\Sigma^\dagger - i(\partial_\mu\Sigma[v^\mu, \Sigma^\dagger] - \text{h.c.}) - [v_\mu, \Sigma][v^\mu, \Sigma^\dagger]\}^2 \\ &= \frac{8\mu_I^2}{f^2}(\partial_\mu\pi_a\partial^\mu\pi_a + 2\partial_\mu\pi_2\partial^\mu\pi_2)\sin^2\alpha \\ &\quad + 16\mu_I^3\left[\frac{\partial_0\pi_2}{f}\sin^3\alpha + \frac{1}{f^2}(3\pi_1\partial_0\pi_2 - \pi_2\partial_0\pi_1)\cos\alpha\sin^2\alpha\right] \\ &\quad + 4\mu_I^4\left\{\sin^4\alpha + 2\sin^2\alpha\left[\frac{\pi_1}{f}\sin 2\alpha + \frac{\pi_a\pi_b}{f^2}(k_{ab} + 2\delta_{a1}\delta_{a2}\cos^2\alpha)\right]\right\},\end{aligned}\tag{20}$$

$$\begin{aligned}\text{Tr}\{\nabla_\mu\Sigma(\nabla_\nu\Sigma)^\dagger\}\text{Tr}\{\nabla^\mu\Sigma(\nabla^\nu\Sigma)^\dagger\} \\ &= \frac{4\mu_I^2}{f^2}(\partial_0\pi_a\partial_0\pi_a + \partial_0\pi_2\partial_0\pi_2 + \partial_\mu\pi_2\partial^\mu\pi_2)\sin^2\alpha \\ &\quad + 16\mu_I^3\left[\frac{\partial_0\pi_2}{f}\sin^3\alpha + \frac{1}{f^2}(3\pi_1\partial_0\pi_2 - \pi_2\partial_0\pi_1)\cos\alpha\sin^2\alpha\right] \\ &\quad + 4\mu_I^4\left\{\sin^4\alpha + 2\sin^2\alpha\left[\frac{\pi_1}{f}\sin 2\alpha + \frac{\pi_a\pi_b}{f^2}(k_{ab} + 2\delta_{a1}\delta_{a2}\cos^2\alpha)\right]\right\},\end{aligned}\tag{21}$$

$$\begin{aligned}\text{Tr}\{\nabla_\mu\Sigma(\nabla^\mu\Sigma)^\dagger\}\text{Tr}\{\chi\Sigma^\dagger + \Sigma\chi^\dagger\} \\ &= 8B_0\bar{m}\left\{2\frac{\partial_\mu\pi_a\partial^\mu\pi_a}{f^2}\cos\alpha + 4\mu_I\left[\frac{\partial_0\pi_2}{2f}\sin 2\alpha + \frac{1}{f^2}(\pi_1\partial_0\pi_2\cos 2\alpha - \pi_2\partial_0\pi_1\cos^2\alpha)\right] \right. \\ &\quad \left. + \mu_I^2\left[2\cos\alpha\sin^2\alpha - 2\frac{\pi_1}{f}\sin\alpha(3\sin^2\alpha - 1) + \frac{1}{f^2}(\pi_1^2[2 - 9\sin^2\alpha] + \pi_2^2[2 - 3\sin^2\alpha] - 3\pi_3^2\sin^2\alpha)\cos\alpha\right]\right\},\end{aligned}\tag{22}$$

$$\text{Tr}\{\chi\Sigma^\dagger + \Sigma\chi^\dagger\}^2 = (8B_0\bar{m})^2\left[\cos^2\alpha - \frac{\pi_1}{f}\sin 2\alpha + \frac{1}{f^2}(\pi_1^2\sin^2\alpha - \pi_a\pi_a\cos^2\alpha)\right],\tag{23}$$

$$\text{Tr}\{\chi\Sigma^\dagger - \Sigma\chi^\dagger\}^2 = -16\left(\frac{2\Delta m B_0\pi_3}{f}\right)^2,\tag{24}$$

$$\begin{aligned}\text{Tr}\{(\chi\Sigma^\dagger)^2 + (\Sigma\chi^\dagger)^2\} \\ &= 16B_0^2\bar{m}^2\left(\cos 2\alpha - 2\frac{\pi_1}{f}\sin 2\alpha - 2\frac{\pi_a\pi_a}{f^2}\cos^2\alpha + 2\frac{\pi_1^2}{f^2}\sin^2\alpha\right) + 16B_0^2\Delta m^2\left(1 - 2\frac{\pi_3^2}{f^2}\right),\end{aligned}\tag{25}$$

$$\text{Tr}\{\chi^\dagger\chi\} = 8B_0^2(\bar{m}^2 + \Delta m^2).\tag{26}$$

## 2 Effective Potential

### Minimizing energy

The value of  $\alpha$  is found by minimizing the free energy. The first approximation to the free energy in the ground state is the static part of the Hamiltonian density  $\mathcal{H}^{(0)}$ , which we get from Eq. (9) through

$$\mathcal{H}_2^{(0)} = -\mathcal{L}_2^{(0)} = -f^2 \left( 2B_0 m \cos \alpha + \frac{1}{2} \mu^2 \sin^2 \alpha \right), \quad (27)$$

The minimum of this function is achieved when

$$\frac{d}{d\alpha} \mathcal{H}_2^{(0)} = f^2 (2B_0 m - \mu_I^2 \cos \alpha) \sin \alpha = 0.$$

This gives the solution set and minimization criterion

$$\alpha = \pi n, \quad n \in \mathbb{Z} \quad \text{or} \quad \cos \alpha = \frac{2B_0 m}{\mu_I^2}. \quad (28)$$

We see that the linear part of the potential from Eq. (10),  $\mathcal{V}^{(1)} = f(\mu_I^2 \cos \alpha - 2B_0 m)\pi_1 \sin \alpha = 0$  if and only if the criterion for minimization is fulfilled.

### Propagator

We may write the quadratic part of the Lagrangian Eq. (11) as <sup>2</sup>

$$\mathcal{L}^{(2)} = \frac{1}{2} \sum_a \partial_\mu \pi_a \partial^\mu \pi_a + \frac{1}{2} m_{12} (\pi_1 \partial_0 \pi_2 - \pi_2 \partial_0 \pi_1) - \frac{1}{2} \sum_a m_a^2 \pi_a^2, \quad (29)$$

where

$$m_1^2 = 2B_0 m \cos \alpha - \mu_I^2 \cos 2\alpha, \quad (30)$$

$$m_2^2 = 2B_0 m \cos \alpha - \mu_I^2 \cos^2 \alpha, \quad (31)$$

$$m_3^2 = 2B_0 m \cos \alpha + \mu_I^2 \sin^2 \alpha, \quad (32)$$

$$m_{12} = 2\mu_I \cos \alpha. \quad (33)$$

The components of the Euler-Lagrange equations of this field are

$$\frac{\partial \mathcal{L}}{\partial \pi_a} = \frac{1}{2} m_{12} (\delta_{a1} \partial_0 \pi_2 - \delta_{a2} \partial_0 \pi_1) - m_a^2 \pi_a, \quad \frac{\partial \mathcal{L}}{\partial (\partial_\mu \pi_a)} = \partial^\mu \pi_a - \frac{1}{2} m_{12} \delta_0^\mu (\delta_{a1} \pi_2 - \delta_{a2} \pi_1).$$

This gives the equation of motion for the field

$$\partial^\mu \partial_\mu \pi_a + m_a^2 \pi_a = m_{12} (\delta_{a1} \partial_0 \pi_2 - \delta_{a2} \partial_0 \pi_1). \quad (34)$$

The propagator of the pion field is defined by

$$[\delta_{ab} (\partial^\mu \partial_\mu + m_a^2) - m_{12} (\delta_{a1} \delta_{b2} - \delta_{a2} \delta_{b1}) \partial_0] D_{bc}(x, x') = -i \delta(x - x') \delta_{ac}. \quad (35)$$

The momentum space propagator, as defined in the Appendix A, fulfills

$$- [\delta_{ab} (p^2 - m_a^2) + i p_0 m_{12} (\delta_{a1} \delta_{b2} - \delta_{a2} \delta_{b1})] \tilde{D}_{bc}(p) := A_{ab} \tilde{D}_{bc}(p) = -i \delta_{ac},$$

where

$$A = - \begin{pmatrix} p^2 - m_1^2 & i p_0 m_{12} & 0 \\ -i p_0 m_{12} & p^2 - m_2^2 & 0 \\ 0 & 0 & p^2 - m_3^2 \end{pmatrix}.$$

---

<sup>2</sup>Summation over isospin index  $(a, b, c)$  will be explicit in this section.

The spectrum of the particles is given by solving  $\det(A) = 0$  for  $p^0$ . With  $p = (p_0, P)$  as the four momentum, this gives

$$\det(A) = A_{33} (A_{11}A_{22} + A_{12}^2) = - (p^2 - m_3^2) [(p^2 - m_1^2)(p^2 - m_2^2) - p_0^2 m_{12}^2] = 0,$$

This equation has the solutions

$$E_0^2 = P^2 + m_2^2, \quad (36)$$

$$E_{\pm}^2 = P^2 + \frac{1}{2} (m_1^2 + m_2^2 + m_{12}^2) \pm \frac{1}{2} \sqrt{4P^2 m_{12}^2 + (m_1^2 + m_2^2 + m_{12}^2)^2 - 4m_1^2 m_2^2}. \quad (37)$$

This gives the effective masses

$$m_0^2 = m_2^2, \quad (38)$$

$$m_{\pm}^2 = \frac{1}{2} [m_1^2 + m_2^2 + m_{12}^2] \pm \frac{1}{2} \sqrt{(m_1^2 + m_2^2 + m_{12}^2)^2 - 4m_1^2 m_2^2}. \quad (39)$$

The propagator may then be obtained as described in Appendix A,

$$\begin{aligned} D = iA^{-1} &= \frac{i}{\det(A)} \begin{pmatrix} A_{22}A_{33} & A_{12}A_{33} & 0 \\ -A_{12}A_{33} & A_{11}A_{33} & 0 \\ 0 & 0 & A_{11}A_{22} + A_{12}^2 \end{pmatrix} \\ &= i \begin{pmatrix} \frac{p^2 - m_2^2}{(p_0^2 - E_+^2)(p_0^2 - E_-^2)} & \frac{-ip_0 m_{12}}{(p_0^2 - E_+^2)(p_0^2 - E_-^2)} & 0 \\ \frac{ip_0 m_{12}}{(p_0^2 - E_+^2)(p_0^2 - E_-^2)} & \frac{p^2 - m_1^2}{(p_0^2 - E_+^2)(p_0^2 - E_-^2)} & 0 \\ 0 & 0 & \frac{1}{p_0^2 - E_0^2} \end{pmatrix}. \end{aligned} \quad (40)$$

### 3 Thermal Field Theory

#### Imaginary-time formalism

In classical mechanics, a thermal system at temperature  $T = 1/\beta$  is described as a ensemble state, which have a probability  $P_n$  of being in state  $n$ , with energy  $E_n$ . In the canonical ensemble, the probability is proportional to  $e^{-\beta E_n}$ . The expectation value of some quantity  $A$ , with value  $A_n$  in state  $n$  is

$$\langle A \rangle = \sum_n A_n P_n = \frac{1}{Z} \sum_n A_n e^{-\beta E_n}, \quad Z = \sum_n e^{-\beta E_n}.$$

$Z$  is the partition function.

In quantum mechanics, an ensemble configuration is described by a non-pure density operator,

$$\hat{\rho} = C \sum_n P_n |n\rangle\langle n|,$$

where  $|n\rangle$  is some basis for the relevant Hilbert space and  $C$  is a constant. Assuming  $|n\rangle$  are energy eigenvectors, i.e.  $\hat{H} |n\rangle = E_n |n\rangle$ , the density operator for the canonical ensemble is

$$\hat{\rho} = C \sum_n e^{-\beta E_n} |n\rangle\langle n| = C e^{-\beta \hat{H}} \sum_n |n\rangle\langle n| = C e^{-\beta \hat{H}}.$$

The expectation value in the ensemble state of a quantity corresponding to the operator  $\hat{A}$  is given by

$$\langle A \rangle = \frac{\text{Tr}\{\hat{\rho}\hat{A}\}}{\text{Tr}\{\hat{\rho}\}} = \frac{1}{Z} \text{Tr}\{\hat{A}e^{-\beta \hat{H}}\} \quad (41)$$



The partition function  $Z$  ensures that the probabilities add up to 1, and is defined as

$$Z = \text{Tr} \left\{ e^{-\beta \hat{H}} \right\}. \quad (42)$$

The grand canonical ensemble takes into account the conserved charges of the system. Conserved charges are a result of Nöther's theorem. Assume we have a set of fields  $\varphi_\alpha$ . Nöther's theorem tells us that if the Lagrangian  $\mathcal{L}[\varphi_\alpha]$  has a *continuous symmetry*, then there is a corresponding conserved current [6, 7]. To define a continuous symmetry of the Lagrangian, we need a one-parameter family of transformations,

$$\varphi_\alpha(x) \longrightarrow \varphi'_\alpha(x; \epsilon) \sim \varphi_\alpha(x) + \epsilon \eta_\alpha(x), \quad \epsilon \rightarrow 0.$$

Here,  $\eta_\alpha(x)$  is some arbitrary function which defines the transformation as  $\epsilon \rightarrow 0$ . Applying this transformation to the Lagrangian will in general change its form,

$$\mathcal{L}[\varphi_\alpha] \rightarrow \mathcal{L}[\varphi'_\alpha] \sim \mathcal{L}[\varphi_\alpha] + \epsilon \delta \mathcal{L}, \quad \epsilon \rightarrow 0.$$

If the change in the Lagrangian can be written as a total derivative, i.e.

$$\delta \mathcal{L} = \partial_\mu K^\mu(x),$$

we say that the Lagrangian has a continuous symmetry. This is because a term of this form will result in a boundary term in the action integral, which does not contribute to the variation of the action. Nöther's theorem states more precisely that the current

$$j^\mu = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \varphi_\alpha)} \eta_\alpha - K^\mu \quad (43)$$

obeys the conservation law

$$\partial_\mu j^\mu = 0. \quad (44)$$

The flux of current through some space-like surface  $V$ , i.e. a surface with a time-like normal vector, defines a conserved charge. This surface is most commonly a surface of constant time in some reference frame. The charge is defined as

$$Q = \int_V d^3x n_\mu j^\mu, \quad \frac{\partial}{\partial t} Q = \nabla \cdot \vec{j}, \quad (45)$$

where  $n^\mu$  is the normal vector of  $V$ . For a surface of constant time,  $n^\mu = (1, 0, 0, 0)$ , and the conserved charged is

$$Q = \int_V dx j^0. \quad (46)$$

In the grand canonical ensemble, a system with  $n$  conserved charges  $Q_i$  has probability proportional to  $e^{-\beta(H - \mu_i Q_i)}$ .  $\mu_i$  are the chemical potentials corresponding to conserved charge  $Q_i$ . This leads to the partition function

$$Z = \text{Tr} \left\{ e^{-\beta(\hat{H} - \mu_i \hat{Q}_i)} \right\}. \quad (47)$$

The partition function may be calculated in a similar way to the path integral approach, in what is called the imaginary-time formalism. This formalism is restricted to time independent problems, and is used to study fields in a volume  $V$ . This volume is taken to infinity in the thermodynamic limit. As an example, take a scalar quantum field theory with the Hamiltonian

$$\hat{H} = \int_V d^3x \hat{\mathcal{H}}[\hat{\varphi}(\vec{x}), \hat{\pi}(\vec{x})], \quad (48)$$

where  $\hat{\varphi}(\vec{x})$  is the field operator, and  $\hat{\pi}(\vec{x})$  is the corresponding canonical momentum operator. These field operators have time independent eigenvectors,  $|\varphi\rangle$  and  $|\pi\rangle$ , defined by

$$\hat{\varphi}(\vec{x}) |\varphi\rangle = \varphi(\vec{x}) |\varphi\rangle, \quad \hat{\pi}(\vec{x}) |\pi\rangle = \pi(\vec{x}) |\pi\rangle. \quad (49)$$

In analogy with regular quantum mechanics, they obey the relations [8]<sup>3</sup>

$$\mathbb{1} = \int \mathcal{D}\varphi(\vec{x}) |\varphi\rangle\langle\varphi| = \int \mathcal{D}\pi(\vec{x}) |\pi\rangle\langle\pi|, \quad (50)$$

$$\langle\varphi|\pi\rangle = \exp\left(i \int_V dx \varphi(\vec{x})\pi(\vec{x})\right), \quad (51)$$

$$\langle\pi_a|\pi_b\rangle = \prod_x \delta(\phi_a(\vec{x}) - \phi_b(\vec{x})), \quad \langle\varphi_a|\varphi_b\rangle = \prod_x \delta(\varphi_a(\vec{x}) - \varphi_b(\vec{x})). \quad (52)$$

The functional integral is defined by starting with  $M$  degrees of freedom,  $\{\varphi_m\}_{m=1}^M$  located at a finite grid  $\{\vec{x}_m\}_{m=1}^M \subset V$ . The integral is then the limit of the integral over all degrees of freedom, as  $M \rightarrow \infty$ :

$$\int \mathcal{D}\varphi(\vec{x}) = \lim_{M \rightarrow \infty} \int \left( \prod_{m=1}^M d\varphi_m \right).$$

The functional Dirac-delta  $\prod_x \delta(f(x))$  is generalization of the familiar Dirac delta function. Given a functional  $\mathcal{F}[f]$ , it is defined by the relation

$$\int \mathcal{D}f(x) \mathcal{F}[f] \prod_x \delta(f(x) - g(x)) = \mathcal{F}[g]. \quad (53)$$

The Hamiltonian is the limit of a sum of Hamiltonians  $\hat{H}_m$  for each point  $\vec{x}_m$

$$\hat{H} = \lim_{M \rightarrow \infty} \sum_{m=1}^M \frac{V}{M} \hat{H}_m(\{\hat{\varphi}_m\}, \{\hat{\pi}_m\}).$$

$H_m$  may depend on the local degrees of freedom  $\hat{\varphi}_m, \hat{\pi}_m$  as well as those at neighboring points. By inserting the completeness relations Eq. (50) into the definition of the partition function, it may be written as

$$Z = \int \mathcal{D}\varphi(\vec{x}) \langle\varphi|e^{-\beta\hat{H}}|\varphi\rangle = \prod_{n=1}^N \left( \int \mathcal{D}\varphi_n(\vec{x}) \int \mathcal{D}\pi_n(\vec{x}) \right) \prod_{n=1}^N \langle\varphi_1|\pi_1\rangle \langle\pi_n|e^{-\epsilon\hat{H}}|\varphi_{n+1}\rangle,$$

where  $\epsilon = \beta/N$  and  $\varphi_1 = \varphi_{n+1}$ . Strictly speaking, we only need to require  $\varphi_1 = e^{i\theta}\varphi_{n+1}$ , as the partition function is only defined up to a constant. As will be shown later, bosons such as the scalar field  $\varphi$ , follow the periodic boundary condition  $\varphi(0, \vec{x}) = \varphi(\beta, \vec{x})$ , i.e.  $e^{i\theta} = 1$ , while fermions follow the anti-periodic boundary condition  $\psi(0, \vec{x}) = -\psi(\beta, \vec{x})$ , i.e.  $e^{i\theta} = -1$ . We now want to exploit the fact that  $|\pi\rangle$  and  $|\varphi\rangle$  are the eigenvectors of the operators that define the Hamiltonian. In our case, as the Hamiltonian density  $\mathcal{H}$  can be written as a sum of functions of  $\varphi$  and  $\pi$  separately,  $\mathcal{H}[\varphi(\vec{x}), \pi(\vec{x})] = \mathcal{F}_1[\varphi(\vec{x})] + \mathcal{F}_2[\pi(\vec{x})]$  we may evaluate it as  $\langle\pi_n|\mathcal{H}[\hat{\varphi}(x), \hat{\pi}(x)]|\varphi_{n+1}\rangle = \mathcal{H}[\varphi_{n+1}(x), \pi_n(x)] \langle\pi_n|\varphi_{n+1}\rangle$ . This relationship does not, however, hold for more general functions of the field operators. In that case, one has to be more careful about the ordering of the operators, for example by using *Weyl ordering* [6]. By series expanding  $e^{-\epsilon\hat{H}}$  and exploiting this relationship, the partition function can be written as, to second order in  $\epsilon$ ,

$$Z = \prod_{n=1}^N \left( \int \mathcal{D}\varphi_n(\vec{x}) \int \mathcal{D}\pi_n(\vec{x}) \right) \exp \left[ -\epsilon \sum_{n=1}^N \int_V d^3x \left( \mathcal{H}(\varphi_n(\vec{x}), \pi_n(\vec{x})) - i\pi_n(\vec{x}) \frac{\varphi_n(\vec{x}) - \varphi_{n+1}(\vec{x})}{\epsilon} \right) \right].$$

We denote  $\varphi_n(\vec{x}) = \varphi(\tau_n, \vec{x})$ ,  $\tau \in [0, \beta]$  and likewise with  $\pi_n(\vec{x})$ . In the limit  $N \rightarrow \infty$ , the expression for the partition function becomes

$$Z = \int_S \mathcal{D}\varphi(\tau, \vec{x}) \int \mathcal{D}\pi(\tau, \vec{x}) \exp \left\{ - \int_0^\beta d\tau \int_V d\vec{x} \{ \mathcal{H}[\varphi(\tau, \vec{x}), \pi(\tau, \vec{x})] - i\pi(\tau, \vec{x}) \dot{\varphi}(\tau, \vec{x}) \} \right\}, \quad (54)$$

where  $S$  is the set of field configurations  $\varphi$  such that  $\varphi(\beta, \vec{x}) = \varphi(0, \vec{x})$ . With a Hamiltonian density of the form  $\mathcal{H} = \frac{1}{2}\pi^2 + \frac{1}{2}(\nabla\varphi)^2 + \mathcal{V}(\varphi)$ , we can evaluate the integral over the canonical momentum  $\pi$  by discretizing

<sup>3</sup>Some authors write  $\mathcal{D}\pi/2\pi$ . This extra factor  $2\pi$  is a convention which in this text is left out for notational clarity.

$$\pi(\tau_n, \vec{x}_m) = \pi_{n,m},$$

$$\begin{aligned} & \int \mathcal{D}\pi \exp \left\{ - \int_0^\beta d\tau \int_V d^3x \left( \frac{1}{2} \pi^2 - i\pi\dot{\varphi} \right) \right\} \\ &= \lim_{M,N \rightarrow \infty} \int \left( \prod_{m,n=1}^{M,N} \frac{d\pi_{m,n}}{2\pi} \right) \exp \left\{ - \sum_{m,n} \frac{V\beta}{MN} \left[ \frac{1}{2} (\pi_{m,n} - i\dot{\varphi}_{m,n})^2 + \frac{1}{2} \dot{\varphi}_{m,n}^2 \right] \right\} \\ &= \lim_{M,N \rightarrow \infty} \left( \frac{MN}{2\pi V\beta} \right)^{MN/2} \exp \left\{ - \int_0^\beta d\tau \int_V d^3x \frac{1}{2} \dot{\varphi}^2 \right\} \end{aligned}$$

where  $\dot{\varphi}_{m,n} = (\varphi_{m,n+1} - \varphi_{m,n})/\epsilon$ . The partition function is then,

$$Z = C \int \mathcal{D}\varphi \exp \left\{ - \int_0^\beta d\tau \int_V d^3x \left[ \frac{1}{2} (\dot{\varphi}^2 + \nabla\varphi^2) + \mathcal{V}(\varphi) \right] \right\}. \quad (55)$$

Here,  $C$  is the divergent constant that results from the  $\pi$ -integral. In the last line, we exploited the fact that the variable of integration  $\pi_{n,m}$  may be shifted by a constant without changing the integral, and used the Gaussian integral

$$\int_{-\infty}^{\infty} dx e^{-ax^2/2} = \sqrt{\frac{2\pi}{a}}.$$

The partition function resulting from this procedure may also be obtained the ground state path integral

$$Z_g = \int \mathcal{D}\varphi \mathcal{D}\pi \exp \left\{ i \int_{\Omega'} d^4x (\pi\dot{\varphi} - \mathcal{H}[\varphi, \pi]) \right\} = C' \int \mathcal{D}\varphi(x) \exp \left\{ i \int_{\Omega'} d^4x \mathcal{L}[\varphi, \partial_\mu \varphi] \right\}.$$

The domain of the functional integral  $\int \mathcal{D}\varphi$  is restricted from *all* (smooth enough) field configurations  $\varphi(t, \vec{x})$ , to only those that obey  $\varphi(\beta, \vec{x}) = e^{i\theta} \varphi(0, \vec{x})$ , which is denoted  $S$ . The action integral is modified by performing a Wick-rotation of the time coordinate  $t$ . This involves changing the domain of  $t$  from the real line to the imaginary line by closing the contour at infinity, and making the change of variable  $it \rightarrow \tau$ . The new variable is then restricted to the interval  $\tau \in [0, \beta]$ . This procedure motivates the introduction of the Euclidean Lagrange density,  $\mathcal{L}_E(\tau, \vec{x}) = -\mathcal{L}(-i\tau, \vec{x})$ , as well as the name “imaginary-time formalism”. The result is the same partition function as before,

$$Z = C \int_S \mathcal{D}\varphi \int \mathcal{D}\pi \exp \left\{ \int_0^\beta d\tau \int_V d^3x [i\dot{\varphi}\pi - \mathcal{H}(\varphi, \pi)] \right\} = C' \int_S \mathcal{D}\varphi \exp \left\{ - \int_0^\beta d\tau \int_V d^3x \mathcal{L}_E(\varphi, \pi) \right\}.$$

## Free scalar field

This section uses notation as described in Appendix A. The procedure for obtaining the thermal properties of an interacting scalar field is similar to that used in scattering theory. One starts with a free theory, which can be solved exactly. Then an interaction term is added, which is accounted for perturbatively by using Feynman diagrams. The Euclidean Lagrangian for a free scalar gas is, after integrating by parts,

$$\mathcal{L}_E = \frac{1}{2} \varphi(X) (-\partial_E^2 + m^2) \varphi(X)$$

Here,  $\Omega$  is the domain  $[0, \beta] \times V$ , and  $X = (\tau, \vec{x})$  is the Euclidean coordinate which is a result of the Wick-rotation. Furthermore, we have introduced the Euclidean Laplace operator,  $\partial_E^2 = \partial_\tau^2 + \nabla^2$ . Following the procedure as described in section 3 to obtain the thermal partition function yields

$$Z = C \int_S \mathcal{D}\varphi(X) \exp \left\{ - \int_\Omega dX \frac{1}{2} \varphi(X) (-\partial_E^2 + m^2) \varphi(X) \right\}.$$

Here,  $K = (\omega_n, \vec{k})$  is the Euclidean Fourier-space coordinate. We then insert the Fourier expansion of  $\varphi$ , and change the functional integration variable to the Fourier components. The integration measures are related by

$$\mathcal{D}\varphi(X) = \det \left( \frac{\delta\varphi(X)}{\delta\tilde{\varphi}(K)} \right) \mathcal{D}\tilde{\varphi}(K).$$

The determinant factor which appears may be absorbed into the constant  $C$ , as the integration variables are related by a linear transform. The action becomes

$$\begin{aligned} S &= - \int_{\Omega} dX \mathcal{L}_e = -\frac{1}{2} V \beta \int_{\Omega} dX \int_{\tilde{\Omega}} dK \int_{\tilde{\Omega}} dK' \tilde{\varphi}(K') \left( \omega_n^2 + \vec{k}^2 + m^2 \right) \tilde{\varphi}(K) e^{iX \cdot (K - K')} \\ &= -\frac{1}{2} V \beta^2 \int_{\tilde{\Omega}} dK \tilde{\varphi}(K)^* \left( \omega_n^2 + \omega_k^2 \right) \tilde{\varphi}(K), \end{aligned}$$

where  $\omega_k^2 = \vec{k}^2 + m^2$ .  $\tilde{\Omega}$  is the reciprocal space corresponding to  $\Omega$ , as described in Appendix A. We used the fact that  $\varphi$  is real, which implies that  $\tilde{\varphi}(-K) = \tilde{\varphi}(K)^*$ , as well as the identity Eq. (118). This gives the partition function

$$Z = C \int_{\tilde{S}} \mathcal{D}\tilde{\varphi}(K) \exp \left\{ -\frac{1}{2} V \int_{\tilde{\Omega}} dK \tilde{\varphi}(K)^* [\beta^2 (\omega_n^2 + \omega_k^2)] \tilde{\varphi}(K) \right\},$$

Going back to before the continuum limit, this integral can be written as a product of Gaussian integrals, and may therefore be evaluated

$$Z = C \prod_{n=-\infty}^{\infty} \prod_{k \in \tilde{V}} \left( \int d\tilde{\varphi}_{n,\vec{k}} \exp \left\{ -\frac{1}{2} \tilde{\varphi}_{n,\vec{k}}^* [\beta^2 (\omega_n^2 + \omega_k^2)] \tilde{\varphi}_{n,\vec{k}} \right\} \right) = C \prod_{n=-\infty}^{\infty} \prod_{k \in \tilde{V}} \sqrt{\frac{2\pi}{\beta^2 (\omega_n^2 + \omega_k^2)}}.$$

The partition function is related to Helmholtz free energy  $F$  through

$$\frac{F}{TV} = -\frac{\ln(Z)}{V} = \frac{1}{2} \int_{\tilde{\Omega}} dK \frac{1}{2} \ln[\beta^2 (\omega_n^2 + \omega_k^2)] + \frac{F_0}{TV},$$

where  $F_0$  is a constant.

A faster and more formal way to get to this result is to compare the partition function to the multidimensional version of the Gaussian integral [8, 6]. The partition function is on the form

$$I_n = \int_{\mathbb{R}^n} d^n x \exp \left\{ -\frac{1}{2} \langle x, Dx \rangle \right\},$$

where  $D$  is a linear operator, and  $\langle \cdot, \cdot \rangle$  an inner product on the corresponding vector space. By diagonalizing  $D$ , we get the result

$$I_n = \sqrt{\frac{(2\pi)^n}{\det(D)}}.$$

We may then use the identity

$$\det(D) = \prod_i \lambda_i = \exp\{\text{Tr}[\ln(D)]\}, \quad (56)$$

where  $\lambda_i$  are the eigenvalues of  $D$ . The trace in this context is defined by the vector space  $D$  acts on. For given an orthonormal basis  $x_n$ , such that  $\langle x_n, x'_n \rangle = \delta_{nn'}$  the trace can be evaluated as  $\text{Tr}\{D\} = \sum_n \langle x_n, Dx_n \rangle$ . Identifying

$$\langle x, Dx \rangle = \int_{\Omega} dX \varphi(X) (-\partial_E^2 + m^2) \varphi(X),$$

we get the formal result

$$Z = C \det(-\partial_E^2 + m^2)^{-1/2},$$

and

$$\beta F = \frac{1}{2} \text{Tr}\{\ln(-\partial_E^2 + m^2)\} + C.$$

The logarithm may then be evaluated by using the eigenvalues of the linear operator. This is found by diagonalizing the operator,

$$\langle x, Dx \rangle = \int_{\Omega} dX \varphi(X) (-\partial_E^2 + m^2) \varphi(X) = V \int_{\tilde{\Omega}} dK \tilde{\varphi}(K)^* [\beta^2 (\omega_k^2 + \omega_n^2)] \tilde{\varphi}(K),$$

leaving us with the same result

$$\beta F = \frac{1}{2} \text{Tr} \{ \ln(-\partial_E^2 + m^2) \} = \frac{1}{2} V \int_{\tilde{\Omega}} dK \ln[\beta^2(\omega_n^2 + \omega_k^2)].$$

How to work this sum is shown in section 3 This gives the free energy density, after dropping the additive constant,

$$\beta \mathcal{F} = \frac{F}{VT} = \frac{1}{2} \int_{\tilde{V}} \frac{d^3 k}{(2\pi)^3} [\beta \omega_k + 2 \ln(1 - e^{-\beta \omega_k})] + \frac{F_0}{VT} \quad (57)$$

## Thermal sum

When evaluating thermal integral, we will often encounter sums of the form

$$j(\omega, \mu) = \frac{1}{2\beta} \sum_{\omega_n} \ln\{\beta^2[(\omega_n + i\mu) + \omega^2]\} + g(\beta), \quad (58)$$

where the sum is over either the bosonic Matsubara frequencies  $\omega_n = 2n\pi/\beta$ ,  $n \in \mathbb{Z}$ , or the fermionic ones,  $\omega_n = (2n+1)\pi/\beta$ ,  $n \in \mathbb{Z}$ .  $\mu \in \mathbb{R}$  is the chemical potential.  $g$  may be a function of  $\beta$ , but we assume it is independent of  $\omega$ . Thus, the factor  $\beta^2$  could strictly be dropped, but it is kept to make the argument within the logarithm dimensionless. We define the function

$$i(\omega, \mu) = \frac{1}{\omega} \frac{d}{d\omega} j(\omega, \mu) = \frac{1}{\beta} \sum_{\omega_n} \frac{1}{(\omega_n + i\mu)^2 + \omega^2}. \quad (59)$$

Assume  $f(z)$  is an analytic function, except perhaps on a set of isolated poles  $\{z_i\}$  located outside the real line. By exploiting the properties of  $in_B(iz)$ , as described in Appendix A, we can rewrite the sum over Matsubara frequencies as a contour integral

$$\frac{1}{\beta} \sum_{\omega_n} f(\omega_n) = \oint_{\gamma} \frac{dz}{2\pi i} f(z) (in_B(iz)),$$

where  $\gamma$  is a contour that goes from  $-\infty - i\epsilon$  to  $+\infty - i\epsilon$ , crosses the real line at  $\infty$ , goes from  $+\infty - i\epsilon$  to  $-\infty + i\epsilon$  before closing the curve. (FIGURE) This result exploits Cauchy's integral formula, by letting the poles of  $in_B(iz)$  at the Matsubara frequencies "pick out" the necessary residues. The integral over  $\gamma$  is equivalent to two integrals along  $\mathbb{R} \pm i\epsilon$ ,

$$\begin{aligned} \frac{1}{\beta} \sum_{\omega_n} f(\omega_n) &= \left( \int_{\infty+i\epsilon}^{-\infty+i\epsilon} \frac{dz}{2\pi} + \int_{-\infty-i\epsilon}^{\infty-i\epsilon} \frac{dz}{2\pi} \right) f(z) n_B(iz), \\ &= \int_{-\infty-i\epsilon}^{\infty-i\epsilon} \frac{dz}{2\pi} [f(z) + (f(z) + f(-z)) n_B(iz)]. \end{aligned}$$

In the last line, we have changed variables  $z \rightarrow -z$  in the first integral, and exploited the property  $n_B(-iz) = -1 - n_B(iz)$ . As  $n_B(iz)$  is analytic on the real line, the result is the sum of residues of  $f$  in the lower half plane. The function

$$f(z) = \frac{1}{(z + i\mu)^2 + \omega^2} = \frac{i}{2\omega} \left( \frac{1}{z + i(\mu + \omega)} - \frac{1}{z + i(\mu - \omega)} \right) \quad (60)$$

obeys the assumed properties, as it has poles at  $z = -i(\mu \pm \omega)$ , with residue  $1/2\omega$ , so the function defined in Eq. (59) may be written <sup>4</sup>

$$i(\omega, \mu) = \frac{1}{2\omega} [1 + n_B(\omega + \mu) + n_B(\omega - \mu)]. \quad (61)$$

Using the anti-derivative of the Bose distribution, we get the final form of Eq. (58)

$$j(\omega, \mu) = \int d\omega' \omega' i(\omega', \mu) = \frac{1}{2} \omega + \frac{1}{2\beta} \left[ \ln(1 - e^{-\beta(\omega + \mu)}) + \ln(1 - e^{-\beta(\omega - \mu)}) \right] + g'(\beta). \quad (62)$$

The extra  $\omega$ -independent term  $g'(\beta)$  is an integration constant. (Noe om antipartikkel)

---

<sup>4</sup>Assuming  $\omega > \mu$ .




Figure 1: The integral contour  $\gamma$ , and the result of deforming it into to contours close to the real line.

## Regulating the free energy

The free energy integral we obtained from the energy for the free scalar Eq. (57), has two parts, Noticing that the integral is spherically symmetric, we may write

$$J_0 = \frac{1}{2} \int_{\Omega} \frac{d^3 k}{(2\pi)^3} \sqrt{k^2 + m^2}, \quad J_T = \frac{1}{\beta} \int_{\Omega} \frac{d^3 k}{(2\pi)^3} \ln \left[ 1 - \exp\left(-\beta \sqrt{k^2 + m^2}\right) \right], \quad (63)$$

The temperature-independent part,  $J_0$ , is clearly divergent, and we must therefore impose a regulator, and then add counter-terms. This differs from the zero-temperature formalism, where there were no need to renormalize the free theory. The second part of the integral, is convergent. To see this, we first exploit that it is spherically symmetric to write

$$J_T = \frac{T^4}{2\pi^2} \int_{\mathbb{R}} dx x^2 \ln \left( 1 - e^{-\sqrt{x^2 + (\beta m)^2}} \right). \quad (64)$$

Using the series expansion  $\ln(1 + \epsilon) \sim \epsilon + \mathcal{O}(\epsilon^2)$ , we can find the leading part of the integrand for large  $k$ 's,

$$x^2 \ln \left( 1 - e^{-\sqrt{x^2 + (\beta m)^2}} \right) \sim -x^2 e^{-x}, \quad (65)$$

which is suppressed exponentially, making the integral convergent. In the limit of  $T \rightarrow \infty$ , we get

$$J_{\infty} \sim \frac{T^4}{2\pi^2} \int_{\mathbb{R}} dx x^2 \ln(1 - e^{-x}) = -T^4 \frac{\pi^2}{90} = -\frac{3}{2} \sigma T^4, \quad (66)$$

where  $\sigma$  is the Stefan-Boltzmann constant.

We use dimensional regularization on the temperature independent term. To that end, we define

$$\Phi(m, d, A) = \int_{\Omega} \frac{d^d k}{(2\pi)^d} (k^2 + m^2)^{-A}, \quad (67)$$

so that  $J_0 = \Phi(m, 3, 1/2)/2$ . Dimensional regularization takes uses the formula for integration of spherically symmetric function in  $d$ -dimensions,

$$\int_{\mathbb{R}^d} d^d x f(r) = \frac{2\pi^{d/2}}{\Gamma(d/2)} \int_{\mathbb{R}} dr r^{d-1} f(r), \quad (68)$$

where  $r = \sqrt{x_i x_i}$  is the radial distance, and  $\Gamma$  is the gamma-function. The factor in the front of the integral comes from the solid-angle. By extending this formula from integer-valued  $d$  to real numbers, the function we defined becomes

$$\Phi = \frac{2\pi^{d/2}}{\Gamma(d/2)} \int_{\mathbb{R}} dk \frac{k^{d-1}}{(k^2 + m^2)^A} = \frac{m^{d-2A}}{(4\pi)^{d/2} \Gamma(d/2)} 2 \int_{\mathbb{R}} dz \frac{z^{d-1}}{(1+z)^A}, \quad (69)$$

where we have made the change of variables  $mz = k$ . We make one more change of variable to the integral,

$$I = 2 \int_{\mathbb{R}} dz \frac{z^{d-1}}{(1+z)^A} \quad (70)$$

Let

$$z^2 = \frac{1}{s} - 1 \implies 2zdz = -\frac{ds}{s^2} \quad (71)$$

Thus,

$$I = \int_0^a ds s^{A-d/2-1} (1-z)^{d/2-1}. \quad (72)$$

This is the beta-function, which can be written in terms of gamma-funcitons [6]

$$I = B\left(A - \frac{d}{2}, \frac{d}{2}\right) = \frac{\Gamma\left(A - \frac{d}{2}\right) \Gamma\left(\frac{d}{2}\right)}{\Gamma(A)}. \quad (73)$$

Putting it all together, this gives

$$\Phi = \frac{(m^2)^{d/2-A} \Gamma\left(A - \frac{d}{2}\right)}{(4\pi)^{d/2} \Gamma(A)}. \quad (74)$$

Inserting  $d = 3 - 2\epsilon$  and  $A = -1/2$ , we get

$$\frac{(m^2)^{3/2-\epsilon-A} \Gamma(-2+\epsilon)}{(4\pi)^{3/2-\epsilon} \Gamma(-1/2)} = \left(\frac{m^2}{4\pi}\right)^{3/2} \frac{m}{-2\pi^{1/2}} \mu^{-2\epsilon} \left(\frac{m^2}{4\pi\mu^2}\right)^{-\epsilon} \frac{\Gamma(\epsilon)}{(\epsilon-2)(\epsilon-1)}, \quad (75)$$

where we have used the defining property  $\Gamma(z+1) = z\Gamma(z)$ , and inserted a parameter  $\mu$  with the dimensions of  $m$ . Expanding around  $\epsilon = 0$  gives

$$\left(\frac{m^2}{4\pi\mu^2}\right)^{-\epsilon} \sim 1 + \epsilon \ln\left(4\pi\frac{\mu^2}{m^2}\right), \quad (76)$$

$$\Gamma(\epsilon) \sim \frac{1}{\epsilon} - \gamma, \quad (77)$$

$$\frac{1}{(\epsilon-2)(\epsilon-1)} \sim \frac{1}{2} \left(1 + \frac{3}{2}\epsilon\right). \quad (78)$$

Inserting this back into the function gives

$$\Phi = -\frac{m^4}{2^5\pi^2} \mu^{-2\epsilon} \left[1 + \epsilon \ln\left(4\pi\frac{\mu^2}{m^2}\right)\right] \left(\frac{1}{\epsilon} - \gamma\right) \left(1 + \frac{3}{2}\epsilon\right) \sim -\frac{m^4}{2^5\pi^2} \mu^{-2\epsilon} \left[\frac{1}{\epsilon} - \gamma + \frac{3}{2} + \ln\left(4\pi\frac{\mu^2}{m^2}\right)\right]. \quad (79)$$

## Interacting scalar

We now study a scalar field with a  $\lambda\varphi^4$  interaction term. We write the Lagrangian in the form

$$\mathcal{L} = \mathcal{L}^{(0)} + \mathcal{L}^{(I)}, \quad \mathcal{L}^{(0)} = \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi + m^2 \varphi^2, \quad \mathcal{L}^{(I)} = -\frac{\lambda}{4!} \varphi^4$$

$\mathcal{L}^{(I)}$  is called the interaction term, and makes it impossible to exactly solve for the partition function. Instead, we turn to perturbation theory. The grand canonical partition function in this theory

$$Z = \text{Tr}\left\{e^{-\beta\hat{H}}\right\} = \int_S \mathcal{D}\varphi \exp\left\{-\int_\Omega dX \left(\mathcal{L}_E^{(0)} + \mathcal{L}_E^{(I)}\right)\right\} = \int_S \mathcal{D}\varphi e^{S_0} e^{S_I}. \quad (80)$$

Here,  $S_0$  and  $S_I$  denote the Euclidean action due to the free and interacting Lagrangian, respectively. The domain of integration  $S$  is again periodic field configurations  $\varphi(\beta, \vec{x}) = \varphi(0, \vec{x})$ . We may write the free energy as

$$-\beta F = \ln \left[ \int_S \mathcal{D}\varphi e^{S_0} \sum_n \frac{1}{n!} S_I^n \right] = \ln[Z_0] + \ln[Z_I],$$

where  $Z_0$  is the partition function from the free theory. The correction to the partition function is thus given by

$$Z_1 = \sum_{n=0}^{\infty} \frac{1}{n!} \langle S_I^n \rangle_0, \quad (81)$$

where

$$\langle A \rangle_0 = \frac{\int_S \mathcal{D}\varphi A e^{S_0}}{\int_S \mathcal{D}\varphi e^{S_0}}. \quad (82)$$

To evaluate expectation values of the form  $\langle \varphi(X_1) \dots \rangle_0$ , we introduce the the partition function with a source term

$$Z[J] = \int_S \mathcal{D}\varphi \exp \left\{ -\frac{1}{2} \int_{\Omega} dX \varphi (-\partial_E^2 + m^2) \varphi + \int_{\Omega} dX J \varphi \right\}. \quad (83)$$

Using the thermal Greens function  $D(X, Y)$ , as defined in Appendix A, we may complete the square to write

$$Z[J] = Z[0] \exp \left\{ \frac{1}{2} \int_{\Omega} dX dY J(X) D_0(X, Y) J(Y) \right\} = Z[0] \exp(W[J]) \quad (84)$$

We can now write

$$\langle \varphi(X) \varphi(Y) \rangle_0 = \frac{1}{Z[0]} \frac{\delta}{\delta J(X)} \frac{\delta}{\delta J(Y)} Z[J] \Big|_{J=0} = D(X, Y), \quad (85)$$

where  $D(X, Y)$  is the thermal propagator, as defined in Appendix A This generalizes to higher order expectation values,

$$\langle \varphi(X_1) \dots \varphi(X_n) \rangle_0 = \left( \prod_{i=1}^n \frac{\delta}{\delta J(X_i)} \right) Z[J] \Big|_{J=0}, \quad (86)$$

The exponential form of  $Z[J]$  leads straight forwardly to Wick's theorem, which states that an expectation value of  $2n$  fields is a sum of *all possible, distinct* combination of  $n$  propagators. To write this in a formal way, we define the functions  $a$  and  $b$ , which define a way to pair up  $2m$  elements. The domain of the functions are the integers between 1 and  $m$ , the image a subset of the integers between 1 and  $2m$  of size  $m$ . A valid pairing is a set  $\{(a(1), b(1)), \dots (a(m), b(m))\}$ , where all elements  $a(i)$  and  $b(j)$  are different, such all integers up to and including  $2m$  are featured. A pair is not directed, so  $(a(i), b(i))$  is the same pair as  $(b(i), a(i))$ . Wick theorem states that,

$$\left\langle \prod_{i=1}^{2m} \varphi(X_i) \right\rangle_0 = \sum_{\{(a,b)\}} \langle \varphi(X_{a(i)}) \varphi(X_{b(i)}) \rangle. \quad (87)$$

where the sum is over all tuples  $(a, b)$  that define a valid and unique pairing. Using Wick's theorem, the expectation values we are evaluating can be written

$$\begin{aligned} \langle S_I^m \rangle &= \left( -\frac{\lambda}{4!} \right)^m \int_{\Omega} dX_1 \dots dX_m \langle \varphi(X_1)^4 \dots \varphi(X_m)^4 \rangle \\ &= \left( -\frac{\lambda}{4!} \right)^m \int_{\Omega} dX_1 \dots dX_m \sum_{\{(a,b)\}} \langle \varphi(X_{a(1)}) \varphi(X_{b(1)}) \rangle \dots \langle \varphi(X_{a(2m)}) \varphi(X_{b(2m)}) \rangle \end{aligned}$$




where  $X_i$  for  $i > m$  is defined as  $X_j$ , where  $j = i \bmod m$ . Inserting the fourier expansions of the field gives

$$\begin{aligned}
\langle S_I^m \rangle &= \left( -\frac{\lambda}{4!} \right)^m \int_{\Omega} dX_1 \dots dX_m (V\beta)^2 \int_{\tilde{\Omega}} dK_1 \dots dK_{2m} \sum_{\{a,b\}} \\
&\quad \langle \varphi(K_{a(1)}) \varphi(K_{b(1)}) \rangle \dots \langle \varphi(K_{a(2m)}) \varphi(K_{b(2m)}) \rangle \exp \left( i \sum_{i=1}^m X_i \cdot K_i \right) \\
&= \left( -\frac{\lambda}{4!} \right)^m \frac{(V\beta)^{2m} \beta^m}{(V\beta^2)^{2m}} \int_{\tilde{\Omega}} dK_1 \dots dK_{2m} \sum_{\{a,b\}} \\
&\quad \tilde{D}(K_{a(1)}) \delta(K_{a(1)} + K_{b(1)}) \dots \tilde{D}(K_{a(2m)}) \delta(K_{a(2m)} + K_{b(2m)}) \prod_{i=1}^m \delta \left( X_i \cdot \sum_{j=0}^3 K_{i+jm} \right) \\
&= \left( -\frac{\lambda\beta}{4!} \right)^m \prod_{i=1}^{2m} \int_{\tilde{\Omega}} \left( dK_i \frac{1}{\beta} \tilde{D}(K_i) \right) \prod_{i=1}^m \delta \left( X_i \cdot \sum_{j=0}^3 K_{i+jm} \right) \sum_{\{a,b\}} \prod_{n=1}^{2m} \delta(K_{a(n)} + K_{b(n)})
\end{aligned}$$


Here we have used that  $V\beta^2 \tilde{D}_0(K, P) = \tilde{D}_0(K) \delta(P + K)$ , and  $\tilde{D}(K)$  is the thermal propagator for the free field, as defined in Appendix A. In this case, the thermal propagator of the free field is

$$D_0(K) = D_0(\omega_n, \vec{k}) = \frac{1}{\omega_k^2 + \omega_n^2}. \quad (88)$$

This expectation value can be represented graphically using Feynman diagrams. The thermal  $\lambda\varphi^2$ -theory gets the prescription




$$= -\lambda\beta\delta\left(\sum_i K_i\right), \quad (89)$$



$$= \frac{1}{\beta} \int_{\tilde{\Omega}} dK D_0(K). \quad (90)$$

The factor  $1/4!$  is removed as a general Feynman diagram represent all diagrams with the same form, but different pairing of the momenta. Some diagrams are more symmetric, such that an exchange of momenta still gives *the same pairing*. This is dealt with by dividing with a symmetry factor  $s$ , which is described in detail in [6].

Calculating  $\langle S_I^n \rangle_0$  boils down to the sum of all possible Feynman diagrams with  $m$  vertices. The first example is



$$\langle S_I \rangle = \text{[Three loop diagrams]} = \frac{3}{4!} \times \text{[Two-loop diagram]}. \quad (91)$$

For higher order, one gets both connected and disconnected diagrams. Let  $\langle S_I^n \rangle_{0,c}$  be only the connected diagrams, i.e. those diagrams in which it is possible to move between all vertices along a series of edges.

A general diagram contain  $n_i$  copies of a connected diagram with the value  $V_i$ . The value of the total diagram is then the product of the value of all its disconnected pieces, but with the caveat that each diagram has a symmetry factor of  $n_i!$ . The sum of all diagrams is thus

$$Z_I = \sum_n \frac{1}{n!} \langle S_I^n \rangle = \sum_{\text{all sets } \{n_i\}} \prod_i \frac{1}{n_i!} V_i^{n_i} = \prod_i \sum_{n_i} \frac{1}{n_i!} V_i^{n_i} = \exp\left(\sum_i V_i\right). \quad (92)$$

Thus, the correction to the free energy is given by the sum of all connected diagrams,

$$-\beta F = \ln(Z_0) + \sum_n \langle S_I^n \rangle_{0,c}. \quad (93)$$

## Rester

Expectation values of field configurations with different momenta vanish,

$$K' \neq K \implies \langle \varphi(K) \varphi(K') \rangle_0 = 0. \quad (94)$$

This is due to the symmetry of the functional integrand. This can be seen by going back to the discrete version of the integral. The integral can then be factored into integrals over each degree of freedom,  $\varphi_{n,m}$ . The integrall over the degrees of freedom corresponding to  $\varphi(K)$  and  $\varphi(K')$  are of the form

$$\int d\varphi_{n,m} \varphi_{n,m} \exp(-S_{n,m}^{(0)}) = 0, \quad (95)$$

as the integral is over the entier real line, and the integrand is anti-symmetric in  $\varphi_{n,m}$ . The generalization of this rule is straight forward. Assuming  $f[\tilde{\varphi}]$  *does not* depend on  $\tilde{\varphi}(K)$ , but possibly on  $\tilde{\varphi}(K')$ ,  $K \neq K'$ , then

$$\left\langle f[\tilde{\varphi}] (\tilde{\varphi}(K))^{(2n+1)} \right\rangle_0 = 0, \quad n \in \mathbb{Z}. \quad (96)$$

Furthermore, as the field at different values of  $K$  are independent, we get the familiar formula for expectation values of independent variables, that is

$$\langle f[\varphi(K)] g[\varphi(K')] \rangle_0 = \langle f[\varphi(K)] \rangle_0 \langle g[\varphi(K')] \rangle_0, \quad K \neq K'. \quad (97)$$

This leads to a version of Wick's theorem for thermal systems, as it lays constrains on integrals of the form

$$\int_{\tilde{\Omega}} dK_1 \dots dK_n \langle \tilde{\varphi}(K_1) \dots \tilde{\varphi}(K_n) \rangle_0 \delta\left(\sum_i K_i\right) = \quad (98)$$

## Fermions

The phase factor  $e^{i\theta}$  that was introduced in (REF IMAG FORM.) can be decided by studying the properties of the thermal greens function, which are defined as

$$G(X_1, X_2) = G(\vec{x}, \vec{y}, \tau_1, \tau_2) = \left\langle e^{-\beta \hat{H}} T_\tau \{ \varphi(X_1) \varphi(X_2) \} \right\rangle.$$

$T_\tau \{ \dots \}$  is the temperature ordering operator, analogous to the time ordering operator from scattering theory. It is defined as

$$T_\tau \{ \varphi(\tau_1) \varphi(\tau_2) \} = \theta(\tau_1 - \tau_2) \varphi(\tau_1) \varphi(\tau_2) + \nu \theta(\tau_2 - \tau_1) \varphi(\tau_2) \varphi(\tau_1),$$

where  $\nu = \pm 1$  for respectively bosons and fermions, and  $\theta(\tau)$  is the Heaviside step function. In the same way that  $i\hat{H}$  generates the time-dependence of a quantum field operator through  $\hat{\varphi}(x) = \hat{\varphi}(t, \vec{x}) = e^{it\hat{H}} \hat{\varphi}(\vec{x}) e^{-it\hat{H}}$ , the imaginary-time formalism implies the relation

$$\hat{\varphi}(X) = \hat{\varphi}(\tau, \vec{x}) = e^{\tau \hat{H}} \hat{\varphi}(\vec{x}) e^{-\tau \hat{H}}. \quad (99)$$

Using  $\mathbb{1} = e^{\tau\hat{H}}e^{-\tau\hat{H}}$  and the cyclic property of the trace, we show that, assuming  $\beta > \tau > 0$ ,

$$\begin{aligned}
G(\vec{x}, \vec{y}, \tau, 0) &= \left\langle e^{-\beta\hat{H}} \text{Tr}_\tau \{ \varphi(\tau, \vec{x}) \varphi(0, \vec{y}) \} \right\rangle \\
&= \frac{1}{Z} \text{Tr} \left\{ e^{-\beta\hat{H}} \varphi(\tau, \vec{x}) \varphi(0, \vec{y}) \right\} \\
&= \frac{1}{Z} \text{Tr} \left\{ \varphi(0, \vec{y}) e^{-\beta\hat{H}} \varphi(\tau, \vec{x}) \right\} \\
&= \frac{1}{Z} \text{Tr} \left\{ e^{-\beta\hat{H}} e^{\beta\hat{H}} \varphi(0, \vec{y}) e^{-\beta\hat{H}} \varphi(\tau, \vec{x}) \right\} \\
&= \frac{1}{Z} \text{Tr} \left\{ e^{-\beta\hat{H}} \varphi(\vec{y}, \beta) \varphi(\tau, \vec{x}) \right\} \\
&= \nu \left\langle e^{-\beta\hat{H}} \text{Tr}_\tau \{ \varphi(\tau, \vec{x}) \varphi(\vec{y}, \beta) \} \right\rangle.
\end{aligned}$$

This implies that  $\varphi(0, x) = \nu\varphi(\beta, \varphi)$ , which show that bosons are periodic in time, as stated earlier, while fermions are anti-periodic.

The Lagrangian density of a free fermion is

$$\mathcal{L} = \bar{\psi} (i\partial\!\!\!/ - m) \psi. \quad (100)$$

This Lagrangian is invariant under the transformation  $\psi \rightarrow e^{-i\alpha}\psi$ , which by Nöther's theorem results in a conserved current

$$j^\mu = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \psi)} \delta\psi = \bar{\psi} \gamma^\mu \psi. \quad (101)$$

The corresponding charge is

$$Q = \int_V d^3x j^0 = \int_V d^3x \psi^\dagger \psi. \quad (102)$$

We can now use our earlier result for the thermal partition function, Eq. (54), only with the substitution  $\mathcal{H} \rightarrow \mathcal{H} - \mu\psi^\dagger\psi$ , and integrate over anti-periodic  $\psi$ 's:

$$Z = \text{Tr} \left\{ e^{-\beta(\hat{H} - \mu\hat{Q})} \right\} = \prod_{ab} \int \mathcal{D}\psi_a \mathcal{D}\pi_b \exp \left\{ \int_\Omega dX \left( i\dot{\psi}\pi - \mathcal{H}(\psi, \pi) + \mu\psi^\dagger\psi \right) \right\},$$

where  $a, b$  are the spinor indices. The canonical momentum corresponding to  $\psi$  is

$$\pi = \frac{\partial \mathcal{L}}{\partial(\partial_0 \psi)} = i\psi^\dagger, \quad (103)$$

and the Hamiltonian density is

$$\mathcal{H} = \pi\dot{\psi} - \mathcal{L} = \bar{\psi}(-\gamma^i\partial_i + m)\psi \quad (104)$$

which gives

$$i\dot{\psi}\pi - \mathcal{H}(\psi, \pi) + \mu\psi^\dagger\psi = \bar{\psi}[-\gamma^0(\partial_\tau - \mu) + i\gamma^i\partial_i - m]\psi, \quad (105)$$

By using the Grassman-version of the Gaussian integral formula, the partition function can be written

$$Z = \prod_{ab} \int \mathcal{D}\psi_a \mathcal{D}i\psi_b^\dagger \exp \left\{ \int_\omega dX \bar{\psi} [-\gamma^0(\partial_\tau - \mu) + i\gamma^i\partial_i - m] \psi \right\} = \prod_{ab} \int \mathcal{D}\psi_a \mathcal{D}i\psi_b^\dagger e^{\langle i\psi^\dagger, D\psi \rangle} = C \det(D),$$

where

$$D = -i\gamma^0[(\mu - \partial_\tau) + i\gamma^i\partial_i - m]. \quad (106)$$

This results in

$$-\beta F = \ln(Z) = \text{Tr}[\ln(D)] + C. \quad (107)$$

Assuming  $\psi_a(X)$  is a orthonormal set, we can calculate the trace

$$\begin{aligned}
\text{Tr}[\ln(D)] &= \int_{\Omega} dX (i\psi_a(X)^\dagger) \ln\{-i\gamma^0[\gamma^0(\mu - \partial_\tau) + i\gamma^i\partial_i - m]\}_{ab}\psi_b(X) \\
&= V \int_{\tilde{\Omega}} dK \ln\{-i\beta\gamma^0[\gamma^0(\mu - i\omega_n) - \gamma^i p_i - m]\}_{aa} \\
&= V \int_{\tilde{\Omega}} dK \ln\{i\beta\gamma^0[i\gamma^0(\omega_n + i\mu) + i(-i\gamma^i)p_i + m]\}_{aa} \\
&= V \int_{\tilde{\Omega}} dK \ln\{i\beta\gamma^0[i\tilde{\gamma}_a p_{n;a} + m]\}_{aa}.
\end{aligned}$$

In the last line, we have introduced the notation  $p_{n;a} = (\omega_n + i\mu, p_i)$  and use the Euclidean gamma matrices, as defined in Appendix A. (JEG ER IKKE SIKKER PÅ DETTE) Furthermore, we use the fact that

$$\ln[i\tilde{\gamma}_a p_a - m] = \ln[i\tilde{\gamma}_a p_a - m] + \ln(\tilde{\gamma}_5 \tilde{\gamma}_5) = \ln[\tilde{\gamma}_5(i\tilde{\gamma}_a p_a - m)\tilde{\gamma}_5] = \ln[-i\tilde{\gamma}_a p_a - m],$$

which allows us to calculate

$$\begin{aligned}
&\ln\{i\beta\gamma^0[i\tilde{\gamma}^0(\omega_n + i\mu) + i(-i\tilde{\gamma}^i)p_i + m]\} \\
&= \ln[(i\tilde{\gamma}_a p_a + m)] + \frac{1}{4} \ln[(i\beta\gamma^0)^4] \\
&= \ln[\beta] + \frac{1}{2} \{\ln[i\tilde{\gamma}^a p^a + m] + \ln[-i\tilde{\gamma}^a p^a + m]\} \\
&= \frac{1}{2} \{\ln[\beta^2(i\tilde{\gamma}^a p^a + m)(-i\tilde{\gamma}^a p^a + m)]\} \\
&= \frac{1}{2} \mathbb{1} \ln\{\beta^2[(\omega_n + i\mu)^2 + \omega^2]\}
\end{aligned}$$

which gives

$$\mathcal{F} = -\frac{2}{\beta} \int_{\tilde{\Omega}} dX \ln\{\beta^2[(\omega_n + i\mu)^2 + \omega^2]\}. \quad (108)$$

Using the fermionic version of the thermal sum, this gives the answer

$$\mathcal{F} = -\frac{2}{\beta} \int \frac{d^3 p}{(2\pi)^3} \left[ \beta\omega + \ln\left(1 + e^{-\beta(\omega+\mu)}\right) + \ln\left(1 + e^{-\beta(\omega-\mu)}\right) \right]. \quad (109)$$

## References

- [1] Steven Weinberg. Phenomenological lagrangians. *Physica A: Statistical Mechanics and its Applications*, 96(1):327–340, 1979.
- [2] S. Scherer. Introduction to chiral perturbation theory. *Advances in Nuclear Physics*, 27:277–538, 2002.
- [3] Martin Aria Mojahed. Two-flavor chiral perturbation theory at finite isospin beyond leading order. Master’s thesis, NTNU, 2020.
- [4] J.S.R. Chisholm. Change of variables in quantum field theories. *Nuclear Physics*, 26(3):469–479, 1961.
- [5] S. Kamefuchi, L. O’Raifeartaigh, and Abdus Salam. Change of variables and equivalence theorems in quantum field theories. *Nuclear Physics*, 28(4):529–549, 1961.
- [6] Michael E. Peskin and Daniel V. Schroeder. *An Introduction to quantum field theory*. Addison-Wesley, Reading, USA, 1995.
- [7] Sean M. Carroll. *Spacetime and Geometry*. Cambridge University Press, 7 2019.
- [8] J. I. Kapusta and Charles Gale. *Finite-temperature field theory: Principles and applications*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2011.

## Appendix A Conventions and notation

Throughout this text, natural units are employed, in which

$$\hbar = c = k_B = 1, \quad (110)$$

where  $\hbar$  is the Planck reduced constant,  $k_B$  is the Boltzmann constant and  $c$  is the speed of light. The Minkowski metric convention used is the “mostly minus”,  $g_{\mu\nu} = \text{diag}(1, -1, -1, -1)$ .

The  $\mathfrak{su}(2)$  basis used is the Pauli matrices,

$$\tau_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \tau_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \tau_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

They obey

$$[\tau_a, \tau_b] = 2i\varepsilon_{abc}\tau_c, \quad \{\tau_a, \tau_b\} = 2\delta_{ab}\mathbb{1}, \quad \text{Tr}[\tau_a] = 0, \quad \text{Tr}[\tau_a\tau_b] = 2\delta_{ab}\mathbb{1}.$$

Together with the identity matrix  $\mathbb{1}$ , the Pauli matrices form a basis for the vector space of all 2-by-2 matrices. An arbitrary 2-by-2 matrix  $M$  may be written

$$M = M_0\mathbb{1} + M_a\tau_a, \quad M_0 = \frac{1}{2}\text{Tr}\{M\}, \quad M_a = \frac{1}{2}\text{Tr}\{\tau_a M\}. \quad (111)$$

The gamma matrices  $\gamma^\mu$ ,  $\mu \in \{0, 1, 2, 3\}$ , obey

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}\mathbb{1}. \quad (112)$$

The “fifth  $\gamma$ -matrix” is defined by

$$\gamma^5 = \frac{i}{4!}\epsilon_{\mu\nu\rho\sigma}\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma = i\gamma^0\gamma^1\gamma^2\gamma^3. \quad (113)$$

The  $\gamma^5$ -matrix obey

$$\{\gamma^5, \gamma^\mu\} = 0, \quad (\gamma^5)^2 = \mathbb{1}, \quad \gamma^{0\dagger} = \gamma^0, \quad \gamma^{i\dagger} = -\gamma^i \quad (114)$$

Their Euclidean counterpart obey

$$\{\tilde{\gamma}_a, \tilde{\gamma}_b\} = 2\delta_{ab}\mathbb{1}, \quad \tilde{\gamma}_a^\dagger = \tilde{\gamma}_a, \quad (115)$$

and they are related by  $\tilde{\gamma}_0 = \gamma^0$ , and  $\tilde{\gamma}_j = -i\gamma^j$ . The Euclidean  $\tilde{\gamma}_5$  is defined as

$$\tilde{\gamma}_5 = \gamma_0\gamma_1\gamma_2\gamma_3 = i\gamma^0\gamma^1\gamma^2\gamma^3 = \gamma^5. \quad (116)$$

It thus also anti-commutes with the Euclidean  $\gamma$ -matrices.

### Fourier transform

The Fourier transform used in this text is defined by

$$\mathcal{F}\{f(x)\}(p) = \tilde{f}(p) = \int dx e^{ipx} f(x), \quad \mathcal{F}^{-1}\{\tilde{f}(p)\}(x) = f(x) = \int \frac{dp}{2\pi} e^{-ipx} \tilde{f}(p).$$

### Fourier series

Imaginary-time formalism is set in a Euclidean space  $\Omega = [0, \beta] \times V$ , where  $V = L_x L_y L_z$  is a space-like volume. The possible momenta in this space are

$$\tilde{V} = \left\{ \vec{k} \in \mathbb{R}^3 \mid \vec{k} = \left( \frac{2\pi n_x}{L_x}, \frac{2\pi n_y}{L_y}, \frac{2\pi n_z}{L_z} \right) \right\}$$

$\omega_n$  are the Matsubara-frequencies, with  $\omega_n = 2n\pi/\beta$  for bosons and  $\omega_n = (2n+1)\pi/\beta$  for fermions. They together form the reciprocal space  $\tilde{\Omega} = \{\omega_n\} \times \tilde{V}$ . The Euclidean coordinates are denoted  $X = (\tau, \vec{x})$  and  $K = (\omega_n, \vec{K})$ , and have the dot product  $X \cdot P = \omega_n \tau + \vec{k} \cdot \vec{x}$ . In the limit  $V \rightarrow \infty$ , we follow the prescription

$$\frac{1}{V} \sum_{\vec{p} \in \tilde{V}} \rightarrow \int_{\mathbb{R}^3} \frac{d^3 p}{(2\pi)^3}.$$

The sum over all degrees of freedom, and the corresponding integrals for the thermodynamic limit are

$$\begin{aligned} \frac{\beta V}{NM} \sum_{n=1}^N \sum_{\vec{x}_m \in V} &\xrightarrow{N, M \rightarrow \infty} \int_0^\beta d\tau \int_{\mathbb{R}^3} d^3 x = \int_{\Omega} dX, \\ \frac{1}{V} \sum_{n=-\infty}^{\infty} \sum_{\vec{k} \in \tilde{V}} &\xrightarrow{V \rightarrow \infty} \sum_{n=-\infty}^{\infty} \int_{\mathbb{R}^3} \frac{d^3 p}{(2\pi)^3} = \int_{\tilde{\Omega}} dK. \end{aligned}$$

The convention used for the Fourier expansion of thermal fields is in accordance with [8]. The prefactor is chosen to make the Fourier components of the field dimensionless, which makes it easier to evaluate the trace correctly. For bosons, the Fourier expansion is

$$\begin{aligned} \varphi(X) &= \sqrt{V\beta} \int_{\tilde{\Omega}} dK \tilde{\varphi}(K) e^{iX \cdot K} = \sqrt{\frac{\beta}{V}} \sum_{n=-\infty}^{\infty} \sum_{\vec{k} \in \tilde{V}} \tilde{\varphi}_n(\vec{p}) \exp\{i(\omega_n \tau + \vec{x} \cdot \vec{k})\}, \\ \tilde{\varphi}(K) &= \sqrt{\frac{1}{V\beta^3}} \int_{\tilde{\Omega}} dX \tilde{\varphi}(X) e^{-iX \cdot K} \end{aligned}$$

while for Fermions it is

$$\psi(X) = \sqrt{V} \int_{\tilde{\Omega}} dK \tilde{\psi}(K) e^{iX \cdot K} = \frac{1}{\sqrt{V}} \sum_{n=-\infty}^{\infty} \sum_{\vec{k} \in \tilde{V}} \psi(\omega_n, \vec{k}) \exp\{i(\omega_n \tau + \vec{x} \cdot \vec{k})\} \quad (117)$$

A often used identity is

$$\int_{\Omega} dX e^{iX \cdot (K - K')} = \beta \delta_{nn'} (2\pi)^3 \delta^3(\vec{k} - \vec{k}') := \beta \delta(K - K'), \quad (118)$$

$$\int_{\tilde{\Omega}} dK e^{iK(X - X')} = \beta \delta(\tau - \tau') \delta^3(\vec{x} - \vec{x}') := \beta \delta(X - X'). \quad (119)$$

## Bose distribution

The Bose distribution is defined as

$$n_B(\omega) = \frac{1}{e^{\beta\omega} - 1}. \quad (120)$$

This function obeys

$$n_B(-i\omega) = -1 - n_B(i\omega). \quad (121)$$

We can expand it around the Bose Matsubara frequencies on the imaginary line:

$$in_B[i(\omega_n + \epsilon)] = \frac{i}{e^{i\beta\epsilon + 2\pi i n} - 1} = i[i\beta\epsilon + \mathcal{O}(\epsilon^2)]^{-1} \sim \frac{1}{\epsilon\beta}. \quad (122)$$

This means that  $in_B(i\omega)$  has a pole on all Matsubara-frequencies, with residue  $1/\beta$ . Furthermore, we have

$$\frac{d}{d\omega} \ln(1 - e^{-\beta\omega}) = \beta n_B(\omega). \quad (123)$$

## Propagators

If  $D^{-1}[f(x)] = 0$  is the equation of motion for some field  $f$ , where  $D^{-1}$  in general is a differential operator, then the propagator  $D(x, x')$  for this field is defined by

$$D^{-1}[D(x, x')] = -i\delta(x - x')\mathbb{1}.$$

Assuming  $A$  is linear and independent of space, we may redefine  $D(x - x', 0) \rightarrow D(x - x')$ , and the Fourier transform with respect to both  $x$  and  $x'$  to obtain

$$\mathcal{F}\{D^{-1}[D(x, x')]\}(p, p') = \tilde{D}^{-1}(p) \tilde{D}(p) \delta(p + p') = -i\delta(p + p'),$$

meaning the momentum space propagator  $\tilde{D}(p) = \mathcal{F}\{D(x)\}(p)$  is given by  $\tilde{D} = -i(\tilde{D}^{-1})^{-1}$ .

For some differential operator  $D^{-1}$ , the thermal propagator is defined as

$$D^{-1}D(X, Y) = \beta\delta(X - Y). \quad (124)$$

The Fourier transformed propagator is, assuming  $D(X, Y) = D(X - Y, 0)$ ,

$$\tilde{D}(K, K') = \frac{1}{V\beta^3} \int_{\Omega} dX dY D(X, Y) \exp(-i[X \cdot K + Y \cdot K']) \quad (125)$$

$$= \frac{1}{V\beta^3} \int_{\Omega} dX' dY' D(X', 0) \exp\left(-i[X' \cdot \frac{1}{2}(K - K') + Y \cdot (K + K')]\right) \quad (126)$$

$$= \frac{1}{V\beta^3} \tilde{D}(K) \beta\delta(K + K'), \quad (127)$$

where

$$\tilde{D}(K) = \int dX e^{iK \cdot X} D(X, 0). \quad (128)$$

Thus,

$$D^{-1}D(X, Y) = \int dK \tilde{D}(K) D^{-1} e^{iK(X-Y)} = \beta\delta(X - Y) = \int dK e^{iK(X-Y)}. \quad (129)$$

The thermal propagator in momentum space,  $\tilde{D}(K)$ , is given by

$$\tilde{D}^{-1}\tilde{D}(K) = 1, \quad (130)$$

where  $\tilde{D}^{-1}$  are the eigenvalues of  $D^{-1}$ .

## Appendix B Covariant derivative

In  $\chi$ PT at finite isospin chemical potential  $\mu_I$ , the covariant derivative acts on functions  $A(x) : \mathcal{M}_4 \rightarrow \text{SU}(2)$ , where  $\mathcal{M}_4$  is the space-time manifold. It is defined as

$$\nabla_{\mu} A(x) = \partial_{\mu} A(x) - i[v_{\mu}, A(x)], \quad v_{\mu} = \frac{1}{2}\mu_I \delta_{\mu}^0 \tau_3. \quad (131)$$

The covariant derivative obeys the product rule, as

$$\nabla_{\mu}(AB) = (\partial_{\mu} A)B + A(\partial_{\mu} B) - i[v_{\mu}, AB] = (\partial_{\mu} A - i[v_{\mu}, A])B + A(\partial_{\mu} B - i[v_{\mu}, B]) = (\nabla_{\mu} A)B + A(\nabla_{\mu} B).$$

Decomposing a 2-by-2 matrix  $M$ , as described in Appendix A, shows that the trace of the commutator of  $\tau_b$  and  $M$  is zero:

$$\text{Tr}\{[\tau_a, M]\} = M_b \text{Tr}\{[\tau_a, \tau_b]\} = 0.$$

Together with the fact that  $\text{Tr}\{\partial_{\mu} A\} = \partial_{\mu} \text{Tr}\{A\}$ , this gives the product rule for invariant traces:

$$\text{Tr}\{A \nabla_{\mu} B\} = \partial_{\mu} \text{Tr}\{AB\} - \text{Tr}\{(\nabla_{\mu} A)B\}.$$

This allows for the use of the divergence theorem when doing partial integration. Let  $\text{Tr}\{K^\mu\}$  be a space-time vector, and  $\text{Tr}\{A\}$  scalar. Let  $\Omega$  be the domain of integration, with coordinates  $x$  and  $\partial\Omega$  its boundary, with coordinates  $y$ . Then,

$$\int_{\Omega} dx \text{Tr}\{A \nabla_{\mu} K^{\mu}\} = \int_{\partial\Omega} dy n_{\mu} \text{Tr}\{A K^{\mu}\} - \int_{\Omega} dx \text{Tr}\{(\nabla_{\mu} A) K^{\mu}\},$$

where  $n_{\mu}$  is the normal vector of  $\partial\Omega$ . [7] This makes it possible to do partial integration and discard surface terms in the  $\chi$ PT Lagrangian, given the assumption of no variation on the boundary.