

An implicit, energy-conserving and asymptotic-preserving time integrator for particle-in-cell simulation of magnetized plasmas

Lee F. Ricketson

In collaboration with: Luis Chacón (LANL)

Lawrence Livermore National Laboratory

Presented at Geometric Algorithms and Methods in Physics
June 24, 2021

This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

Context:

- Kinetic equations for plasmas
- Particle in Cell methods
 - Recent implicit version
- Motion in a strong magnetic field

Implicit, AP time integrator:

- “Effective force” to capture ∇B drift
- Considerable care needed to conserve energy
- Time-step restrictions, FLR effects, and adaptivity

Numerical Examples:

- Single particle motion in magnetic mirror, tokamak equilibrium, etc.

Kinetic Plasma Simulation

We're interested in solving

$$\partial_t f + \mathbf{v} \cdot \nabla_{\mathbf{x}} f + \frac{q}{m} (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \nabla_{\mathbf{v}} f = 0 \quad (1)$$

for $f(\mathbf{x}, \mathbf{v}, t)$, particularly in cases for which the magnetic field \mathbf{B} may be “very large”.

High-dimensionality motivates **particle-based** methods: Postulate


$$f \approx \sum_p S(\mathbf{x} - \mathbf{x}_p) \delta(\mathbf{v} - \mathbf{v}_p), \quad (2)$$

so each “particle” traces characteristics:

$$\dot{\mathbf{x}}_p = \mathbf{v}_p, \quad \dot{\mathbf{v}}_p = \frac{q}{m} (\mathbf{E} + \mathbf{v}_p \times \mathbf{B}) \quad (3)$$

Computing \mathbf{E}, \mathbf{B} directly via particle-particle interactions is an $O(N_p^2)$ operation (at least, without FMM), so...

PIC Summary



- Interpolate particle properties onto grid
- Compute \mathbf{E}, \mathbf{B} on grid (FEM, FFT, FMM, etc.)
- Interpolate force back onto particle positions
- Update particle \mathbf{x}, \mathbf{v} using fields:

$$\mathbf{v}_{n+1}^p = \mathbf{v}_n^p + (\mathbf{E}^p + \mathbf{v}^p \times \mathbf{B}_p) \Delta t$$

$$\mathbf{x}_{n+1}^p = \mathbf{x}_n^p + \mathbf{v}_{n+1}^p \Delta t$$

Work of Chen, Chacón and others shows that a Crank-Nicolson particle push

$$\begin{aligned}\mathbf{x}^{n+1} &= \mathbf{x}^n + \Delta t \mathbf{v}^{n+1/2}, \\ \mathbf{v}^{n+1} &= \mathbf{v}^n + \Delta t \frac{q}{m} \left[\mathbf{E}^{n+1/2} + \mathbf{v}^{n+1/2} \times \mathbf{B}^{n+1/2} \right]\end{aligned}\tag{4}$$

coupled to a carefully chosen field solve and their novel technique of *particle enslavement* leads to an implicit PIC scheme that is


- Computationally feasible
- Exactly energy conserving *for all* Δt , Δx
- Resistant to the finite-grid instability
- Decouples particle and field time-steps
- **BUT** does not capture guiding center motion for $\Omega_c \Delta t \gg 1$

Fast Gyration

- Aside from high-dimensionality, plasmas suffer from a zoo of disparate time-scales
- We focus on a particular time-scale gap here:
 - In strong magnetic fields, gyration about the field line has frequency

$$\Omega_c = \frac{qB}{m} \quad (5)$$

that is faster than anything else of interest.



The Traditional Approach: (Drift/Gyro)kinetics

The traditional approach - called (drift/gyro)kinetics - is:

- Analytically remove the fast time-scale from the governing equations by asymptotic expansion
- Numerically solve the new, reduced equations

This has been very successful, but presents two important problems:

- Boundary conditions
- What if $\Omega_c^{-1} \ll \tau_{\text{interesting}}$ is only true in a *portion* of the domain? e.g. tokamak edge

Asymptotic Preserving (AP) Schemes

Asymptotic preserving (AP) schemes take a different approach to confronting the fast scale:

- Numerically simulate the full, original governing equations
- **BUT** make sure your numerical scheme respects asymptotic limit(s)
- Then, you can take large time-steps compared to fast scale

This avoids both previous issues:

- Boundary conditions are just those for the original equations
- Simple to step over fast scale only when justifiable

Single particle motion

Single particle motion in prescribed electromagnetic fields satisfies

$$\begin{aligned}\frac{d\mathbf{x}}{dt} &= \mathbf{v}, \\ \frac{d\mathbf{v}}{dt} &= \frac{q}{m} (\mathbf{E} + \mathbf{v} \times \mathbf{B}).\end{aligned}\tag{6}$$

When gyroradius \ll length-scales and $\Omega_c^{-1} \ll$ time-scales, asymptotic expansion gives

$$\mathbf{v} \approx \mathbf{u} + v_{\parallel} \mathbf{b} + \mathbf{v}_E + \mathbf{v}_I + \mathbf{v}_B.\tag{7}$$

- \mathbf{u} = Gyration about field line
- $\mathbf{v}_E = \mathbf{E} \times \mathbf{B} / B^2$.
- \mathbf{v}_I = slow drifts induced by changes in \mathbf{v}_E and $\mathbf{b} = \mathbf{B} / B$.
- \mathbf{v}_B = slow drift induced by spatial variation in B .

Existing integrators

Properties of existing 2nd-order time-stepping schemes for $\Omega_c \Delta t \gg 1$:

	Boris/Leapfrog	Crank-Nicolson	BFV ¹	MI ²
Energy conserving		✓		
Correct gyroradius		✓	✓	✓
$\mathbf{E} \times \mathbf{B}$ drift	✓	✓	✓	✓
Inertial drifts	✓	✓	✓	✓
∇B drift	✓		✓	✓

Our goal: A scheme that checks all the boxes!

1: Brackbill/Forslund '85, Vu/Brackbill '90

2: Genoni et al, '10

Starting point: BFV

- Fundamentally, ∇B drift and mirror force arise from “effective” force acting on the guiding center $\mathbf{F} = -\mu \nabla B$ (where $\mu = mu^2/2B$) that C-N misses for $\Omega_c \Delta t \gg 1$.
- BFV attempts to put that force back in:

$$\mathbf{v}^{n+1} = \mathbf{v}^n + \Delta t \frac{q}{m} \left[\mathbf{E}^{n+1/2} + \mathbf{v}^{n+1/2} \times \mathbf{B}^{n+1/2} + \mathbf{F}_{BFV} \right], \quad (8)$$

where

$$\mathbf{F}_{BFV} = -\frac{m \|\mathbf{v}_{\perp}^{n+1} - \mathbf{v}_{\perp}^n\|^2}{8B^{n+1/2}} \nabla B^{n+1/2} \quad (9)$$

- Satisfies

$$\begin{aligned} \Omega_c \Delta t \ll 1 &\implies \mathbf{F}_{BFV} = O(\Delta t^2), \\ \Omega_c \Delta t \gg 1 &\implies \mathbf{F}_{BFV} \approx -\mu \nabla B. \end{aligned} \quad (10)$$

BUT: $\mathbf{F}_{BFV} \cdot \mathbf{v}^{n+1/2} \neq 0$, breaking energy conservation!

Overconstrained

One would really like an effective force \mathbf{F}_{eff} satisfying

- $\Omega_c \Delta t \ll 1 \implies \mathbf{F}_{eff} = O(\Delta t^2)$
- $\Omega_c \Delta t \gg 1 \implies \mathbf{F}_{eff} \approx -\mu \nabla B$
- $\mathbf{F}_{eff} \cdot \mathbf{v}^{n+1/2} = 0$ for all Δt

Unfortunately, the last two constraints are **incompatible**:

$$\nabla B \cdot \mathbf{v}^{n+1/2} \neq 0 \quad (11)$$

We can settle for a weaker version of the second constraint:

$$\mathbf{F}_{eff} \approx -\mu \nabla B \longrightarrow \langle \mathbf{F}_{eff} \rangle \approx -\mu \nabla B,$$

where $\langle \cdot \rangle$ denotes average over gyration.

The idea: Time-stepping samples a variety of gyrophases for us, so the average of \mathbf{F}_{eff} over several time-steps approximates the gyroaverage operator for us. We call this “**implicit gyroaveraging**”.

Conservative Effective Force

We postulate an effective force of the form

$$\mathbf{F}_{eff} = \left(\mathbf{b} - \frac{v_{\parallel} \mathbf{v}_{\perp}}{v_{\perp}^2} \right) G_{\parallel} + \left(\mathbf{I} - \frac{\mathbf{v}_{\perp} \mathbf{v}_{\perp}}{v_{\perp}^2} \right) \cdot \mathbf{G}_{\perp} \quad (12)$$

- Conserves energy $\forall \mathbf{G}$ and avoids mixing parallel and perpendicular motion

Constraints + algebra \implies form of \mathbf{G} :

$$\begin{aligned} \mathbf{G} = & F_{BFV,\parallel} \mathbf{b} + \frac{(\mathbf{I} - \hat{\mathbf{v}}_E \hat{\mathbf{v}}_E) \cdot \mathbf{F}_{BFV,\perp}}{1 - \eta^2/2} \\ & + \frac{2}{\eta^2} \hat{\mathbf{v}}_E \left(\hat{\mathbf{v}}_E \cdot \mathbf{F}_{BFV,\perp} + \mathbb{1}_{v_E > u} F_{BFV,\parallel} \frac{v_{\parallel}}{v_E} \right) \end{aligned} \quad (13)$$

where $\eta = \min\{1, u/v_E\}$, $\hat{\mathbf{v}}_E = \mathbf{v}_E/v_E$.

Going Further: FLR Effects

Scheme just derived captures *drift*-kinetic limit. To get *gyrokinetic*, need to allow *finite Larmor radius* (FLR) effects:

$$\left(\frac{\|\nabla \mathbf{E}\|}{E} \right)^{-1} \sim \text{gyroradius}. \quad (14)$$


- Changes to asymptotic limit expression: $\mathbf{E} \longrightarrow \langle \mathbf{E} \rangle$
- Challenge for scheme: \mathbf{E} has structure, so can't sample *random* gyrophases, must force time-stepping to sample *equispaced* gyrophases

Going further: FLR effects

Two changes to scheme required:

- Alternate large ($\Omega_c \Delta t \gg 1$) and smaller ($\Omega_c \delta t \sim 1$) time-steps to rig appropriate gyrophase traversal
- Change \mathbf{E} evaluation from midpoint to trapezoidal so evaluation is on gyro-orbit, not at gyrocenter

$$\mathbf{E}^{n+1/2} = \mathbf{E} \left(\frac{\mathbf{x}^n + \mathbf{x}^{n+1}}{2} \right) \longrightarrow \frac{\mathbf{E}(\mathbf{x}^n) + \mathbf{E}(\mathbf{x}^{n+1})}{2}$$



Time-step Restrictions

Of course, one still cannot take arbitrarily large time-steps. Must resolve spatiotemporal variation in \mathbf{E} , \mathbf{B} ... but also:

- The time it takes for **implicit gyroaverage** to be computed accurately must be small
- Anomalous displacement due to inaccuracy in **implicit gyroaverage** must be small

Time-step Restrictions

Quantitative expressions for the maximum allowable $\Omega_c \Delta t$ are challenging to find in general, but can be done in two important limits:

	Displacement restriction	Avg. restriction
$u^{n+1/2} \gg v_E$	$\min \{ 2(\delta_\perp)^{-1}, \sqrt{2}(\delta_\parallel)^{-1/2} \}$	$2 \frac{\Omega_c \tau_{res}}{\pi}$
$u^{n+1/2} \ll v_E$	$\sqrt{2} (\delta_E + \delta_\parallel)^{-1/2}$	$\Omega_c \tau_{res}$

where τ_{res} is the smallest time-scale you wish to resolve,

$$\delta_\perp = \frac{u}{\Omega_c} \frac{\|(\nabla B)_\perp\|}{B}, \quad \delta_\parallel = \frac{v_\parallel}{\Omega_c} \frac{\|(\nabla B)_\parallel\|}{B}, \quad \delta_E = \frac{v_E}{\Omega_c} \frac{\|(\nabla B)_\perp\|}{B}. \quad (15)$$

These restrictions inform adaptive time-stepping procedure.

So far, scheme primarily tested on single-particle motion... Several different field configurations:


- Magnetic mirror
- Non-adiabatic jump in μ
- Tokamak equilibrium with electric field

Implementation in self-consistent implicit PIC scheme is in progress with promising initial results.

Magnetic Mirror

Configured so that particles with $v_{\parallel} < v_{crit}$ are confined, others are “passing”.

$$\underline{v_{\parallel} = 0.999 v_{crit}}$$




New scheme predicts trapped passing boundary to within 0.05%

Magnetic Mirror

Configured so that particles with $v_{\parallel} < v_{crit}$ are confined, others are “passing”.


$$\underline{v_{\parallel} = 0.98 v_{crit}}$$



Also get improved prediction of “bounce frequency”.



Unmagnetized Region

$$\mathbf{E} = 2\hat{\mathbf{y}}, \quad \mathbf{B} = \hat{\mathbf{z}}\sqrt{1 + 4(x - 12)^2}$$




$\mathbf{E} \times \mathbf{B}$ drift pushes particle through unmagnetized region...

Unmagnetized Region



- Size of jump in μ is *very* sensitive to initial gyrophase
- Can't get jump right for any particular gyrophase b/c $\Omega_c \Delta t > 1$ inherently misses gyrophase information
- **BUT** can hope to get the jump correct **on average**


Unmagnetized Region



- Reproduce gyroaveraged μ jump within 0.8%
- Standard deviation within 14%
- (drift/gyro)kinetics would predict **no** change in μ

Tokamak equilibrium with **E** field


- **B** is a Solov'ev equilibrium field + toroidal component $\propto 1/r$
- **E** has $k_{\perp}\rho \approx 1.2$



New version captures small FLR corrections

Tokamak equilibrium with \mathbf{E} field - Even larger $k_{\perp}\rho$

- \mathbf{B} is a Solov'ev equilibrium field + toroidal component $\propto 1/r$
- \mathbf{E} has $k_{\perp}\rho \approx 2.4$



FLR corrections still improve accuracy, but working on sensitivity to adaptive time-stepping

Conclusions

- First-of-kind scheme that both preserves asymptotic properties and conserves energy
- An alternative to (drift/gyro)kinetic PIC that can handle arbitrary magnetization
- Energy conservation is **not** just pedantry, has **qualitative** consequences on orbits
- Work still to-do:
 - Continue in-progress implementation in implicit PIC schemes
 - Improve preconditioning of implicit solve
 - Further refinement/generalization of adaptive time-stepping

For more detail, see: L.F. Ricketson, L. Chacón, "An energy-conserving and asymptotic-preserving charged-particle orbit implicit time integrator for arbitrary electromagnetic fields." *JCP* (2020): 109639



Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.