



Co-funded by
the European Union

Degenerate Variational Integrators Variations on a Motif

Michael Kraus (michael.kraus@ipp.mpg.de)

Acknowledgements: Joshua Burby, Leland Ellison, Omar Maj, Hiroaki Yoshimura, ...

Outline


1. Degenerate Lagrangian Systems
2. Variational Integrators
3. Degenerate Variational Integrators
4. Discontinuous Galerkin DVIs
5. Remarks on Projection Methods
6. Summary and Outlook

Degenerate Lagrangian Systems

Hamilton's Principle of Stationary Action

- Physical equations are often derived from an action principle
- The action \mathcal{A} is a functional of a trajectory $q : [0, T] \rightarrow \mathbb{R}^m$

$$\mathcal{A}[q] = \int_0^T L(q(t), \dot{q}(t)) dt$$




Hamilton's principle of stationary action

Among all possible trajectories connecting q_0 and q_T , the physical trajectory makes the action integral \mathcal{A} stationary.

Hamilton's Principle of Stationary Action

- computing variations

$$\delta \mathcal{A}[q] = \int_0^T \left[\frac{\partial L}{\partial q} \cdot \delta q + \frac{\partial L}{\partial \dot{q}} \cdot \delta \dot{q} \right] dt = 0 \text{ for all } \delta q$$



- integration by parts (endpoints fixed: $\delta q(0) = \delta q(T) = 0$)

$$\delta \mathcal{A}[q] = \int_0^T \left[\frac{\partial L}{\partial q} - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \right) \right] \cdot \delta q dt + \left[\frac{\partial L}{\partial \dot{q}} \cdot \delta q \right]_0^T = 0 \text{ for all } \delta q$$

- requiring stationarity of the action leads to the Euler-Lagrange equations of motion

$$\frac{\partial L}{\partial q}(q, \dot{q}) - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}}(q, \dot{q}) \right) = 0$$

Regular Lagrangians

- for regular Lagrangians, the velocity space Hessian M is invertible for each (q, \dot{q})

$$M_{ij}(q, \dot{q}) = \frac{\partial^2 L}{\partial \dot{q}^i \partial \dot{q}^j}(q, \dot{q})$$

- this implies that the Euler–Lagrange equations are a system of m 2nd-order ODEs

$$\ddot{q}^i = M_{ij}^{-1} \left(\frac{\partial L}{\partial q^j} - \frac{\partial^2 L}{\partial \dot{q}^j \partial q^k} \dot{q}^k \right)$$

following from

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}^i}(q, \dot{q}) \right) = M_{ij} \ddot{q}^j + \frac{\partial^2 L}{\partial \dot{q}^i \partial q^k} \dot{q}^k$$

- the solution of this system of equations requires $2m$ initial conditions

Phasespace Lagrangians

- in the following, we consider phasespace Lagrangians, which have the general form

$$L(q, \dot{q}) = \vartheta(q) \cdot \dot{q} - H(q)$$

- as L is linear in velocities the velocity-space Hessian is zero everywhere
- the Euler-Lagrange equations are first order ordinary differential equations

$$\frac{d}{dt}\vartheta(q) = \nabla\vartheta(q) \cdot \dot{q} - \nabla H(q) \quad \text{or} \quad \dot{q}^i \omega_{ij}(q) = \frac{\partial H}{\partial q^i}(q)$$

with the $m \times m$ noncanonical symplectic matrix ω given by

$$\omega_{ij} = \frac{\partial \vartheta_i}{\partial q^j} - \frac{\partial \vartheta_j}{\partial q^i}$$

- the solution of this system of equations requires m initial conditions

Variational Integrators

Discrete Variational Principle

- discrete action and discrete Lagrangian with discrete trajectory $q_d = \{q_n\}_{n=0}^N$

$$\mathcal{A}_d[q_d] = \sum_{n=0}^{N-1} L_d(q_n, q_{n+1}) \quad \text{e.g.} \quad L_d(q_n, q_{n+1}) = h L\left(\frac{q_n + q_{n+1}}{2}, \frac{q_{n+1} - q_n}{h}\right)$$

- requiring stationarity of the discrete action,

$$\delta \mathcal{A}_d[q_d] = \delta \sum_{n=0}^{N-1} L_d(q_n, q_{n+1}) = 0 \quad \text{for all} \quad \delta q_n,$$

with $\delta q_0 = \delta q_N = 0$, leads to the discrete Euler-Lagrange equations

$$D_2 L_d(q_{n-1}, q_n) + D_1 L_d(q_n, q_{n+1}) = 0 \quad \text{for all} \quad n$$

Discrete Variational Principle

- provided the discrete Hessian

$$M_{ij}(q_n, q_{n+1}) = \frac{\partial L_d}{\partial q_n^i \partial q_{n+1}^j}(q_n, q_{n+1})$$

is invertible, the discrete Euler-Lagrange equations define a mapping

$$\Psi_{L_d} : (q_{n-1}, q_n) \mapsto (q_n, q_{n+1}^*(q_{n-1}, q_n))$$

- solving the discrete Euler-Lagrange equations requires $2m$ initial conditions

→ same number of initial conditions required to solve the continuous Euler-Lagrange equations for a regular Lagrangian

Discrete Variational Principle and Phasespace Lagrangians

- for phasespace Lagrangians

$$L(q, \dot{q}) = \vartheta(q) \cdot \dot{q} - H(q)$$


the mapping

$$\Psi_{L_d} : (q_{n-1}, q_n) \mapsto (q_n, q_{n+1}^*(q_{n-1}, q_n))$$

corresponds to a multi-step variational integrator

- the variational integrator requires $2m$ initial conditions even though the continuous Euler–Lagrange equations (being first order ODEs) require only m initial conditions
- susceptible to parasitic modes driving simulations unstable

Lotka–Volterra System with Variational Integrator



Discrete Variational Principle and Phasespace Lagrangians

Problem

Most discrete Lagrangians L_d approximating a phasespace Lagrangian have an invertible discrete Hessian M .

Degenerate Variational Integrators

Special Phasespace Lagrangians

- consider the special phasespace Lagrangian

$$L(q, \dot{q}) = \vartheta(q) \cdot \dot{q} - H(q), \quad q \in \mathbb{R}^{2d},$$

where $d = m/2$ of the components of ϑ vanish, in particular

$$\vartheta_\mu = 0 \quad \text{for} \quad \mu = d+1, \dots, 2d$$

- for clarity sake we will consider a phasespace Lagrangian with $d = 1$

$$L(q^1, q^2, \dot{q}^1, \dot{q}^2) = \vartheta_1(q^1, q^2) \dot{q}^1 - H(q^1, q^2)$$

Degenerate Variational Integrators

- consider the discrete degenerate Lagrangian

$$L_d(q_n, q_{n+1}) = h \left[\vartheta_1(q_n^1, q_n^2) \frac{q_{n+1}^1 - q_n^1}{h} - H(q_n^1, q_n^2) \right] = 0$$

- the discrete Hessian is degenerate (rank $d = 1$)
- discrete Euler–Lagrange equations look like a multi-step method

$$\frac{\vartheta_1(q_n^1, q_n^2) - \vartheta_1(q_{n-1}^1, q_{n-1}^2)}{h} = D_1 \vartheta_1(q_n^1, q_n^2) \frac{q_{n+1}^1 - q_n^1}{h} + D_1 H(q_n^1, q_n^2)$$

for $n = 1, \dots, N-1$,

$$0 = D_2 \vartheta_1(q_n^1, q_n^2) \frac{q_{n+1}^1 - q_n^1}{h} + D_2 H(q_n^1, q_n^2)$$

for $n = 0, \dots, N-1$

Degenerate Variational Integrators

- for sake of clarity introduce a velocity

$$v_n^1 = \frac{q_{n+1}^1 - q_n^1}{h},$$

→ assuming appropriate invertibility conditions, v can be solved for by

$$v_n^1 = \left(D_2 \vartheta_1(q_n^1, q_n^2) \right)^{-1} D_2 H(q_n^1, q_n^2)$$

→ the discrete Euler–Lagrange equations become

$$\vartheta_1(q_n^1, q_n^2) = \vartheta_1(q_{n-1}^1, q_{n-1}^2) + h D_1 \vartheta_1(q_n^1, q_n^2) v_n^1 - h D_1 H(q_n^1, q_n^2) \quad \text{for } n = 1, \dots, N-1$$

$$v_n^1 = \left(D_2 \vartheta_1(q_n^1, q_n^2) \right)^{-1} D_2 H(q_n^1, q_n^2) \quad \text{for } n = 0, \dots, N-1$$

$$q_{n+1}^1 = q_n^1 + h v_n^1 \quad \text{for } n = 0, \dots, N-1$$

→ underdetermined system of equations: no functional dependence on q_N^2

Degenerate Variational Integrators

- close the discrete Euler–Lagrange equations by

$$\vartheta_1(q_N^1, q_N^2) = \vartheta_1(q_{N-1}^1, q_{N-1}^2) + hD_1\vartheta_1(q_N^1, q_N^2) \cdot v_N^1 - hD_1H(q_N^1, q_N^2),$$
$$v_N^1 = \left(D_2\vartheta_1(q_N^1, q_N^2)\right)^{-1} D_2H(q_N^1, q_N^2)$$

→ motivated by the discrete symplecticity condition following from the boundary values in the action principle

$$\delta\mathcal{A}_d[q_{DEL}] = -\left[\vartheta_1(q_0) - D_1\vartheta_1(q_0) \cdot v_0^1 + D_1H(q_0)\right] \delta q_0^1 + \left[\vartheta_1(q_{N-1})\right] \delta q_N^1$$

→ conservation of the discrete symplectic structure $\omega_d = d\vartheta_d$ with

$$\vartheta_d(q_n) = \left[\vartheta_1(q_n^1, q_n^2) - \nabla_1\vartheta_1(q_n^1, q_n^2) \cdot v_n^1 + \nabla_1H(q_n^1, q_n^2)\right] dq_n^1$$

Degenerate Variational Integrators

- allows to obtain one-step methods for degenerate Lagrangians directly from a discrete action principle
- alternative derivation from discrete Hamilton–Pontryagin principle that includes the velocities in the formulation of the action principle
- first order accurate
- not composable: adjoint methods preserve a different symplectic structure
- preservation of a discrete symplectic structure

Degenerate Variational Integrators

2nd-Order Leapfrog DVIs

Leapfrog Degenerate Variational Integrators

- centred, staggered discretisations of the Lagrangian
- midpoint discretisation (MDVI)

$$L_d(q_n^1, q_{n+1/2}^2, q_{n+1}^1) = h \left[\vartheta_1 \left(\frac{q_n^1 + q_{n+1}^1}{2}, q_{n+1/2}^2 \right) \frac{q_{n+1}^1 - q_n^1}{h} - H \left(\frac{q_n^1 + q_{n+1}^1}{2}, q_{n+1/2}^2 \right) \right]$$

- trapezoidal discretisation (TDVI)

$$L_d(q_n^1, q_{n+1/2}^2, q_{n+1}^1) = \frac{h}{2} \left[\left(\vartheta_1(q_n^1, q_{n+1/2}^2) + \vartheta_1(q_{n+1}^1, q_{n+1/2}^2) \right) \frac{q_{n+1}^1 - q_n^1}{h} - H(q_n^1, q_{n+1/2}^2) - H(q_{n+1}^1, q_{n+1/2}^2) \right]$$


- the discrete Hessian is degenerate (rank $d = 1$)

Leapfrog Degenerate Variational Integrators

- allows to obtain one-step methods for degenerate Lagrangians directly from a discrete action principle
- second order accurate
- composable: symmetric and self-adjoint
- requires processing for initialisation of half steps
- obtaining the genuine one-step form of the integrator not always straightforward
- preservation of a discrete symplectic structure

Discontinuous Galerkin DVIs

Discontinuous Galerkin Approximation



- piecewise-continuous polynomial basis $q_h|_{\mathcal{I}_n} \in \mathbb{P}^r(\mathcal{I}_n)$ with $\mathcal{I}_n = (t_n, t_{n+1})$
- allow for discontinuities at t_n and t_{n+1}

Discontinuous Galerkin Degenerate Variational Integrators

- consider a Galerkin discretisation of the Lagrangian, formally written as

$$L_h(q_n^+, q_{n+1}^-) = h \sum_{i=1}^s b_i L(q_h(t_n + c_i h), \dot{q}_h(t_n + c_i h)),$$

where q_h is a piecewise polynomial approximation of the trajectory q with

$$q_n^+ = \lim_{t \downarrow t_n} q_h(t), \quad q_{n+1}^- = \lim_{t \uparrow t_{n+1}} q_h(t)$$

- the following construction works in a similar way also for Runge–Kutta discretisations

Discontinuous Galerkin Degenerate Variational Integrators

- the discontinuous approximation of the Lagrangian needs to be connected by appropriate jump terms

$$L_d(q_n, q_{n+1}) = L_h(q_n^+, q_{n+1}^-) + \vartheta(\langle\langle q \rangle\rangle_n^+) \cdot \llbracket q \rrbracket_n^+ + \vartheta(\langle\langle q \rangle\rangle_{n+1}^-) \cdot \llbracket q \rrbracket_{n+1}^- ,$$

with averages and jumps defined as

$$\begin{aligned} \langle\langle q^\mu \rangle\rangle_n^+ &= \begin{cases} q_n^{+, \mu} & \text{for } \mu = 1, \dots, d, \\ q_n^\mu & \text{for } \mu = d+1, \dots, 2d, \end{cases} & \llbracket q \rrbracket_n^+ &= q_n^+ - q_n, \\ \langle\langle q^\mu \rangle\rangle_n^- &= \begin{cases} q_n^{-, \mu} & \text{for } \mu = 1, \dots, d, \\ q_n^\mu & \text{for } \mu = d+1, \dots, 2d, \end{cases} & \llbracket q \rrbracket_n^- &= q_n - q_n^- \end{aligned}$$

- the q_n are varied as independent variables

Discontinuous Galerkin Degenerate Variational Integrators

- the discontinuous Galerkin Lagrangian

$$L_d(q_n, q_{n+1}) = L_h(q_n^+, q_{n+1}^-) + \vartheta(\llbracket q \rrbracket_n^+) \cdot (q_n^+ - q_n) + \vartheta(\llbracket q \rrbracket_{n+1}^-) \cdot (q_{n+1} - q_{n+1}^-)$$

is guaranteed to be degenerate as the discrete Hessian is degenerate (rank 0)

- discrete Euler–Lagrange equations

$$\delta q_n^1: \quad 0 = \vartheta_1(\llbracket q \rrbracket_n^+) - \vartheta_1(\llbracket q \rrbracket_n^-),$$

$$\delta q_n^2: \quad 0 = D_2 \vartheta_1(\llbracket q \rrbracket_n^{+,1}) \cdot (q_n^{+,1} - q_n^1) + D_2 \vartheta_1(\llbracket q \rrbracket_n^{-,1}) \cdot (q_n^1 - q_n^{-,1})$$

plus discrete Euler–Lagrange equations resulting from variations w.r.t. internal variables of $L_h(q_n^+, q_{n+1}^-)$

Discontinuous Galerkin Degenerate Variational Integrators

- discrete Euler–Lagrange equations from jump terms

$$\delta q_n^1: \quad 0 = \vartheta_1(q_n^{+,1}, q_n^2) - \vartheta_1(q_n^{-,1}, q_n^2),$$

$$\delta q_n^2: \quad 0 = D_2 \vartheta_1(q_n^{+,1}, q_n^2) \cdot (q_n^{+,1} - q_n^1) + D_2 \vartheta_1(q_n^{-,1}, q_n^2) \cdot (q_n^1 - q_n^{-,1})$$

- for sufficiently small time steps and some mild assumptions on ϑ the first equation implies $\langle\langle q \rangle\rangle_n^+ = \langle\langle q \rangle\rangle_n^-$ and thus $q_n^{+,1} = q_n^{-,1}$
 - this this, the second equation trivialises, leaving us with a system that is under-determined by Nd equations
- allows us to add the continuity condition $q_n^1 = q_n^{+,1} = q_n^{-,1}$

Discontinuous Galerkin Degenerate Variational Integrators

- the boundary terms of the discrete action principle read

$$d\mathcal{A}_d[q_{DEL}] = -\vartheta(\langle\langle q \rangle\rangle_0^+) \cdot dq_0 + \vartheta(\langle\langle q \rangle\rangle_N^-) \cdot dq_N$$

- identifying q_n^1 with $q_n^{+,1} = q_n^{-,1}$, this becomes

$$d\mathcal{A}_d[q_{DEL}] = -\vartheta(q_0) \cdot dq_0 + \vartheta(q_N) \cdot dq_N$$

→ DG-DVIs preserve the continuous symplectic structure

Discontinuous Galerkin Degenerate Variational Integrators

- no closed-form expression for degenerate discrete Lagrangians:
obtaining one-step methods requires implicit arguments
(but is otherwise straightforward)
- arbitrary order accurate
- composable: includes symmetric and self-adjoint methods
- no processing for initialisation required
- preservation of the continuous symplectic structure

Discontinuous Galerkin DVIs

Some Examples

Hamilton–Pontryagin Principle

- Hamilton–Pontryagin principle: action principle on $T\mathcal{M} \oplus T^*\mathcal{M}$

$$\delta \int_0^T \left[L(q, v) + \langle p, \dot{q} - v \rangle \right] dt = 0$$

- variations of v are left free, a kinematic constraint ensures the second-order condition $v = \dot{q}$ with the momentum p as a Lagrange multiplier
(Hamilton's action principle: variations $\delta \dot{q}$ are induced by variations δq)
- requiring stationarity of the Hamilton–Pontryagin action, leads to the implicit Euler–Lagrange equations

$$\dot{q} = v, \quad p = \frac{\partial L}{\partial v}, \quad \dot{p} = \frac{\partial L}{\partial q}$$

(second-order condition, fibre derivative, Euler-Lagrange equations)

First-Order DVIs (Variant A)

- piecewise linear polynomial approximation of q

$$q_h(t)|_{(t_n, t_{n+1})} = \frac{t_{n+1} - t}{t_{n+1} - t_n} q_n^+ + \frac{t - t_n}{t_{n+1} - t_n} q_{n+1}^-$$



- continuous linear polynomial approximation of v

$$v_h(t)|_{[t_n, t_{n+1}]} = \frac{t_{n+1} - t}{t_{n+1} - t_n} v_n + \frac{t - t_n}{t_{n+1} - t_n} v_{n+1}$$

- use left Riemann quadrature on L_h and right Riemann quadrature on the Pontryagin constraint

$$\begin{aligned} L_d(q_n, q_{n+1}) = & h \left[\vartheta_1(q_n^{+,1}, q_n^{+,2}) \cdot v_n^1 - H(q_n^{+,1}, q_n^{+,2}) \right] + h p_{n+1}^1 \cdot \left[\frac{q_{n+1}^{-,1} - q_n^{+,1}}{h} - v_{n+1}^1 \right] \\ & + \vartheta_1(q_n^{+,1}, q_n^2) \cdot (q_n^{+,1} - q_n^1) + \vartheta_1(q_{n+1}^{-,1}, q_{n+1}^2) \cdot (q_{n+1}^1 - q_{n+1}^{-,1}) \end{aligned}$$

Lotka–Volterra System with DVI1A



First-Order DVIs (Variant B)

- piecewise linear polynomial approximation of q

$$q_h(t)|_{(t_n, t_{n+1})} = \frac{t_{n+1} - t}{t_{n+1} - t_n} q_n^+ + \frac{t - t_n}{t_{n+1} - t_n} q_{n+1}^-$$



- continuous linear polynomial approximation of v

$$v_h(t)|_{[t_n, t_{n+1}]} = \frac{t_{n+1} - t}{t_{n+1} - t_n} v_n + \frac{t - t_n}{t_{n+1} - t_n} v_{n+1}$$

- use right Riemann quadrature on L_h and left Riemann quadrature on the Pontryagin constraint

$$\begin{aligned} L_d(q_n, q_{n+1}) = & h \left[\vartheta_1(q_{n+1}^{-,1}, q_{n+1}^{-,2}) \cdot v_{n+1}^1 - H(q_{n+1}^{-,1}, q_{n+1}^{-,2}) \right] + h p_n^1 \cdot \left[\frac{q_{n+1}^{-,1} - q_n^{+,1}}{h} - v_n^1 \right] \\ & + \vartheta_1(q_n^{+,1}, q_n^{+,2}) \cdot (q_n^{+,1} - q_n^1) + \vartheta_1(q_{n+1}^{-,1}, q_{n+1}^{+,2}) \cdot (q_{n+1}^1 - q_{n+1}^{-,1}) \end{aligned}$$

Lotka–Volterra System with DVI1B



Trapezoidal DVI

- the approximation of q is given by

$$q_h^1(t)|_{(t_n, t_{n+1})} = \frac{t_{n+1} - t}{t_{n+1} - t_n} q_n^{+,1} + \frac{t - t_n}{t_{n+1} - t_n} q_{n+1}^{-,1},$$

$$q_h^2(t)|_{(t_n, t_{n+1})} = q_{n+1/2}^2$$

- use trapezoidal quadrature on the Lagrangian

$$\begin{aligned} L_d(q_n, q_{n+1}) = & \frac{h}{2} \left[\left(\vartheta_1(q_n^{+,1}, q_{n+1/2}^2) + \vartheta_1(q_{n+1}^{-,1}, q_{n+1/2}^2) \right) \cdot v_{n+1/2}^1 - H(q_n^{+,1}, q_{n+1/2}^2) - H(q_{n+1}^{-,1}, q_{n+1/2}^2) \right] \\ & + h p_{n+1/2}^1 \cdot \left[\frac{q_{n+1}^{-,1} - q_n^{+,1}}{h} - v_{n+1/2}^1 \right] + \vartheta_1(q_n^{+,1}, q_n^2) \cdot (q_n^{+,1} - q_n^1) + \vartheta_1(q_{n+1}^{-,1}, q_{n+1}^2) \cdot (q_{n+1}^1 - q_{n+1}^{-,1}) \end{aligned}$$

Trapezoidal DVI

$$p_{n+1/2}^1 = \vartheta_1(q_n^1, q_n^2) + \frac{h}{2} D_1 \vartheta_1(q_n^1, q_{n+1/2}^2) \cdot v_{n+1/2}^1 - \frac{h}{2} D_1 H(q_n^1, q_{n+1/2}^2),$$



$$\begin{aligned} \vartheta_1(q_{n+1}^1, q_{n+1}^2) &= \vartheta_1(q_n^1, q_n^2) + \frac{h}{2} \left(D_1 \vartheta_1(q_n^1, q_{n+1/2}^2) + D_1 \vartheta_1(q_{n+1}^1, q_{n+1/2}^2) \right) \cdot v_{n+1/2}^1 \\ &\quad - \frac{h}{2} \left(D_1 H(q_n^1, q_{n+1/2}^2) + D_1 H(q_{n+1}^1, q_{n+1/2}^2) \right), \end{aligned}$$

$$p_{n+1/2}^1 = \frac{1}{2} \left(\vartheta_1(q_n^1, q_{n+1/2}^2) + \vartheta_1(q_{n+1}^1, q_{n+1/2}^2) \right),$$

$$\begin{aligned} v_{n+1/2}^1 &= \left(D_2 \vartheta_1(q_n^1, q_{n+1/2}^2) + D_2 \vartheta_1(q_{n+1}^1, q_{n+1/2}^2) \right)^{-1} \left(D_2 H(q_n^1, q_{n+1/2}^2) \right. \\ &\quad \left. + D_2 H(q_{n+1}^1, q_{n+1/2}^2) \right), \end{aligned}$$

$$q_{n+1}^1 = q_n^1 + h v_{n+1/2}^1$$

Lotka–Volterra System with DG-TDVI



Midpoint DVI

- the approximation of q is given by

$$q_h^1(t)|_{(t_n, t_{n+1})} = \frac{t_{n+1} - t}{t_{n+1} - t_n} q_n^{+,1} + \frac{t - t_n}{t_{n+1} - t_n} q_{n+1}^{-,1},$$

$$q_h^2(t)|_{(t_n, t_{n+1})} = q_{n+1/2}^2$$

- use midpoint quadrature on the Lagrangian

$$\begin{aligned} L_d(q_n, q_{n+1}) = h & \left[\vartheta_1(\bar{q}_{n+1/2}^1, q_{n+1/2}^2) \cdot v_{n+1/2}^1 - H(\bar{q}_{n+1/2}^1, q_{n+1/2}^2) \right] + h p_{n+1/2}^1 \cdot \left[\frac{q_{n+1}^{-,1} - q_n^{+,1}}{h} - v_{n+1/2}^1 \right] \\ & + \vartheta_1(q_n^{+,1}, q_n^2) \cdot (q_n^{+,1} - q_n^1) + \vartheta_1(q_{n+1}^{-,1}, q_{n+1}^2) \cdot (q_{n+1}^1 - q_{n+1}^{-,1}) \end{aligned}$$

with

$$\bar{q}_{n+1/2}^1 = \frac{q_n^{+,1} + q_{n+1}^{-,1}}{2}$$

Midpoint DVI

$$\vartheta_1(\bar{q}_{n+1/2}^1, q_{n+1/2}^2) = \vartheta_1(q_n^1, q_n^2) + \frac{h}{2} D_1 \vartheta_1(\bar{q}_{n+1/2}, q_{n+1/2}^2) \cdot v_{n+1/2}^1 - \frac{h}{2} D_1 H(\bar{q}_{n+1/2}, q_{n+1/2}^2),$$



$$\vartheta_1(q_{n+1}^1, q_{n+1}^2) = \vartheta_1(q_n^1, q_n^2) + h D_1 \vartheta_1(\bar{q}_{n+1/2}, q_{n+1/2}^2) \cdot v_{n+1/2}^1 - h D_1 H(\bar{q}_{n+1/2}, q_{n+1/2}^2),$$

$$v_{n+1/2}^1 = \left(D_2 \vartheta_1(\bar{q}_{n+1/2}^1, q_{n+1/2}^2) \right)^{-1} D_2 H(\bar{q}_{n+1/2}^1, q_{n+1/2}^2),$$

$$q_{n+1}^1 = q_n^1 + h v_{n+1/2}^1,$$

$$\bar{q}_{n+1/2}^1 = \frac{q_n^1 + q_{n+1}^1}{2}$$

Lotka–Volterra System with DG-MDVI



Discontinuous Galerkin DVIs

Symplectic Runge–Kutta Methods

Symplectic Runge–Kutta Methods for Phasespace Lagrangians

$$Q_{n,i} = q_n + h \sum_{j=1}^s a_{ij} V_{n,j},$$

$$P_{n,i} = p_n + h \sum_{j=1}^s a_{ij} F_{n,j},$$

$$q_{n+1}^{\mu} = q_n^{\mu} + h \sum_{i=1}^s b_i V_{n,i}^{\mu},$$

$$p_{n+1}^{\mu} = p_n^{\mu} + h \sum_{i=1}^s b_i F_{n,i}^{\mu},$$

$$p_{n+1}^{\mu} = \vartheta^{\mu}(q_{n+1}),$$

$$P_{n,i} = \frac{\partial L}{\partial \dot{q}}(Q_{n,i}, V_{n,i}),$$

$$F_{n,i} = \frac{\partial L}{\partial q}(Q_{n,i}, V_{n,i}),$$

$$\mu = 1, \dots, d,$$

$$\mu = 1, \dots, d,$$

$$\mu = 1, \dots, 2d$$

Remarks on Projection Methods

Position–Momentum Form

- use discrete fibre derivative to obtain position-momentum form

$$p_n = -D_1 L_d(q_n, q_{n+1}),$$

$$p_{n+1} = D_2 L_d(q_n, q_{n+1})$$

- can be solved as the discrete Lagrangian L_d is not degenerate

→ provides an update rule of the form

$$\tilde{\Psi}_{L_d} : (q_n, p_n) \mapsto (q_{n+1}, p_{n+1})$$

- use continuous fibre derivative to obtain an exact second initial condition p_0

$$p_0 = \frac{\partial L}{\partial \dot{q}}(q_0) = \vartheta(q_0)$$

Position–Momentum Form

- position-momentum form: rewrite the equations of motion as an index-2 DAE

$$\begin{aligned}\dot{z} &= \Omega^{-1}(\nabla H(z) + \nabla \phi^T(z) \lambda), \\ 0 &= \phi(z),\end{aligned}$$

with




$$z = (q, p), \quad \phi(q, p) = p - \vartheta(q), \quad \Omega = \begin{pmatrix} 0 & -\mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}$$

- the variational integrator does not preserve the constraint $\phi(q, p) = 0$



→ the numerical solution drifts away from the constraint submanifold

$$p_n \neq \vartheta(q_n) \quad \text{for } n \geq 1, \text{ even though } p_0 = \vartheta(q_0)$$

Lotka–Volterra System with Variational Integrator



Symmetric Projection on Primary Constraint



$$\tilde{z}_n = z_n + h \Omega^{-1} \nabla \phi^T(z_n) \lambda_{n+1}$$

perturb

$$\tilde{z}_{n+1} = \Psi_h(\tilde{z}_n)$$

apply arbitrary one-step method

$$z_{n+1} = \tilde{z}_{n+1} + h R(\infty) \Omega^{-1} \nabla \phi^T(z_{n+1}) \lambda_{n+1}$$

project on constraint submanifold

$$0 = \phi(z_{n+1})$$

constraint

$(R(\infty) = \pm 1$ is the stability function of Ψ_h)

Symmetric Projection and Symplecticity

- symplecticity condition

$$\begin{aligned} d\mathbf{q}_n^\mu \wedge d\mathbf{p}_n^\mu - d(\lambda_{n+1}^T \phi_{q^\mu}(\mathbf{p}_n, \mathbf{q}_n)) \wedge d(\lambda_{n+1}^T \phi_{p^\mu}(\mathbf{p}_n, \mathbf{q}_n)) = \\ = d\mathbf{q}_{n+1}^\mu \wedge d\mathbf{p}_{n+1}^\mu - |R(\infty)|^2 d(\lambda_{n+1}^T \phi_{q^\mu}(\mathbf{p}_{n+1}, \mathbf{q}_{n+1})) \wedge d(\lambda_{n+1}^T \phi_{p^\mu}(\mathbf{p}_{n+1}, \mathbf{q}_{n+1})) \end{aligned}$$

- for $|R(\infty)| = 1$ a variational integrator with symmetric projection is symplectic iff

$$\begin{aligned} \lambda_{n+1}^T \phi_{p^\mu}(\mathbf{p}_n, \mathbf{q}_n) &= \lambda_{n+1}^T \phi_{p^\mu}(\mathbf{p}_{n+1}, \mathbf{q}_{n+1}) && \text{for all } \mu, \\ \lambda_{n+1}^T \phi_{q^\mu}(\mathbf{p}_n, \mathbf{q}_n) &= \lambda_{n+1}^T \phi_{q^\mu}(\mathbf{p}_{n+1}, \mathbf{q}_{n+1}) && \text{for all } \mu \end{aligned}$$

- the first condition is always satisfied
- for special phasespace Lagrangians, λ_{n+1}^1 vanishes to machine precision while ϑ^2 is zero by assumption so that also the second condition is satisfied

Summary and Outlook

Summary and Outlook

- Degenerate Variational Integrators (DVIs)
 - one-step methods for degenerate Lagrangians obtained directly from a discrete action
 - original work (C. L. Ellison): 1st order, not composable
 - leapfrog methods (J. W. Burby): 2nd order, composable, require processing for initialisation
 - preservation of a discrete symplectic structure
- Discontinuous Galerkin Degenerate Variational Integrators (DG-DVIs)
 - Galerkin- and Runge–Kutta methods with arbitrary order
 - recover 1st-order DVIs, symplectic Runge–Kutta methods, and some projection methods
 - preservation of the continuous symplectic structure
- Open Problems
 - generalisation to arbitrary degenerate Lagrangians
 - closed-form expression for degenerate discrete Lagrangians

References

■ References

- C. L. Ellison et al. Degenerate variational integrators for magnetic field line flow and guiding center trajectories, Physics of Plasmas 25, 052502, 2018
- C. L. Ellison. Development of Multistep and Degenerate Variational Integrators for Applications in Plasma Physics, Doctoral Thesis, Princeton University, 2016
- J. W. Burby et al. Improved accuracy in degenerate variational integrators for guiding center and magnetic field line flow, 2021
- MK. Projected Variational Integrators for Degenerate Lagrangian Systems, arXiv:1708.07356
- MK. Symplectic Runge–Kutta Methods for Degenerate Lagrangian Systems, in preparation
- MK. Discontinuous Galerkin Variational Integrators for Degenerate Lagrangians, in preparation
- MK. On Action Principles and Degenerate Lagrangians: Continuous and Discrete, in preparation

■ Implementation

- <https://github.com/JuliaGNI/GeometricIntegrators.jl>
- <https://github.com/JuliaGNI/GeometricProblems.jl>
- <https://github.com/JuliaPlasma/ChargedParticleDynamics.jl>
- <https://github.com/JuliaPlasma/ElectroMagneticFields.jl>