

What can 1 week of seismic data reveal
about Strokkur's eruption rhythm?

Seismic Analysis of the Strokkur Geyser

Marwan Saad, Master Student


Under Supervision of Prof. Dr. Eva Eibl

Seismological Data Science with Application to Volcano Systems

Strokkur Geyser

- **Location:** in the Haukadalur geothermal area, southwest Iceland.
- **Eruption:** repeatedly every few minutes, Heated groundwater, steam, and gas build pressure in the conduit.
- **Seismic production:** Bubble collapse and rapid fluid movement generate seismic vibrations.
- **Seismic Monitor usefulness:** enables continuous, nonvisual eruption detection and recurrence analysis.

Workflow



Initial Data Inspection

- Sampling rate: **100 Hz**.
- **7 days** of Seismic recordings.
- **63** Traces.

Station–Component Diagram




Figure created with generative AI for illustrative purposes.

preprocessing

- Demean
- Detrend
- Apply Hann Taper
- Apply 10-40 Hz band-pass Filter

-Preprocessing reduced drift, edge effects, and low-frequency noise.

Filter Selection

Why 10-40 Hz?

- Inspected **waveform, spectrum, & spectrogram.**
- Useful event energy mainly between **10-40 Hz.**
- Filter preserved the transient & **reduced background noise.**

Triggering Algorithms in ObsPy

- Applied to **continuous data**.
- Detected **sudden** seismic transients.
- Compared across **3** stations.
- **Retained** coherent network events.

Grouping detected events

```
[56]: catalog_week = pd.DataFrame(all_week_events)
```

```
catalog_week["time"] = pd.to_datetime(
    catalog_week["time"].astype(str)
)
```

```
catalog_week
```

```
[56]:
```

	day	time	duration	stations
0	1	2026-01-01 00:01:09.890000+00:00	3.74	[GE4, GE5, GE2]
1	1	2026-01-01 00:05:59.260000+00:00	3.55	[GE4, GE2, GE5]
2	1	2026-01-01 00:09:29.900000+00:00	5.07	[GE2, GE4, GE5]
3	1	2026-01-01 00:13:59.210000+00:00	3.58	[GE2, GE4, GE5]
4	1	2026-01-01 00:18:30.450000+00:00	3.67	[GE4, GE2, GE5]
...	...	...	...	...
3336	7	2026-01-07 10:59:54.910000+00:00	4.05	[GE5, GE2, GE4]
3337	7	2026-01-07 11:01:43.260000+00:00	6.10	[GE5, GE4, GE2]
3338	7	2026-01-07 11:05:25.070000+00:00	3.73	[GE4, GE2, GE5]
3339	7	2026-01-07 11:05:45.810000+00:00	3.24	[GE4, GE2, GE5]
3340	7	2026-01-07 11:05:55.470000+00:00	2.73	[GE4, GE2, GE5]

3341 rows × 4 columns

- **Closely spaced** detections combined.
- Grouping window: **30 S**.
- Compared across **3** stations.
- **Repeated** triggers treated as **1 episode**.
- **3341** transients—**2904** episodes.

Screenshot from the Jupyter Notebook analysis.


Event cataloging

- **Grouped** episodes within **2 m**.
- **Compared** network peak **amplitudes**.
- Kept the **strongest** event per cluster.
- **2904** episodes – **1946** principal candidates.

Recurrence analysis

- Stable rate: **12-13** candidates/hr.
- **Mean** Interval: **4.78** min.
- **Median** interval: **4.27** min.
- **Most** interval: **3-6** min.

-Recurrence seismic activity remained relatively regular throughout the week




Screenshot from the Jupyter Notebook analysis.

Rep., event selection


- Selected from **final** candidate **catalog**.
- Clear at all **3** stations.
- **Strong** signal to noise **ratio**.
- **Visible** on HHE, HHN, HHZ.

-Selected event: 2 January 2026 – 09:50:23.67 UTC


2026-01-02T09:50:13.67 - 2026-01-02T09:50:53.67



Screenshot from the Jupyter Notebook analysis.




Waveform & Spectrogram



Screenshot from the Jupyter Notebook analysis.

Particle motion analysis



Interpretation

The data show regular, persistent seismic activity, with most candidates repeating every 3–6 minutes and the selected event showing a clear waveform, frequency signal, and directional motion.

Limitations

- Candidates were not **visually** confirmed eruptions.
- **No** station **coordinates** were provided.
- **Back azimuth** retained a **180°** ambiguity.
- The **30 s** and **2 min** thresholds were project-specific choices.
- **Day 7** contained only about **12 hours** of data.

Conclusions

- **Continuous** seismic monitoring successfully detected recurring Strokkur-related events **throughout the week**. The **principal candidates** occurred mainly every **3–6 minutes**, while **waveform**, **spectrogram**, and **particle-motion** analysis clearly characterized **one** representative **event**. The results show that seismic data can effectively monitor regular geyser activity, although independent **visual confirmation** and station **coordinates** would **improve the interpretation**.

References

- Eibl, E. P. S., et al. (2020). *Eruption interval monitoring at Strokkur Geyser, Iceland*. **Geophysical Research Letters**, **47**, e2019GL085266.
- Eibl, E. P. S., et al. (2024). *Video camera and seismic monitoring of water-bulge explosions at Strokkur Geyser, Iceland*. **Volcanica**, **7**(1).
- GFZ German Research Centre for Geosciences. (2020). *Seismological experiment at Strokkur from 2020—Network 2Z*.
- ObsPy Development Team. *ObsPy documentation: Python framework for processing seismological data*.

Thank you!

Do not be hesitant to ask any **Questions!**