

Clinical analysis report


Patient & case information

Doctor	sama
Patient	jana
Patient code	PT 45
Age	20 years
Gender	Female
Case number	#80
Case date	25 May 2026
Created	25 May 2026 · 09:23


Side view — measurement summary

Measurement	Value	Classification	Normal range	Clinical note
Nasiolabial angle	134.3°	Increased (>110°)	90° – 110°	Possible high columella or retruded maxilla. Class III cases may require surgical evaluation depending on full diagnosis.
Profile convexity angle	158.2°	Normal (151°–171°)	151° – 171°	Within normal range. No treatment implication from this angle alone.
Total facial convexity angle	132.7°	Normal (127°–137°)	127° – 137°	Within normal range. No treatment implication from this angle alone.
Mentolabial angle	123.7°	Normal (110°–130°)	110° – 130°	Within normal range. No treatment implication from this angle alone.

Side view — measurement visualizations




Nasiolabial angle
134.3°
Normal: 90° – 110°
Increased (>110°)
Possible high columella or retruded maxilla.
Class III cases may require surgical evaluation depending on full diagnosis.



Profile convexity angle
158.2°
Normal: 151° – 171°
Normal (151°–171°)
Within normal range.
No treatment implication from this angle alone.

Side view — measurement visualizations (continued)


**Total facial convexity angle**
132.7°

Normal: 127° – 137°

Normal (127°–137°)

Within normal range.

No treatment implication from this angle alone.

**Mentolabial angle**
123.7°

Normal: 110° – 130°

Normal (110°–130°)

Within normal range.


No treatment implication from this angle alone.

Frontal non-smile analysis

Frontal non-smile — measurement summary

Measurement	Value	Interpretation
Middle vs lower facial third (proxy)	Middle 382.7 px, lower 383.8 px, ratio 0.997	Approximate vertical proportions; interpret with clinical context and growth pattern.
Upper / lower lip height ratio	0.143	Values far from ~1.0 may suggest disproportion; correlate with dental and soft-tissue exam.
Interpupillary vs commissure line	Interpupillary 0.0°, commissure 0.0°	Parallel within 5.0° (lines appear level).
Facial midline (RMS deviation)	126.2 px	Midline deviation index is elevated; assess chin, nose, and dental midlines clinically.
Rule of fifths (approximate)	Ideal fifth width 154.2 px	Compare eye widths and intercanthal distance to one-fifth of bizygomatic width; rough aesthetic guide only.
Facial width (bizygomatic)	771.0 px	Useful for proportional checks; correlate with arch form and transverse skeletal pattern.
Mandibular width (bigonial)	771.0 px	Wide or narrow lower face relative to cheekbones informs transverse diagnosis.
Bizygomatic / bigonial ratio	1.000	Typical values often near ~1.2–1.4 in many samples; extremes suggest disproportion worth clinical review.
Facial index (height / width)	0.735	Longer narrower vs shorter wider facial form; integrate with vertical skeletal and growth assessment.
Interlabial gap (approx.)	84.0 px	Approximate soft-tissue opening at rest; verify with clinical lip posture exam.

Frontal visualization & interpretations



Annotated frontal overlay

Middle vs lower facial third (proxy)

Middle 382.7 px, lower 383.8 px, ratio 0.997

Approximate vertical proportions; interpret with clinical context and growth pattern.

Upper / lower lip height ratio

0.143

Values far from ~1.0 may suggest disproportion; correlate with dental and soft-tissue exam.

Interpupillary vs commissure line

Interpupillary 0.0°, commissure 0.0°

Parallel within 5.0° (lines appear level).

Facial midline (RMS deviation)

126.2 px

Midline deviation index is elevated; assess chin, nose, and dental midlines clinically.

Rule of fifths (approximate)

Ideal fifth width 154.2 px

Compare eye widths and intercanthal distance to one-fifth of bizygomatic width; rough aesthetic guide only.

Facial width (bizygomatic)

771.0 px

Useful for proportional checks; correlate with arch form and transverse skeletal pattern.

Mandibular width (bigonial)

771.0 px

Wide or narrow lower face relative to cheekbones informs transverse diagnosis.

Bizygomatic / bigonial ratio

1.000

Typical values often near ~1.2–1.4 in many samples; extremes suggest disproportion worth clinical review.

Facial index (height / width)

0.735

Longer narrower vs shorter wider facial form; integrate with vertical skeletal and growth assessment.

Interlabial gap (approx.)

84.0 px

Approximate soft-tissue opening at rest; verify with clinical lip posture exam.