

Clinical analysis report


Patient & case information

Doctor	younna
Patient	jana
Patient code	PT 1
Age	21 years
Gender	Female
Case number	#83
Case date	25 May 2026
Created	25 May 2026 · 11:44

Side view — measurement summary

Measurement	Value	Classification	Normal range	Clinical note
Nasiolabial angle	94.2°	Normal (90°–110°)	90° – 110°	Within normal range. No treatment implication from this angle alone.
Profile convexity angle	164.7°	Normal (151°–171°)	151° – 171°	Within normal range. No treatment implication from this angle alone.
Total facial convexity angle	115.9°	Decreased (<127°)	127° – 137°	Possible retruded chin or prominent nasal effect. May suggest Class II tendency; requires full orthodontic evaluation.
Mentolabial angle	150.4°	Increased (>130°)	110° – 130°	Possible retruded chin, proclined lower teeth, or advanced mandible pattern. May require orthodontic correction, genioplasty, or surgery depending on full diagnosis.

Side view — measurement visualizations




Nasiolabial angle 94.2°

Normal: 90° – 110°

Normal (90°–110°)

Within normal range.

No treatment implication from this angle alone.



Profile convexity angle 164.7°


Normal: 151° – 171°

Normal (151°–171°)

Within normal range.

No treatment implication from this angle alone.

Side view — measurement visualizations (continued)


**Total facial convexity angle**
115.9°

Normal: 127° – 137°

Decreased (<127°)

Possible retruded chin or prominent nasal effect.

May suggest Class II tendency; requires full orthodontic evaluation.

**Mentolabial angle**
150.4°

Normal: 110° – 130°

Increased (>130°)

Possible retruded chin, proclined lower teeth, or advanced mandible pattern.


May require orthodontic correction, genioplasty, or surgery depending on full diagnosis.

Frontal non-smile analysis

Frontal non-smile — measurement summary

Measurement	Value	Interpretation
Middle vs lower facial third (proxy)	Middle 230.9 px, lower 200.0 px, ratio 1.154	Approximate vertical proportions; interpret with clinical context and growth pattern.
Upper / lower lip height ratio	0.250	Values far from ~1.0 may suggest disproportion; correlate with dental and soft-tissue exam.
Interpupillary vs commissure line	Interpupillary 0.0°, commissure -9.4°	Difference 9.4° exceeds typical cosmetic parallelism threshold (5.0°); clinical correlation suggested.
Facial midline (RMS deviation)	8.7 px	Midline landmarks cluster closely; gross asymmetry unlikely from this proxy.
Rule of fifths (approximate)	Ideal fifth width 123.0 px	Compare eye widths and intercanthal distance to one-fifth of bizygomatic width; rough aesthetic guide only.
Facial width (bizygomatic)	615.0 px	Useful for proportional checks; correlate with arch form and transverse skeletal pattern.
Mandibular width (bigonial)	460.0 px	Wide or narrow lower face relative to cheekbones informs transverse diagnosis.
Bizygomatic / bigonial ratio	1.337	Typical values often near ~1.2–1.4 in many samples; extremes suggest disproportion worth clinical review.
Facial index (height / width)	0.700	Longer narrower vs shorter wider facial form; integrate with vertical skeletal and growth assessment.
Interlabial gap (approx.)	40.0 px	Approximate soft-tissue opening at rest; verify with clinical lip posture exam.

Frontal visualization & interpretations



Annotated frontal overlay

Middle vs lower facial third (proxy)

Middle 230.9 px, lower 200.0 px, ratio 1.154

Approximate vertical proportions; interpret with clinical context and growth pattern.

Upper / lower lip height ratio

0.250

Values far from ~1.0 may suggest disproportion; correlate with dental and soft-tissue exam.

Interpupillary vs commissure line

Interpupillary 0.0°, commissure -9.4°

Difference 9.4° exceeds typical cosmetic parallelism threshold (5.0°); clinical correlation suggested.

Facial midline (RMS deviation)

8.7 px

Midline landmarks cluster closely; gross asymmetry unlikely from this proxy.

Rule of fifths (approximate)

Ideal fifth width 123.0 px

Compare eye widths and intercanthal distance to one-fifth of bizygomatic width; rough aesthetic guide only.

Facial width (bizygomatic)

615.0 px

Useful for proportional checks; correlate with arch form and transverse skeletal pattern.

Mandibular width (bigonial)

460.0 px

Wide or narrow lower face relative to cheekbones informs transverse diagnosis.

Bizygomatic / bigonial ratio

1.337

Typical values often near ~1.2–1.4 in many samples; extremes suggest disproportion worth clinical review.

Facial index (height / width)

0.700

Longer narrower vs shorter wider facial form; integrate with vertical skeletal and growth assessment.

Interlabial gap (approx.)

40.0 px

Approximate soft-tissue opening at rest; verify with clinical lip posture exam.