

Clinical analysis report

Patient & case information

Doctor	yassin
Patient	zee tome
Patient code	12345
Age	18 years
Gender	Male
Case number	#184
Case date	19 Jun 2026
Created	19 Jun 2026 · 01:12

Medical History

Conditions:

• High blood pressure • Currently taking regular medications • Previous major surgeries

Previous Surgeries:

tttt

Additional Notes:


gfgkvgtg

Medical history reflects the information currently stored in the patient profile at the time this report was generated.

Side view — measurement summary

Measurement	Value	Classification	Normal range	Clinical note
Nasiolabial angle	149.2°	Increased (>110°)	90° – 110°	Possible high columella or retruded maxilla. Class III cases may require surgical evaluation depending on full diagnosis.
Profile convexity angle	164.3°	Normal (151°–171°)	151° – 171°	Within normal range. No treatment implication from this angle alone.
Total facial convexity angle	136.6°	Normal (127°–137°)	127° – 137°	Within normal range. No treatment implication from this angle alone.
Mentolabial angle	141.1°	Increased (>130°)	110° – 130°	Possible retruded chin, proclined lower teeth, or advanced mandible pattern. May require orthodontic correction, genioplasty, or surgery depending on full diagnosis.

Side view — measurement visualizations




Nasiolabial angle
149.2°

Normal: 90° – 110°

Increased (>110°)

Possible high columella or retruded maxilla.

Class III cases may require surgical evaluation depending on full diagnosis.




Profile convexity angle
164.3°

Normal: 151° – 171°

Normal (151°–171°)

Within normal range.

No treatment implication from this angle alone.




Total facial convexity angle
136.6°

Normal: 127° – 137°

Normal (127°–137°)

Within normal range.

No treatment implication from this angle alone.



Mentolabial angle
141.1°

Normal: 110° – 130°

Increased (>130°)

Possible retruded chin, proclined lower teeth, or advanced mandible pattern.

May require orthodontic correction, genioplasty, or surgery depending on full diagnosis.


Frontal non-smile analysis

Frontal non-smile — measurement summary

Frontal non-smile — measurement summary (continued)

Measurement	Value	Interpretation
Middle vs lower facial third (proxy)	Middle 401.2 px, lower 337.0 px, ratio 1.191	Approximate vertical proportions; interpret with clinical context and growth pattern.
Upper / lower lip height ratio	0.347	Values far from ~1.0 may suggest disproportion; correlate with dental and soft-tissue exam.
Interpupillary vs commissure line	Interpupillary 0.0°, commissure 91.7°	Difference 91.7° exceeds typical cosmetic parallelism threshold (5.0°); clinical correlation suggested.
Facial midline (RMS deviation)	101.4 px	Midline deviation index is elevated; assess chin, nose, and dental midlines clinically.
Rule of fifths (approximate)	Ideal fifth width 125.6 px	Compare eye widths and intercanthal distance to one-fifth of bizygomatic width; rough aesthetic guide only.
Facial width (bizygomatic)	628.1 px	Useful for proportional checks; correlate with arch form and transverse skeletal pattern.
Mandibular width (bigonial)	774.0 px	Wide or narrow lower face relative to cheekbones informs transverse diagnosis.
Bizygomatic / bigonial ratio	0.811	Typical values often near ~1.2–1.4 in many samples; extremes suggest disproportion worth clinical review.
Facial index (height / width)	1.070	Longer narrower vs shorter wider facial form; integrate with vertical skeletal and growth assessment.
Interlabial gap (approx.)	91.7 px	Approximate soft-tissue opening at rest; verify with clinical lip posture exam.

Frontal visualization & interpretations



Annotated frontal overlay

Middle vs lower facial third (proxy)

Middle 401.2 px, lower 337.0 px, ratio 1.191

Approximate vertical proportions; interpret with clinical context and growth pattern.

Upper / lower lip height ratio

0.347

Values far from ~1.0 may suggest disproportion; correlate with dental and soft-tissue exam.

Interpupillary vs commissure line

Interpupillary 0.0°, commissure 91.7°

Difference 91.7° exceeds typical cosmetic parallelism threshold (5.0°); clinical correlation suggested.

Facial midline (RMS deviation)

101.4 px

Midline deviation index is elevated; assess chin, nose, and dental midlines clinically.

Rule of fifths (approximate)

Ideal fifth width 125.6 px

Compare eye widths and intercanthal distance to one-fifth of bizygomatic width; rough aesthetic guide only.

Facial width (bizygomatic)

628.1 px

Useful for proportional checks; correlate with arch form and transverse skeletal pattern.

Mandibular width (bigonial)

774.0 px

Wide or narrow lower face relative to cheekbones informs transverse diagnosis.

Bizygomatic / bigonial ratio

0.811

Typical values often near ~1.2–1.4 in many samples; extremes suggest disproportion worth clinical review.

Frontal interpretations (continued)

Facial index (height / width)

1.070

Longer narrower vs shorter wider facial form; integrate with vertical skeletal and growth assessment.

Interlabial gap (approx.)

91.7 px

Approximate soft-tissue opening at rest; verify with clinical lip posture exam.