


CL=CSCI 160

CLASS 9

## Gates




OR




Inverter




AND



NOR



NAND



XOR  
Exclusive OR

## Example - logic diagram

Remember:

AND:


•	0	1
0	0	0
1	0	1

OR:

+	0	1
0	0	1
1	1	1

Problem:

Find the truth table and expression of the function F.



Solution

I recommend finding the expression of F first, and then deduce the truth table from the expression.

We have:  $(xy)' + z = x' + y' + z$

$(xz)' + y = x' + z' + y$

$(yz)' + x = y' + z' + x$

What is then F?

$$F = (x' + y' + z)(x' + z' + y)(y' + z' + x)$$

What can we say about this product?

It is mostly = 0, like every product.

It is = 0 when any one of its factors = 0

$x' + y' + z = 0 \leftrightarrow x'=0 \text{ and } y'=0 \text{ and } z=0$   
 $x=1 \text{ and } y=1 \text{ and } z=0$

Similarly we have:

$x' + y + z' = 0 \leftrightarrow x=1 \text{ and } y=0 \text{ and } z=1$

$x + y' + z' = 0 \leftrightarrow x=0 \text{ and } y=1 \text{ and } z=1$

In rest, for all other values of x, y, z, F = 1

x	y	z	F
0	0	0	1
0	0	1	1
0	1	0	1
0	1	1	0
1	0	0	1
1	0	1	0
1	1	0	0
1	1	1	1

Truth table

## HW 8.1

### Partial Solution + Hint

#### (a) Prove: Inhibition is not commutative

Table gives two functions:

$F_2 = xy'$	$x/y$	Inhibition	$x$ , but not $y$
$F_4 = x'y$	$y/x$	Inhibition	$y$ , but not $x$

Let's choose  $F_2 = x/y = x y'$

$$\begin{array}{l} \text{F}_2 \text{ commutative} \stackrel{\text{Det}}{\Leftrightarrow} \boxed{F_2(x, y) = F_2(y, x) \quad \forall x, y} \stackrel{\text{Det}}{\Leftrightarrow} \boxed{x/y = y/x \quad \forall x, y} \\ \text{F}_2 \text{ not commutative} \stackrel{\text{Det}}{\Leftrightarrow} \exists x, y: F_2(x, y) \neq F_2(y, x) \stackrel{\text{Det}}{\Leftrightarrow} \exists x, y: x/y \neq y/x \end{array}$$

In other words we need to find a counterexample to  $x/y = y/x$ , that is find values in  $\{0, 1\}$  for  $x, y$ , such that we have  $x/y \neq y/x \stackrel{\text{Det}}{\Leftrightarrow} xy' \neq yx'$  as every B.A. has 0 and 1 as elements!

Make one side 1 and at the same time the other side 0.

$x = 1$  and  $y = 0$  imply:

$$LS = 1 \cdot 1 = 1 \quad RS = 0 \cdot 0 = 0 \quad \rightarrow LS \neq RS \quad \rightarrow \text{F}_2 \text{ is not commutative}$$

Done

## HW 8.1 - continued      Partial Solution + Hint - continued

(b) Prove: Exclusive OR (XOR) is associative      Table:

$F_6 = xy' + x'y$        $x \oplus y$       Exclusive-OR       $x$  or  $y$ , but not both

$$F_6(x, y) = x y' + x' y = x \oplus y$$

$$F_6 \text{ associative} \stackrel{\text{Det}}{\Leftrightarrow} (x \oplus y) \oplus z = x \oplus (y \oplus z) \quad \forall x, y, z$$

$$\stackrel{\text{Det}}{\Leftrightarrow} F_6(F_6(x, y), z) = F_6(x, F_6(y, z)) \quad \forall x, y, z$$

We need to prove this for all BAs, so we need to use the axioms and properties of a BA.

$$LS = (x y' + x' y) \oplus z = (x y' + x' y) z' + (x y' + x' y)' z =$$

$$= xy'z' + x'yz' + (x' + y)(x + y')z = xy'z' + x'yz' + x'y'z + xyz$$

$$RS = x \oplus (yz' + y'z) = x(yz' + y'z)' + x'(yz' + y'z) = x(y' + z)(y + z') + x'yz' + x'y'z =$$

$$= xy'z' + xyz + x'yz' + x'y'z \quad \rightarrow \quad LS = RS \quad \rightarrow \quad \underline{F_6 \text{ is associative}}$$

Done

### HW 7.3 (continuation from class 8)

2-9 b)

$$\overline{((\bar{v}w + x)y + \bar{z})} = \underbrace{((v + \bar{w}) \cdot \bar{x} + \bar{y})}_{{v\bar{x} + \bar{w}\bar{x}}} z =$$

$$= (v\bar{x} + \bar{w}\bar{x} + \bar{y})z = \underline{v\bar{x}z + \bar{w}\bar{x}z + \bar{y}z}$$

c)  $\overline{(w x (\bar{y} z + y \bar{z}) + \bar{w} \bar{x} (\bar{y} + z)(y + \bar{z}))} =$

$$= (\bar{w} + \bar{x} + (y + \bar{z})(\bar{y} + z))(w + x + y\bar{z} + \bar{y}z) =$$

$$= (\bar{w} + \bar{x} + yz + \bar{y}\bar{z})(w + x + y\bar{z} + \bar{y}z) =$$

$$= \bar{w}x + \bar{w}y\bar{z} + \bar{w}\bar{y}z + \bar{x}w + \bar{x}y\bar{z} + \bar{x}\bar{y}z + wyz + xyz + \\ + w\bar{y}\bar{z} + x\bar{y}\bar{z}$$

### HW 7.3 (continuation)

2-9 d)

$$\begin{aligned} & \overbrace{((A + \bar{B} + c)(\bar{A}\bar{B} + c)(A + \bar{B}\bar{c}))} = \\ &= \bar{A}B\bar{c} + (A+B)\cdot\bar{c} + \bar{A}\cdot(B+c) \quad (4) \\ &= \bar{A}B\bar{c} + A\bar{c} + B\bar{c} + \bar{A}B + \bar{A}c \quad (12) \\ &= A\bar{c} + B\bar{c} + \bar{A}B + \bar{A}c \end{aligned}$$

What now? Expand and then contract again:

$$\begin{aligned} &= \cancel{AB\bar{c}} + A\bar{B}\bar{c} + \cancel{AB\bar{c}} + \cancel{\bar{A}B\bar{c}} + \cancel{\bar{A}B\bar{c}} + \bar{A}B\bar{c} + \\ &+ \bar{A}Bc + \bar{A}\bar{B}c = \underline{A\bar{B}\bar{c}} + \underline{AB\bar{c}} + \underline{\bar{A}B\bar{c}} + \\ &+ \underline{\bar{A}Bc} + \underline{\bar{A}\bar{B}c} = \underline{A\bar{c}} + \underline{\bar{A}B} + \underline{\bar{A}c} \end{aligned}$$