

# The symptomatology of fever: A step towards qualitative definition of fever.

## Article Metadata

**Authors:** Kapoor M, Kumar N, Panda PK

**Journal:** J Family Med Prim Care

**Year:** 2022

**Volume/Issue:** 11/10

**PMID:** 36618218

**DOI:** 10.4103/jfmprc.jfmprc\_360\_22

**PubMed URL:** <https://ncbi.nlm.nih.gov/pubmed/36618218>

## Full Text (from PubMed Central)

<https://pmc.ncbi.nlm.nih.gov/api/oai/v1/mh/>

oai:pubmedcentral.nih.gov:9810860

J Family Med Prim Care

J Family Med Prim Care

J Family Med Prim Care

Journal of Family Medicine and Primary Care

Wolters Kluwer -- Medknow Publications

10.4103/jfmprc.jfmprc\_360\_22

The symptomatology of fever: A step towards qualitative definition of fever

Department of Internal Medicine (Infectious Disease Division), All India Institute of Medical Sciences, Rishikesh, Uttarakhand, India

Address for correspondence:

motherprasanna@rediffmail.com

Copyright: © 2022 Journal of Family Medicine and Primary Care

<https://creativecommons.org/licenses/by-nc-sa/4.0/>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

The old definitions of fever are based on cross-sectional surveys of the population without analyzing the associated symptomatology as fever is a sign, not symptom. Therefore, a longitudinal follow-up study is the need of hour to analyze associated symptoms with fever.

In a longitudinal study over one year, 196 participants recorded three temperature readings daily, one after waking up, one between 12 and 3 PM, one before sleeping, and filled the symptomatology questionnaire in a thermometry diary.

Per protocol analysis was done for febrile participants (n = 144). Fatigue (50.3%), warmth (47.3%), headache/head heaviness (47.0%), feeling malaise/general weakness (46.7%), loss of appetite (46.5%), muscle cramps/muscle aches (45.6%), chills/shivering (44.6%), increased sweating (43.0%), nausea (42.5%), irritability (38.9%), increased breathing rate (37.1%), and restlessness/anxiety/palpitations (36.5%) were the symptoms maximally seen during the febrile phase. A higher number of associated symptoms are associated with higher temperature readings. Dehydration suggested the numerically highest temperature values ( $100.86 \pm 1.70^{\circ}\text{F}$ ) but seen in few febrile patients.

Incorporation of symptom analysis in febrile patients is the need of the hour. Fatigue and warmth are found to be the most prevalent symptoms during febrile phase. Associated symptoms can help in predicting the intensity of fever also.

temperature recording

pmc-prop-legally-suppressed

pmc-prop-has-supplement

pmc-prop-suppress-copyright

pmc-prop-is-real-version

Fever has long been regarded as an important vital sign. Humans maintain their standard body temperature in a narrow range for the smooth functioning of the body.[

The ideal way to define fever is to follow-up healthy subjects prospectively. Moreover, when such individuals develop rise in baseline temperature with associated symptoms, fever is set to begin.

Thereby, in the study, we recruited healthy participants and longitudinally followed them up and analyzed their temperature in association with their symptoms that were present at that time. We analyzed which symptomatology suggested fever and whether there was any relationship between number of symptoms and the absolute temperature values.

A longitudinal study was conducted in a tertiary care hospital from July 2019 to September 2020. A sample size of 192 was calculated based on a prior study done by Mackowiak

The study was conducted in three phases: Non-febrile, febrile, and post-febrile phase. The participants recorded their temperature readings in all three phases. Minimum three oral temperature readings were recorded daily, one after waking up (AM), one between 12 and 3 PM (AN), and one before sleeping (PM). Participants filled a questionnaire related to the symptomatology of fever along with every temperature recording in provided clinical thermometry diary [

#### Supplemental Material 1

The febrile phase data were analyzed as per-protocol analysis (for those participants only who developed the febrile phase during the study) using Statistical Package for Social Sciences (SPSS) Version 23. The frequency distribution of the various symptoms associated with fever was analyzed. The association of absolute values of temperature with the various symptoms was also done. Taking confidence interval as 95%, a

The study screened 350 participants, and 144 were included in the analysis. The mean age of the participants was  $24.24 \pm 5.92$  years (8-58 years). 26% (36) were <20 years, 72.1% (104) were in the age group 20-40 years, and 1.9% (4) were >40 years of age. 52.1% (75) were males. Various symptoms were present during the febrile phase, arranged in descending order [

#### Symptomatology of three temperature phases among participants (

(a and b): Box and Whisker plots: Association between febrile symptoms and temperature readings

We found that fatigue is the most common symptom associated with the febrile phase (50.3%), followed by warmth (47.3%). The febrile phase was associated with at least one symptom in 98.5% of the readings, whereas  $\geq$  two were seen in 95.8%. The number of symptoms seen in various phases was:  $0.01 \pm 0.08$  (non-febrile),  $6.08 \pm 2.42$  (febrile), and  $0.07 \pm 0.36$  (post-febrile) ( $P < 0.001$ ). One symptom was confirmly present when mean temperature was  $99.91 \pm 1.55^{\circ}\text{F}$ , while mean temperature for two associated symptoms was  $100.22 \pm 1.46^{\circ}\text{F}$ . As per our knowledge, no other study has analyzed

the symptoms associated with fever.

As per classical teaching, fever is defined as an early morning temperature of  $>37.2^{\circ}\text{C}$  ( $>98.9^{\circ}\text{F}$ ) or an evening temperature of  $>37.7^{\circ}\text{C}$  ( $>99.9^{\circ}\text{F}$ ).[

It is essential to derive a symptom-based definition of fever, where patients can be taught about a specific set of symptoms to look out for, suspecting fever. In the present scenario of COVID-19, it gathers more importance as due to the shortage of amenities, even thermometers cannot be ensured in every nook and corner. Hence, the patients can lookout for a few symptoms which may most likely predict fever. The present study confirms fatigue is most common associated symptom (50.3%) during febrile phase followed by warmth (47.3%), headache/head heaviness (47.0%), feeling malaise/generalized weakness (46.7%), loss of appetite (46.5%), muscle cramps/muscle aches (45.6%), chills/shivering (44.6%), increased sweating (43.0%), nausea (42.5%), irritability (38.9%), increased breathing rate (37.1%), and restlessness/anxiety/palpitations (36.5%).

The presence of  $\geq$  two symptoms was associated with higher temperature readings ( $100.22^{\circ}\text{F}$ ) compared to just one symptom ( $99.91^{\circ}\text{F}$ ). Similarly, dehydration correlated with higher temperature ( $100.86^{\circ}\text{F}$ ) followed by increase in breathing rate ( $100.61 \pm 1.38^{\circ}\text{F}$ ). These findings can help in the early prediction of patients having a higher temperature and more likely to get sick. This may help in triaging the patients in the presently overburdened scenario of the COVID-19 pandemic also.

Our study has limitations too. Subjects were allowed to take antipyretics; hence, the symptoms could have resolved post drug use. The categorization among phases (febrile and non-febrile) was based on subjective parameters, varying from person to person. However, this itself forms the basis of our study as we propose that fever must be defined based on an individual's assessment of the symptoms. Future studies may establish objective measurements of these associated symptoms.

Fever must not be defined as a mere rise of temperature, as this may occur physiologically also. As fever is a sign, various symptomatology of fever should be defined. Fatigue is found to be the most prevalent symptom (50.3%) during febrile phase followed by warmth, headache, malaise, loss of appetite, muscle aches, chills, increased sweating, nausea, irritability, increased breathing rate, and palpitations. Simultaneously higher rise in temperature is associated with more number of symptoms, thus predicting the intensity of fever.

Fever is a sign, not a symptom.

Symptomatology of fever should be defined.

Fatigue is the most prevalent symptom during febrile phase.

Greater rise in temperature correlates with more symptoms, thereby predicting the intensity of fever.

MK contributed to the data analysis and was involved in manuscript writing. NK contributed to the data collection and reviewing the manuscript. PKP gave the concept, interpreted analysis, critically reviewed the draft, and approved it for publication along with all authors.

## Ethics and data sharing

The study was done after institute ethical approval and as per declaration of Helsinki. After obtaining approval from corresponding author, de-identified data can be shared.

## Financial support and sponsorship

## Conflicts of interest

There are no conflicts of interest.

## Supplemental Material 1

Thanks all participants for recording daily temperature in thermometry diary and Mr Gopal for data preparation from diary to excel. Thanks to Dr Ajeet, Dr Minakshi, and Dr Yogesh for helping in protocol development.

## Ganong's Review of Medical Physiology, 26e

A critical appraisal of 98.6°F, the upper limit of the normal body temperature, and other legacies of Carl Reinhold August Wunderlich

## Physiology, Ovulation, Basal Body Temperature

## StatPearls Publishing

## Harrison's Principles of Internal Medicine, 20e

## The pathophysiological basis and consequences of fever

10.1186/s13054-016-1375-5

## Normal Body Temperatures in N. India

10.1136/bmj.1.4458.909

## Decreasing human body temperature in the United States since the industrial revolution

## Range for normal body temperature in the general population of Pakistan

## Comparison of body temperature of elderly with adult in South India

## Mpl J Nurs Health Sci

Is older colder or colder older? The association of age with body temperature in 18,630 individuals

J Gerontol A Biol Sci Med Sci

10.1093/gerona/glr001

Tropical fevers in Indian intensive care units:A prospective multicenter study

Indian J Crit Care Med

10.4103/ijccm.IJCCM\_324\_17

Tropical fevers:Management guidelines

Indian J Crit Care Med

10.4103/0972-5229.126074

## Citation

Kapoor M, Kumar N, Panda PK. The symptomatology of fever: A step towards qualitative definition of fever.. J Family Med Prim Care. 2022. DOI: 10.4103/jfmpc.jfmpc\_360\_22

*This document was generated from PubMed Central (PMC) full-text for Component 3, Iteration 5. Source: Open-access article 9810860. This is a complete full-text article from PMC.*