

Hazardous asteroids forecast via Markov random fields

Project for the exam: Probabilistic Modelling (DSE)

Marzio De Corato

Introduction

- **Final Goal** Assessment of forecasts and interpretability for different machine learning algorithms, including the probabilistic models
- **Method** Use a dataset for which the laws that interconnects the different features are known from general principles
- **Dataset** CNEOS asteroids dataset for more than 3500 asteroids
- **Theoretical laws** Celestial mechanics
- **Algorithms involved - probabilistic models** GLASSO, mgm, minforest, mmod
- **Algorithms involved - others** Random forest, Support Vector Machines, Quadratic Discriminant Analysis, Logistic Regression

Celestial mechanics

$$\mathbf{F}_1 = \mathcal{G} \cdot \frac{m_1 m_2}{r^3} \mathbf{r} = m_1 \ddot{\mathbf{r}}_1 \quad (1)$$

$$\mathbf{F}_2 = -\mathcal{G} \cdot \frac{m_1 m_2}{r^3} \mathbf{r} = m_1 \ddot{\mathbf{r}}_2 \quad (2)$$

If we consider the motion of the second item with respect to the first one

$$\ddot{\mathbf{r}} = \ddot{\mathbf{r}}_2 - \ddot{\mathbf{r}}_1 \quad \mu = \mathcal{G}(m_1 + m_2) \quad (3)$$

$$\frac{d^2 \mathbf{r}}{dt^2} + \mu \frac{\mathbf{r}}{r^3} = 0 \quad (4)$$

$$\mathbf{r} \times \ddot{\mathbf{r}} = 0 \implies \mathbf{r} \text{ and } \dot{\mathbf{r}} \text{ lies in the same plane}$$

Celestial mechanics

With polar coordinates $\hat{\mathbf{r}}$ and $\hat{\boldsymbol{\theta}}$

$$\mathbf{r} = r\hat{\mathbf{r}} \quad (5)$$

$$\dot{\mathbf{r}} = \dot{r}\hat{\mathbf{r}} + r\dot{\theta}\hat{\boldsymbol{\theta}} \quad (6)$$

$$\ddot{\mathbf{r}} = \left(\ddot{r} - r\dot{\theta}^2\right)\hat{\mathbf{r}} + \left[\frac{1}{r}\frac{d}{dt}\left(r^2\dot{\theta}\right)\right]\hat{\boldsymbol{\theta}} \quad (7)$$

$$\mathbf{h} = r^2\dot{\theta}\hat{\mathbf{z}} \quad (8)$$

$$h = r^2\dot{\theta} \quad (9)$$

Celestial mechanics [4]: 2th Kepler law




Figure 1: [4]

$$\delta A \approx \frac{1}{2}r(r + dr)\sin(\delta\theta) \approx \frac{1}{2}r^2\delta\theta \quad (10)$$

$$\frac{dA}{dt} = \frac{1}{2} r^2 \frac{d\theta}{dt} = \frac{1}{2} h \quad (11)$$

h is constant $\implies 2^{th}$ Kepler law

Celestial mechanics [4]: 1th Kepler law

Using the substitution $u = \frac{1}{r}$ $h = r^2 \dot{\theta}$

$$\dot{r} = -\frac{1}{u} \frac{du}{d\theta} \dot{\theta} = -h \frac{du}{d\theta} \quad (12)$$

$$\ddot{r} = -h \frac{d^2 u}{d\theta^2} \dot{\theta} = -h^2 u^2 \frac{d^2 u}{d\theta^2} \quad (13)$$

$$\frac{d^2 u}{d\theta^2} + u = \frac{\mu}{h^2} \quad (14)$$

$$u = \frac{\mu}{h^2} [1 + e \cos(\theta - \phi)] \quad (15)$$

Celestial mechanics [4]: 1th Kepler law

$$r = \frac{p}{1 + e \cos(\theta - \phi)} \quad (16)$$

e is **eccentricity**

- circle: $e = 0$ $p = a$
- ellipse: $0 < e < 1$
 $p = a(1 - e^2)$
- parabola: $e = 1$ $p = 2q$
- hyperbola: $e > 1$
 $p = a(e^2 - 1)$

a is the **semi-major axis** of the conic




Figure 2: [4]

Celestial mechanics [4]: 3th Kepler law




Figure 3: [4]

$$b^2 = a^2(1 - e^2) \quad (17)$$

$$r = \frac{a(1 - e^2)}{1 + e \cdot \cos(\theta - \phi)} \quad (18)$$

Area swept in one **orbital period** T

$$A = \pi ab$$

We know that: $hT/2 \quad h^2 = \mu a(1 - e^2)$

Therefore

$$T^2 = \frac{4\pi^2}{\mu} a^3 \quad (19)$$

3th Kepler law

$$\frac{m_c + m}{m_c + m'} = \left(\frac{a}{a'}\right) \left(\frac{T'}{T}\right)^2 \quad (20)$$

But since $m, m' \ll m_c$

$$(a')^3 \approx (T')^2 \quad (21)$$

And therefore

$$T' \approx a'^{3/2} \quad (22)$$

The mass of the asteroid is not involve

d

Orbital parameters

Mean motion

$$n = \frac{2\pi}{T} \quad (23)$$

$$v_{perihelion} = na\sqrt{\frac{1+e}{1-e}} \quad (24)$$

$$v_{aphelion} = na\sqrt{\frac{1-e}{1+e}} \quad (25)$$

Orbital parameters

Mean anomaly

$$M = n(t - \tau) \quad (26)$$

- $M = f = 0 \quad t = \tau$ Perihelion
- $M = f = \pi \quad t = \tau + T/2$ Aphelion




Figure 4: [4]

$$M = E - e \sin E \quad (27)$$

Jupiter Tisserard invariant

$$T_P = \frac{a_p}{a} + 2 \cos I \sqrt{\frac{a}{a_p} (1 - e^2)} \quad (28)$$

Orbital parameters




Figure 5: [4]

I : inclination of the orbit

Ω : longitude of the ascending node

Magnitude

$$\Phi = \frac{L}{4\pi r^2} \quad (29)$$

$$m = -2.5 \log_{10} \Phi + C \quad (30)$$

$$m_1 - m_2 = -2.5 \log_{10} \frac{\Phi_1}{\Phi_2} \quad (31)$$

$$M - m = -2.5 \log_{10} \frac{\Phi \cdot d^2}{\Phi \cdot 10^2} \quad (32)$$

$$M = m + 5 - 5 \log_{10} d \quad (33)$$

Where Φ is the flux for a sphere of radius r , m the relative magnitude and M the **Absolute magnitude**

Magnitude

$$\Phi = \frac{L}{4\pi r^2} \quad (34)$$

$$m = -2.5 \log_{10} \Phi + C \quad (35)$$

$$m_1 - m_2 = -2.5 \log_{10} \frac{\Phi_1}{\Phi_2} \quad (36)$$

$$M - m = -2.5 \log_{10} \frac{\Phi \cdot d^2}{\Phi \cdot 10^2} \quad (37)$$

$$M = m + 5 - 5 \log_{10} d \quad (38)$$

Where Φ is the flux for a sphere of radius r , m the relative magnitude and M the **Absolute magnitude**

Classification

Amors

Earth-approaching NEAs with orbits exterior to Earth's but interior to Mars' (named after asteroid (1221) Amor)



$$a > 1.0 \text{ AU} \\ 1.017 \text{ AU} < q < 1.3 \text{ AU}$$

Apollos

Earth-crossing NEAs with semi-major axes larger than Earth's (named after asteroid (1862) Apollo)



$$a > 1.0 \text{ AU} \\ q < 1.017 \text{ AU}$$

Atens

Earth-crossing NEAs with semi-major axes smaller than Earth's (named after asteroid (2062) Aten)



$$a < 1.0 \text{ AU} \\ Q > 0.983 \text{ AU}$$

Atiras

NEAs whose orbits are contained entirely within the orbit of the Earth (named after asteroid (163693) Atira)



$$a < 1.0 \text{ AU} \\ Q < 0.983 \text{ AU}$$

(q = perihelion distance, Q = aphelion distance, a = semi-major axis)

Classification

- **Potentially Hazardous Asteroids:** $\text{MOID} \leq 0.05 \text{ au}$
 $M \leq 22.0$ NEAs whose Minimum Orbit Intersection Distance (MOID) with the Earth is 0.05 au or less and whose absolute magnitude (M) is 22.0 or brighter

Classification


- The asteroid dataset was retrieved from Kaggle [2], which reports into a more machine readable form the dataset of The Center for Near-Earth Object Studies (CNEOS) [3], a NASA research centre.
- 3552 Asteroids
- Among the 40 the features, the ones connected only to the other name of the asteroid, or connected only to the name of the orbit and the one connected with the orbiting planet (since for all it was the Earth) were discarded
- The proportion hazardous/not hazardous was set 1:5
- The continuous measures were standardised and demeaned

Features


Features	Type
Neo Reference ID	not used
Absolute Magnitude	Continuous
Est Dia in KM (min)	Continuous
Est Dia in KM (max)	Continuous
Close Approach Date	Continuous
Epoch Date Close Approach	Continuous
Relative_Velocity	Continuous
Miss_Dist	Continuous
Min_Orbit_Intersection	Continuous
Jupiter_Tisserand_Invariant	Continuous
Epoch_Osculation	Continuous
Eccentricity	Continuous

Features


Features	Type
Semi Major Axis	Continuous
Inclination	Continuous
Asc Node Longitude	Continuous
Orbital Period	Continuous
Perihelion Distance	Continuous
Perihelion Arg	Continuous
Perihelion Time	Continuous
Mean_Anomaly	Continuous
Mean_Motion	Continuous
Hazardous	Categorical (Binary)




a) Perihelion Distance



b) Eccentricity






c) Min orbit intersection




d) Absolute magnitude

FAMD






GLASSO




$$\rho=0.2$$




$$\rho=0.2$$

GLASSO




$$\rho=0.3$$





$$\rho=0.4$$

Mixed interactions: mgm





Mixed interactions: minforest







Mixed interactions


The mgm model is the one that has the list of connection more coherent with the celestial mechanics laws.





Random Forest





Random Forest





Support Vector Machines



Quadratic Discriminant Analysis (QDA)



Logistic regression



ϕ coefficient

Table 1: ϕ coefficient (also known as Matthews correlation coefficient)

Algorithm	ϕ
RF	0.9876
SVM	0.7111
logistic	0.6173
mgm	0.5997
QDA	0.5562

forecast performances vs intepretability

- The mgm algorithm is not the best one in term of performances, but it provides the connections between the features. On the other side, except for the variable importance in RF, the other are black box one
- The mgm model, as the other graphical model is open to a true scientific validation, the other not.
- The probabilistic models lack in the forecast is definitely compensated by their interetability
- This is meaningful since this two features are in conflict
- The probabilistic models provide a good trade-off between intepretability and forecast performances, as long as one is interest to produce a really scientific result (e.g if the only aim is the forecast the RF is definitely better. However how long one can trust to the RF result ?)

References I

- [1] https://cneos.jpl.nasa.gov/about/neo_groups.html.
- [2] <https://www.kaggle.com/shrutimehta/nasa-asteroids-classification>.
- [3] <https://cneos.jpl.nasa.gov/>.
- [4] Carl D Murray and Stanley F Dermott. *Solar system dynamics*. Cambridge university press, 1999.