

- [17] H. Class et al. A benchmark study on problems related to CO₂ storage in geologic formations. *Comput. Geosci.*, 13(4):409–434, 2009.
- [18] P. J. Cook, A. Rigg, and J. Bradshaw. Putting it back where it came from: Is geological disposal of carbon dioxide an option for Australia. *Australian Petroleum Production and Exploration Association Journal*, 1, 2000.
- [19] G. T. Eigestad, H. K. Dahle, B. Hellevang, F. Riis, W. T. Johansen, and E. Øian. Geological modeling and simulation of CO₂ injection in the Johansen formation. *Computational Geosciences*, 13(4):435–450, 2009.
- [20] M. Flett, R. Gurton, and G. Weir. Heterogeneous saline formations for carbon dioxide disposal: Impact of varying heterogeneity on containment and trapping. *Journal of Petroleum Science and Engineering*, 57(1-2):106–118, 2007.
- [21] A. Förster, B. Norden, K. Zinck-Jørgensen, P. Frykman, J. Kulenkampff, E. Spangenberg, J. Erzinger, M. Zimmer, J. Kopp, G. Borm, et al. Baseline characterization of the CO₂SINK geological storage site at Ketzin, Germany. *Environmental Geosciences*, 13(3):145–161, 2006.
- [22] O. Gharbi, B. Bijeljic, E. Boek, and M. J. Blunt. Changes in pore structure and connectivity induced by CO₂ injection in carbonates: a combined pore-scale approach. In *Proceedings of the international conference on greenhouse gas technologies (GHGT)*, volume 11, 2012.
- [23] A. L. Goater, B. Bijeljic, and M. J. Blunt. Dipping open aquifers—the effect of top-surface topography and heterogeneity on CO₂ storage efficiency. *International Journal of Greenhouse Gas Control*, 17:318–331, 2013.
- [24] W. D. Gunter, S. Bachu, and S. Benson. The role of hydrogeological and geochemical trapping in sedimentary basins for secure geological storage of carbon dioxide. *Geological Society, London, Special Publications*, 233(1):129–145, 2004.
- [25] B. Hitchon. *Aquifer disposal of carbon dioxide: hydrodynamic and mineral trapping: proof of concept*. Geoscience Pub., 1996.
- [26] S. D. Hovorka, C. Doughty, S. M. Benson, K. Pruess, and P. R. Knox. The impact of geological heterogeneity on CO₂ storage in brine formations: a case study from the Texas Gulf Coast. *Geological Society, London, Special Publications*, 233(1):147–163, 2004.
- [27] J. A. Howell, A. Skorstad, A. MacDonald, A. Fordham, S. Flint, B. Fjellvoll, and T. Manzocchi. Sedimentological parameterization of shallow-marine reservoirs. *Petroleum Geoscience*, 14(1):17–34, 2008.
- [28] R. Juanes, E. J. Spiteri, F. M. Orr Jr, and M. J. Blunt. Impact of relative permeability hysteresis on geological CO₂ storage. *Water Resources Research*, 42:W12418, 2006.
- [29] T. Kempka, H. Class, U.-J. Görke, B. Norden, O. Kolditz, M. Kühn, L. Walter, W. Wang, and B. Zehner. A dynamic flow simulation code intercomparison based on the revised static model of the Ketzin pilot site. *Energy Procedia*, 40(0):418–427, 2013.
- [30] T. Kempka, M. Kühn, H. Class, P. Frykman, A. Kopp, C. Nielsen, and P. Probst. Modelling of {CO₂} arrival time at Ketzin—Part I. *International Journal of Greenhouse Gas Control*, 4(6):1007–1015, 2010.
- [31] A. Kopp, H. Class, and R. Helmig. Investigations on CO₂ storage capacity in saline aquifers: Part 1. dimensional analysis of flow processes and reservoir characteristics. *International Journal of Greenhouse Gas Control*, 3(3):263–276, 2009.
- [32] S. Krevor, R. Pini, L. Zuo, and S. M. Benson. Relative permeability and trapping of CO₂ and water in sandstone rocks at reservoir conditions. *Water Resources Research*, 48(2), 2012.
- [33] H. Law and S. Bachu. Hydrological and numerical analysis of CO₂ disposal in deep aquifer systems in the Alberta sedimentary basin. *Ener. Convers. Mgmt*, 37(6-8):1167–1174, 1996.