

- [49] E. Spiteri, R. Juanes, M. Blunt, and F. Orr. Relative-permeability hysteresis: Trapping models and application to geological CO₂ sequestration. In *SPE Annual Technical Conference and Exhibition*, 2005.
- [50] A. R. Syversveen, H. M. Nilsen, K.-A. Lie, J. Tveranger, and P. Abrahamsen. A study on how top-surface morphology influences the storage capacity of CO₂ in saline aquifers. In P. Abrahamsen, R. Hauge, and O. Kolbjørnsen, editors, *Geostatistics Oslo 2012*, volume 17 of *Quantitative Geology and Geostatistics*, pages 481–492. Springer Netherlands, 2012.
- [51] L. Van der Meer. The CO₂ storage efficiency of aquifers. *Energy Conversion and Management*, 36(6):513–518, 1995.
- [52] L. Wei and F. Saaf. Estimate co2 storage capacity of the johansen formation: numerical investigations beyond the benchmarking exercise. *Computational Geosciences*, 13(4):451–467, 2009.
- [53] L. Zuo, S. Krevor, R. W. Falta, and S. M. Benson. An experimental study of CO₂ exsolution and relative permeability measurements during CO₂ saturated water depressurization. *Transport in porous media*, 91(2):459–478, 2012.