

$$\notag=\frac{8\pi G}{c^4}T^{\mu\nu}/\log x$$

$$\begin{array}{l} {}^t\mathop{\mathrm{f}\!\!\!\mathrm{f}\!\!\!\mathrm{f}}\nolimits\mathrm{cohomD}_\chi[\mathrm{I_m}]\\ =\oint (px^n+qx+r)^{\nabla l}\end{array}$$

$$\frac{d}{dl}L(x,y)=2\int ||\sin 2x||^2d\tau$$

$$\frac{d}{d\gamma}\Gamma$$

$$||ds^2||=\lim_{x\rightarrow\infty}[\delta(x)\int\int\int\pi\left(\sum_{k=0}^{\infty}\frac{{}^n\sqrt{p},x}{n}\right)^{\frac{1}{2}}d\tau]^{\mu\nu}$$

$$\oint\!\!\!\!\!\oint\cong ||ds^2||=\lim_{x\rightarrow\infty}[\delta(x)\int\int\int\pi\left(\sum_{k=0}^{\infty}\frac{{}^n\sqrt{p},x}{n}\right)^{\frac{1}{2}}d\tau]^{\mu\nu}$$

$$e^{-2\pi T||\psi||}[\eta+\bar{h}]dx^\mu dx^\nu+T^2d^2\psi$$

$$\frac{-16\pi G}{c^4}T^{\mu\nu}/\log x=\notag$$

$$\frac{-16\pi G}{c^4}T^{\mu\nu}/e^{-2\pi T||\psi||}$$

$$=4\pi G\rho$$

$$\frac{\partial}{\partial x\partial y}=\nabla_i\nabla_j$$

$$\square\int\!\!\!\int\!\!\!\int={^t\mathop{\mathrm{f}\!\!\!\mathrm{f}\!\!\!\mathrm{f}}\nolimits}$$

$$\frac{\partial}{\partial x}\int\!\!\!\int\!\!\!\int=\nabla_i\nabla_j\int\nabla f(x)d\eta$$

$$\frac{\partial}{\partial f} \iiint = \square \iiint$$

$$\mathbb{T}|\Gamma, \mathbb{B}|B$$

$\mathfrak{E}|E, \mathfrak{C}|C$

$\mathbb{T}|F, \mathbb{A}|\beta$

$$\mathfrak{A}|_D, {}^t\mathcal{F} \cong \bigoplus_D \bigoplus^L$$

$$\square + \cancel{\square} = \emptyset$$

$$\cancel{\square} \mid \emptyset = \square$$

$$\begin{pmatrix} 1 & 0 \\ 0 & -i \end{pmatrix}^{\frac{1}{2}} \left| \begin{pmatrix} 1 & 0 \\ 0 & -i \end{pmatrix} \right.$$

$$x^{\frac{1}{2}} \cong x, \emptyset^{\frac{1}{2}} = \emptyset$$