

Miss mass consist with other dimension,
general relativity theory emerge with this value,
and zeta function contrast with Higgs field and
Miss mass material energy.

Masaaki Yamaguchi

The equation of gravity have with a value in mass of material essence, this component of system belong with being instance of accessority, this value mention to have with miss understand of value oneself is impossible with function anbrade with assembly of the other dimension consist with another patters, this partners be able to open the door, this one asist with be scan of AdS5 manifold and three manifold of norm, and this norm relate with general relativity theory of being have with miss value of function involved with accessority of sun and marcurey of planet near with mass of field curved from graivty of surface on this function system.

For instance,

$$a = \frac{4GM}{c^2 R} = \theta[rad], \text{ this one surranging of } 2\pi[rad], l = r\theta, l = 2\pi r = 2\pi r^2 \theta 2\pi[rad], l = r\theta, l = 2\pi r = 2\pi r^2 \theta$$

$$r^2 \theta = \frac{1}{2\pi} l$$

$$\text{sun of harf rounding of pi} = 60 \times 60 \times 360[s/\theta]$$

$$\left(\frac{\frac{a=8.47 \times 10^{-6}}{2}}{3.14}\right) \times 60 \times 60 \times 360 = 1.75[s/\theta]$$

$$||ds^2|| = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

This point result consist with frobenius theorem, assure and volume have with surface of graivty in norm, this resolve with point after converter of accessority from movement of theorem.

$$cR = 8\pi G \left(\frac{a}{c^2 R} \cdot \frac{V}{S} \right)$$

cR is world line of surface with miss value of function.

$$\begin{aligned} \frac{\partial}{\partial u^k} \Gamma_{ij}^i - \frac{\partial}{\partial u^i} \Gamma_{kj}^i + \Gamma_{km}^i \Gamma_{lj}^m - \Gamma_{lm}^i \Gamma_{kj}^m &= \Gamma_{lkj}^{'i} - \Gamma_{klj}^{'i} \\ \frac{d}{df} F &= \frac{d}{d\gamma} \Gamma = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \geq \frac{1}{2} i + \frac{1}{2} \\ \pi(\chi, x) &= [i\pi(\chi, x), f(x)] \end{aligned}$$

$$\begin{aligned}
&= \int \frac{1}{(x \log x)} dx = i \int x \log x dx - \int \frac{1}{(x \log x)} dx \\
&\int \Gamma(\gamma)' dx_m = \int \Gamma dx_m + \frac{d}{d\gamma} \Gamma \geq \int \Gamma dx_m \cdot \frac{d}{d\gamma} \Gamma \\
&= e^{-x \log x} + e^{x \log x} \geq e^{x \log x} - e^{-x \log x}
\end{aligned}$$

These equation consolemet with result of Fubni theorem,

$$\int \Gamma dx_m + \frac{d}{d\gamma} \Gamma \geq \int \Gamma dx_m \cdot \frac{d}{d\gamma} \Gamma = \frac{\partial}{\partial u^k} \Gamma_{ij}^i - \frac{\partial}{\partial u^l} \Gamma_{kj}^i + \Gamma_{km}^i \Gamma_{lj}^m - \Gamma_{lm}^i \Gamma_{kj}^m = \Gamma_{lkj}^{'i} - \Gamma_{klj}^{'i}$$

Non commutative theorem mention to resolve with Lie algebra equation, and this equation escort with general relativity and global partial integrality theorem, and this theorem hit from ideas are Thurston Perelman conjugater theorem, this theorem mention to result from Heisuke Hironaka theorem of delusion and visuality of Perelman theorems. This manifold deconstructed into all of seed dimension journey systems.

$$\begin{aligned}
T_{AB} &= - \int_a^b dt = - \int_a^b \frac{1}{c(1 - \frac{r_s}{r})} dr = - \frac{1}{c} \int_a^b \frac{r}{r - r_s} dr = - \frac{1}{c} \int_a^b (1 + \frac{r_s}{r - r_s}) dr \\
&\quad - \frac{1}{c} [r + r_s \log(r - r_s)]_a^b \\
&\quad \frac{1}{c} (a - b) + r_s \log \left(\frac{a - r_s}{b - r_s} \right) \\
&= \int C dx_m = \int \kappa T^{\mu\nu} dx_m = T^{\mu\nu} T^{\mu\nu'} \\
\int T_{AB} dx_m &= \left(\frac{1}{c} (a - b) + r_s \log \left(\frac{a - r_s}{b - r_s} \right) \right) \left(\frac{1}{c} (a - b) + r_s \log \left(\frac{a - r_s}{b - r_s} \right) \right)' \\
\frac{2\pi}{1 - \epsilon} &= 2\pi(1 + \epsilon) = 2\pi + 2\pi\epsilon \\
&= e^{x \log x} + e^{-x \log x} \geq e^{x \log x} - e^{-x \log x} \\
&(\int b dx_m)^b, B^{B'} = B^b \\
\frac{1}{1 - \epsilon} &= 1 + \epsilon \\
l &= \frac{r(0) + r(\pi)}{2} \\
&= \left(\frac{a}{1 + \epsilon} + \frac{a}{1 - \epsilon} \right) + 2 \\
&= \frac{a}{1 - e^x} = \frac{c^2}{GM(1 - e^x)} \\
= \frac{d}{df} F &= \frac{d}{d\gamma} \Gamma = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \geq \frac{1}{2} i + \frac{1}{2} \\
\beta(p, q) &= - \int \frac{1}{t^2} dt, \frac{d}{dt} g_{ij}(t) = -2R_{ij}
\end{aligned}$$